

Multiplier systems for Hermitian modular groups

Eberhard Freitag, Universität Heidelberg, Mathematisches Institut
2020

Abstract

Let $\Gamma_{F,n}$ be the Hermitian modular group of degree $n > 1$ in sense of Hel Braun with respect to an imaginary quadratic field F . Let r be a natural number. There exists a multiplier system of weight $1/r$ (equivalently a Hermitian modular form of weight $k + 1/r$, k integral) on some congruence group if and only if $r = 1$ or 2 . This follows from a much more general construction of Deligne [De] combining it with results of Hill [Hi], Prasad [P] and Prasad-Rapinchuk [PR]. As far as we know, the systems of weight $1/2$ have not yet been described explicitly. Remarkably Haowu Wang [Wa] gave an example of a modular form of half integral weight. Actually he constructs a Borcherds product of weight $23/2$ for a group of type $O(2, 4)$. This group is isogenous to the group $U(2, 2)$ that contains the Hermitian modular groups of degree two.

In this paper we want to study such multiplier systems. If one restricts them to the unimodular group

$$\mathcal{U} = \left\{ U; \begin{pmatrix} \bar{U}'^{-1} & 0 \\ 0 & U \end{pmatrix} \in \Gamma \right\},$$

one obtains a usual character. Our main result states that the kernel of this character is a non-congruence subgroup. For sufficiently small Γ it coincides with the group described by Kubota [Ku] in the case $n = 2$ and by Bass Milnor Serre [BMS] in the case $n > 2$.

Introduction

In the paper [FH] we gave a simple proof of a special case of a theorem of Deligne [De1] that states that the weights of multiplier systems on subgroups of finite index of the Siegel modular group of degree $n > 1$ are integral or half integral. The same result holds for other modular groups as for the Hermitian modular groups, the quaternary modular groups and the orthogonal groups $O(2, n)$, $n \geq 3$. In all these cases multiplier systems of half integral weight do exist. This follows from results of Deligne [De2], Prasad and Rapinchuk [PR] and Prasad [P]. In the case of the Siegel modular forms these multipliers are obtained as theta multiplier systems which can be expressed by means of a

symplectic Gauss sum. In other cases no such explicit description is known as far as I know. In this paper we study the multiplier systems in the Hermitian modular case. Our main result is that the restriction to the unimodular group is a character whose kernel is a non-congruence subgroup as it has been described in [Ku], [BMS].

Our proof rests on techniques from the paper [FH]. Part of this paper can be generalized from the Siegel to the Hermitian modular group. In the first part we generalize results from [FH] to the Hermitian case. Instead of proofs we refer to the corresponding result in [FH] if the generalization is straightforward.

The main difference of the two cases is that the pair (R, \mathfrak{q}) for the ring R of integers of an imaginary quadratic field with an distinguished non-zero ideal \mathfrak{q} admits a non trivial Mennicke symbol (Definition 5.1) whereas every Mennicke symbol in the case $R = \mathbb{Z}$ is trivial.

We fix a natural number n (which later will be 2). We denote by $E = E^{(n)}$ the $n \times n$ -unit matrix and by

$$I = I^{(n)} = \begin{pmatrix} 0 & -E \\ E & 0 \end{pmatrix}$$

the standard alternating matrix. The unitary group $U(n, n)$ consists of all $M \in \mathrm{GL}(2n, \mathbb{C})$ with the property $\bar{M}'IM = I$. (This is equivalent to $\bar{M}'HN = H$, where H is the Hermitian form iI of signature (n, n)). The special unitary group $SU(n, n)$ is the subgroup of elements with determinant one. One has $SU(1, 1) = \mathrm{SL}(2, \mathbb{R})$.

From now on we fix an imaginary quadratic field $F = \mathbb{Q}(\sqrt{d})$ of discriminant $d < 0$ and denote by

$$\mathfrak{o} = \mathbb{Z} + \mathbb{Z}\omega, \quad \omega = \frac{d + \sqrt{d}}{2},$$

its ring of integers. The Hermitian modular group $\Gamma_{F,n}$ is the subgroup of $U(n, n)$ of matrices with entries in \mathfrak{o} . Let $\mathfrak{q} \subset \mathfrak{o}$ be a non zero ideal. Then

$$\Gamma_{F,n}[\mathfrak{q}] = \mathrm{kernel}(\Gamma_{F,n} \longrightarrow \mathrm{GL}(2n, \mathfrak{o}/\mathfrak{q}))$$

is the (principal) congruence subgroup of level \mathfrak{q} . Since the field F is fixed, we can omit the label F and write

$$\Gamma_n := \Gamma_{F,n} \quad \text{and} \quad \Gamma_n[\mathfrak{q}] = \Gamma_{F,n}[\mathfrak{q}].$$

For sufficiently small \mathfrak{q} the group $\Gamma_n[\mathfrak{q}]$ is contained in $\mathrm{SL}(2n, \mathfrak{o})$. Then the group $\Gamma_1[\mathfrak{q}]$ is the usual principal congruence subgroup of the elliptic modular group $\mathrm{SL}(2, \mathbb{Z})$ of level $\mathfrak{q} \cap \mathbb{Z}$.

1. Multiplier systems

We consider the usual action $MZ = (AZ + B)(CZ + D)^{-1}$ of the unitary group $U(n, n)$ on the Hermitian upper half plane

$$\mathcal{H}_n = \{Z \in \mathbb{C}^{n \times n}; \quad Z = X + iY, \quad X = \bar{X}', \quad Y = \bar{Y}' > 0 \text{ (positive definite)}\}.$$

This is an open convex domain in $\mathbb{C}^{n \times n}$. The function

$$J(M, Z) = \det(CZ + D)$$

has no zeros on the half plane. Since the half plane is convex, there exists a continuous choice $L(M, Z) = \arg J(M, Z)$ of the argument. We normalize it such that it is the principal value for $Z = iE$ where E denotes the unit matrix. Recall that the principal value $\text{Arg}(a)$ is defined such that it is in the interval $(-\pi, \pi]$. So we have

$$L(M, iE) = \text{Arg}(J(M, i)) \in (-\pi, \pi].$$

We consider

$$w(M, N) := \frac{1}{2\pi} ((L(MN, Z) - L(M, NZ) - L(N, Z))).$$

Obviously,

$$e^{2\pi i w(M, N)} = 1.$$

Hence $w(M, N)$ is independent of Z and $w(M, N) \in \mathbb{Z}$. Usually we will compute $w(M, N)$ by evaluation at $Z = iE$.

$$w(M, N) = \frac{1}{2\pi} (\text{Arg } J(MN, iE) - \arg J(M, N(iE)) - \text{Arg } J(N, iE)),$$

where $\arg J(M, N(iE))$ is obtained through continuous continuation of the principal value $\text{Arg } J(M, iE)$ along a path from iE to $N(iE)$. Usually one takes the straight line.

1.1 Remark. *The function $w : U(n, n) \times U(n, n) \rightarrow \mathbb{Z}$ is a cocycle in the following sense:*

$$\begin{aligned} w(M_1 M_2, M_3) + w(M_1, M_2) &= w(M_1, M_2 M_3) + w(M_2, M_3), \\ w(E, M) &= w(M, E) = 0. \end{aligned}$$

The computation of $w(M, N)$ in degree 1 is easy for the following reason. From the definition we have

$$2\pi w(M, N) = \text{Arg}((c\alpha + d\gamma)i + c\beta + d\gamma) - \arg(cN(i) + d) - \text{Arg}(\gamma i + \delta)$$

for

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad N = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$$

where $\arg(cN(i) + d)$ is obtained from the principal value $\text{Arg}(ci + d)$ through continuous continuation. But $cz + d$ for z in the upper half plane never crosses the real axis. Hence the result of the continuation is the principal value too. So all three arguments in the definition of $w(M, N)$ are the principal values (in degree 1). This makes it easy to compute w . We rely on tables for the values of w which have been derived by Petersson and reproduced by Maass [Ma], Theorem 16.

1.2 Lemma. *Let $M = \begin{pmatrix} * & * \\ m_1 & m_2 \end{pmatrix}$, $S = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ be two real matrices with determinant 1 and (m'_1, m'_2) the second row of the matrix MS . Then*

$$4w(M, S) = \begin{cases} \text{sgn } c + \text{sgn } m_1 - \text{sgn } m'_1 - \text{sgn}(m_1 c m'_1) & \text{if } m_1 c m'_1 \neq 0, \\ -(1 - \text{sgn } c)(1 - \text{sgn } m_1) & \text{if } c m_1 \neq 0, m'_1 = 0, \\ (1 + \text{sgn } c)(1 - \text{sgn } m_2) & \text{if } c m'_1 \neq 0, m_1 = 0, \\ (1 - \text{sgn } a)(1 + \text{sgn } m_1) & \text{if } m_1 m'_1 \neq 0, c = 0, \\ (1 - \text{sgn } a)(1 - \text{sgn } m_2) & \text{if } c = m_1 = m'_1 = 0. \end{cases}$$

Corollary. *Assume that $m_1 c m'_1 \neq 0$ and that $m_1 m'_1 > 0$ or $m_1 c < 0$. Then $w(M, S) = 0$.*

We give an example.

1.3 Lemma. *We have*

$$w\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}\right) = w\left(\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) = 0.$$

2. Some special values of the cocycle

We give some examples for values of w in degree $n > 1$.

2.1 Lemma. *One has*

$$w\left(\begin{pmatrix} E & S \\ 0 & E \end{pmatrix}, M\right) = 0.$$

The proof is trivial. □

2.2 Lemma. *Let*

$$P = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Set

$$z := \det(iC + D).$$

Then

$$w(P, M) = w(M, P) = \text{Arg}(-z) - \text{Arg}(z) - \text{Arg}(-1).$$

Proof. Compare [FH], Lemma 1.2. □

2.3 Definition. *The **Siegel parabolic group** consists of all elements from $\text{SU}(n, n)$ of the form*

$$\begin{pmatrix} A & B \\ 0 & D \end{pmatrix}.$$

There is a character on the Siegel parabolic group

$$\varepsilon \begin{pmatrix} A & B \\ 0 & D \end{pmatrix} = \det(D).$$

For an element M of the Siegel parabolic group, the expression $\det(CZ + D) = \det(D)$ is independent of Z . Hence

$$L(M, Z) = 0 \quad \text{if } \varepsilon(M) = 1.$$

An immediate consequence is the following lemma.

2.4 Lemma. *For two elements P, Q of the Siegel parabolic group we have $w(P, Q) = 0$ if $\varepsilon(P) = 1$.*

The proof is trivial. □

We have to consider two embeddings $\iota_1, \iota_2 : \text{SL}(2, \mathbb{R}) \rightarrow \text{SU}(2, 2)$, namely

$$\iota_1 \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a & 0 & b & 0 \\ 0 & 1 & 0 & 0 \\ c & 0 & d & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \iota_2 \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & a & 0 & b \\ 0 & 0 & 1 & 0 \\ 0 & c & 0 & d \end{pmatrix}.$$

2.5 Lemma. *Let M be in the image of one of the embeddings ι_ν and N a Siegel parabolic matrix with $\varepsilon(N) = 1$. Then $w(M, N) = 0$.*

Proof. Compare [FH], Lemma 3.1.

2.6 Lemma. *Assume $n = 2$. Let*

$$M = \begin{pmatrix} E & S \\ 0 & E \end{pmatrix}, \quad S = \bar{S}'.$$

Then

$$w(I, M) = \begin{cases} 0 & \text{if } \operatorname{tr}(S) \geq 0, \\ -1 & \text{else.} \end{cases}$$

Proof. Compare [FH], Lemma 1.6. □

2.7 Lemma. *Assume $n = 2$. Let*

$$M = \begin{pmatrix} E & 0 \\ S & E \end{pmatrix}, \quad S = \bar{S}'.$$

Then

$$w(M, I) = \begin{cases} -1 & \text{if } \operatorname{tr}(S) > 0, \\ 0 & \text{else.} \end{cases}$$

Proof. Compare [FH], Lemma 1.7. □

3. Multipliers

3.1 Definition. *Let $\Gamma \subset \mathrm{U}(n, n)$ be an arbitrary subgroup and let r be a real number. A system $v(M)$, $M \in \Gamma$, of complex numbers of absolute value 1 is called a **multiplier system** of weight r if*

$$v(MN) \equiv v(M)v(N)\sigma(M, N)$$

where

$$\sigma(M, N) = \sigma_r(M, N) := e^{2\pi i r w(M, N)}.$$

Let now Γ be a normal subgroup of finite index of Γ_n , $n \geq 2$. Since the congruence subgroup property holds we know that Γ contains a congruence subgroup $\Gamma_n[\mathfrak{q}]$. It is easy to show that weights r of multiplier systems are rational [Ch]. Hence a suitable power of v is trivial on some congruence subgroup. This shows that there exists a natural number l such the all values of v are l th roots of unity.

For any $L \in \Gamma_n$ we can consider a conjugate multiplier system on Γ that is defined by

$$\tilde{v}(M) = v(LML^{-1}) \frac{\sigma(LML^{-1}, L)}{\sigma(L, M)}.$$

It is easy to check that this is a multiplier system and that this defines an action of Γ_n on the set of all multiplier systems on Γ . The quotient of two multiplier systems of the same weight is a homomorphism, as we know into a finite group. Since the congruence subgroup problem has been solved for the Hermitian modular group, we obtain $\tilde{v}(M) = v(M)$ on some congruence subgroup. Since the Hermitian modular group is finitely generated, they agree on $\Gamma_n[\mathfrak{q}]$, \mathfrak{q} suitable.

3.2 Lemma. *Let v be a multiplier system on a subgroup $\Gamma \subset \Gamma_n$ of finite index. In the case $n \geq 2$ there exists an ideal $\mathfrak{q} \neq 0$ such that $\Gamma[\mathfrak{q}] \subset \Gamma$ and such that*

$$v(M) = v(LML^{-1}) \frac{\sigma(LML^{-1}, L)}{\sigma(L, M)} \quad (M \in \Gamma[\mathfrak{q}])$$

for all $L \in \Gamma_n$.

Several times we will replace \mathfrak{q} by a smaller ideal. We then just say “for suitable \mathfrak{q} ”. We always assume that suitable \mathfrak{q} have the property that the only unit of \mathfrak{o} that is congruent to 1 mod \mathfrak{q} is 1. Then each Siegel parabolic $M \in \Gamma_n[\mathfrak{q}]$ has the property $\varepsilon(M) = 1$. We also assume that $\mathfrak{q} \subset 4\mathfrak{o}$.

3.3 Proposition. *Let v be a multiplier system on a subgroup $\Gamma \subset \Gamma_2$ of finite index. For suitable \mathfrak{q} the group $\Gamma_2[\mathfrak{q}]$ is contained in Γ and for each matrix M from $\Gamma_2[\mathfrak{q}]$ of the form*

$$M = \begin{pmatrix} E & 0 \\ * & E \end{pmatrix}.$$

we have $v(M) = 1$.

Proof. Compare [FH], Proposition 2.4. □

3.4 Proposition. *Let v be a multiplier system on a subgroup $\Gamma \subset \Gamma_2$ of finite index. For suitable \mathfrak{q} we have $\Gamma_2[\mathfrak{q}] \subset \Gamma$ and such the following holds. Let U be an element from the subgroup that is generated by the matrices $\begin{pmatrix} 1 & \mathfrak{q} \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} 1 & 0 \\ \mathfrak{q} & 1 \end{pmatrix}$ for $\mathfrak{q} \in \mathfrak{q}$ and let*

$$M = \begin{pmatrix} \bar{U}'^{-1} & * \\ 0 & U \end{pmatrix}.$$

Then $v(M) = 1$.

Proof. Compare [FH], Proposition 2.4. □

4. Embedded subgroups

We restrict now to the case $n = 2$. The case $n > 2$ can be derived from this easily. Besides the embeddings ι_1, ι_2 we have to consider the embedding

$$\iota : \mathrm{GL}(2, \mathbb{C}) \longrightarrow \mathrm{U}(2, 2), \quad \iota(U) = \begin{pmatrix} \bar{U}'^{-1} & 0 \\ 0 & U \end{pmatrix}.$$

This gives us an embedding $\mathrm{SL}(2, \mathfrak{o}) \hookrightarrow \Gamma_2$. We use the notation

$$\mathrm{SL}(2, \mathfrak{o})[\mathfrak{q}] = \ker(\mathrm{SL}(2, \mathfrak{o}) \longrightarrow \mathrm{SL}(2, \mathfrak{o}/\mathfrak{q})).$$

We have $w(\iota(U), \iota(V)) = 1$. Hence, for suitable \mathfrak{q}

$$\mathrm{SL}(2, \mathfrak{o})[\mathfrak{q}] \longrightarrow S^1, \quad U \longmapsto v(\iota(U)),$$

is a homomorphism. We mentioned that the values of v are l th roots of unity. Hence the kernel is a subgroup of finite index in $\mathrm{SL}(2, \mathfrak{o})[\mathfrak{q}]$.

Our method depends on some game between the embeddings ι_1, ι_2 and ι . We have

$$P\iota_1(M)P^{-1} = \iota_2(M), \quad P = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

From Lemma 2.2 follows $w(\iota_2(M), P) = w(P, \iota_1(M))$. Hence we obtain from Lemma 3.2 the following result.

4.1 Lemma. *Let v be a multiplier system on a subgroup $\Gamma \subset \Gamma_2$ of finite index. For suitable \mathfrak{q} we have $\Gamma_2[\mathfrak{q}] \subset \Gamma$ and*

$$v(\iota_1(M)) = v(\iota_2(M))$$

for $M \in \Gamma_1[\mathfrak{q}]$.

For sake of simplicity we write

$$v(M) = v(\iota_1(M)) = v(\iota_2(M)).$$

This is a multiplier system in degree 1. We have

$$w(M, N) = w(\iota_\nu(M), \iota_\nu(N)), \quad \text{for } \nu = 1, 2.$$

4.2 Lemma. *Let v be a multiplier system on a subgroup $\Gamma \subset \Gamma_2$ of finite index. For suitable \mathfrak{q} the value $v(M)$, $M \in \Gamma_1[\mathfrak{q}]$, depends only on the second row of M .*

Proof. Compare [FH], Lemma 3.2. □

4.3 Lemma. *Let v be a multiplier system on a subgroup $\Gamma \subset \Gamma_2$ of finite index. For suitable \mathfrak{q} we have $\Gamma_2[\mathfrak{q}] \subset \Gamma$ and for any*

$$M_1 \in \begin{pmatrix} a & b_1 \\ c_1 & d_1 \end{pmatrix} \in \mathrm{SL}(2, \mathfrak{o})[\mathfrak{q}] \quad \text{and} \quad M_2 = \begin{pmatrix} a & b_2 \\ c_2 & d_2 \end{pmatrix} \in \Gamma_1[\mathfrak{q}]$$

(in particular $a \in \mathbb{Z}$) the relation

$$v \begin{pmatrix} \bar{d}_1 & -\bar{c}_1 & 0 & 0 \\ -\bar{b}_1 & a & 0 & 0 \\ 0 & 0 & a & b_1 \\ 0 & 0 & c_1 & d_1 \end{pmatrix} \cdot v \begin{pmatrix} a & 0 & b_2 & 0 \\ 0 & 1 & 0 & 0 \\ c_2 & 0 & d_2 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = v \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & a & 0 & b_1 \bar{b}_1 b_2 \\ 0 & 0 & 1 & 0 \\ 0 & c_1 \bar{c}_1 c_2 & 0 & y \end{pmatrix}$$

holds (where $y = d_2 - b_2 c_2 (d_1 + \bar{d}_1) + a b_2 c_2 d_1 \bar{d}_1$).

Proof. Compare [FH], Lemma 3.3. \square

Now we assume that the multiplier system is of half integral weight. We can restrict it to a subgroup of finite index of the Siegel modular group. Since this group has the congruence subgroup property, v must agree with the theta multiplier system on a suitable congruence subgroup. Its restriction to the embedded $\Gamma_1[q]$ is the multiplier system of the classical theta function

$$1 + 2e^{2\pi iz} + 2e^{2\pi i4z} + 2e^{2\pi i9z} + \dots,$$

which, for $q \equiv 0 \pmod{4}$, is given by the Kronecker symbol $\left(\frac{c}{d}\right)$ (see [Di] for details). We will need it only for $c \neq 0$ and for odd d . We collect some properties (always assuming this condition)

$$\left(\frac{c_1 c_2}{d}\right) = \left(\frac{c_1}{d}\right) \left(\frac{c_2}{d}\right), \quad \left(\frac{c}{d_1 d_2}\right) = \left(\frac{c}{d_1}\right) \left(\frac{c}{d_2}\right).$$

Assume that m, n are odd coprime numbers such that at least one of them is not negative. Then the usual reciprocity law

$$\left(\frac{m}{n}\right) \left(\frac{n}{m}\right) = (-1)^{\frac{m-1}{2}} (-1)^{\frac{n-1}{2}}$$

holds. Assume $d > 0$ or $c_1 c_2 > 0$. Then

$$\left(\frac{c_1}{d}\right) = \left(\frac{c_2}{d}\right) \quad \text{if} \quad c_1 \equiv c_2 \pmod{d}.$$

Also the relation

$$\left(\frac{c}{d_1}\right) = \left(\frac{c}{d_2}\right) \quad \text{if} \quad \begin{cases} d_1 \equiv d_2 \pmod{4c} \text{ and } c \equiv 0 \pmod{4} \text{ or} \\ d_1 \equiv d_2 \pmod{c} \text{ and } c \equiv 2 \pmod{4} \end{cases}$$

is valid. Finally we mention

$$\left(\frac{c}{-1}\right) = \begin{cases} 1 & \text{for } c > 0, \\ -1 & \text{for } c < 0. \end{cases}$$

We obtain the existence of a natural number $q \equiv 0 \pmod{4}$ such that $\mathfrak{q} \subset (q)$ and such that

$$v(\iota_1 M) = v(\iota_2 M) = \left(\frac{c}{d}\right) \quad \text{for } M \in \Gamma_1[q].$$

Since $ad \equiv 1 \pmod{bc}$ and $b \equiv 0 \pmod{4}$, we get from one of the above rules

$$\left(\frac{c}{ad}\right) = 1 \quad \text{hence} \quad \left(\frac{c}{a}\right) = \left(\frac{c}{d}\right).$$

Next we assume that v is trivial on a congruence subgroup inside \mathcal{U} . So we can get $v(H_2) = 1$. From Lemma 4.3 we get the following result.

There exists a natural number $q \equiv 0 \pmod{4}$ such that for

$$\begin{aligned} a &\in \mathbb{Z}, \quad c_1 \in \mathfrak{o}, \quad c_2 \in \mathbb{Z}, \\ a &\equiv 1 \pmod{q}, \quad c_1 \equiv 0 \pmod{q}, \quad c_2 \equiv 0 \pmod{q}, \\ a\mathfrak{o} + c_1\mathfrak{o} &= \mathfrak{o}, \quad a\mathbb{Z} + c_2\mathbb{Z} = \mathbb{Z} \end{aligned}$$

the relation

$$\left(\frac{c_2}{a}\right) = \left(\frac{c_1 \bar{c}_1 c_2}{a}\right)$$

holds. This implies

$$\left(\frac{c\bar{c}}{a}\right) = 1 \quad \text{for } a \in 1 + q\mathbb{Z}, \quad c \in q\mathfrak{o}, \quad (a, c) = 1.$$

One can apply this relation to qc for an arbitrary $c \in \mathfrak{o}$ to obtain

$$\left(\frac{c\bar{c}}{a}\right) = 1 \quad \text{for } a \in 1 + q\mathbb{Z}, \quad c \in \mathfrak{o}, \quad (a, c) = 1.$$

It is known that there are infinitely many primes of the form $p = c\bar{c}$ [Co]. We choose one such that p and q are coprime. Then we have

$$\left(\frac{p}{a}\right) = \left(\frac{a}{p}\right)$$

if p and a are coprime. We choose α such that $\left(\frac{\alpha}{p}\right) = -1$. We can solve the congruence $1 + xq \equiv \alpha \pmod{p}$. Then $a = 1 + xq$ is coprime to p and we have

$$\left(\frac{c\bar{c}}{a}\right) = \left(\frac{p}{a}\right) = \left(\frac{a}{p}\right) = \left(\frac{\alpha}{p}\right) = -1.$$

This is a contradiction. This gives the first part of our main results (for $n = 2$ and as a consequence for arbitrary n).

4.4 Theorem. *Let $\Gamma \subset \Gamma_n$ be any subgroup of finite index of a Hermitian modular group of degree $n \geq 2$. Let v be a multiplier system of half integral weight. The restriction of v to the subgroup*

$$\mathcal{U} = \left\{ U; \begin{pmatrix} \bar{U}'^{-1} & 0 \\ 0 & U \end{pmatrix} \in \Gamma \right\}$$

is a usual character. Its kernel is a non-congruence subgroup of finite index. On a suitable congruence subgroup it agrees with the subgroup constructed in [Ku] in the case $n = 2$ and in [BMS] in the general case.

In the rest of the paper we will give the proof of the second part. (The case $n = 2$ is enough.)

5. Mennicke symbol

We recall the notion of a Mennicke symbol. Let R be a Dedekind domain and $\mathfrak{q} \subset R$ a non-zero ideal. We introduce the set

$$\mathcal{C}(R, \mathfrak{q}) := \{ (a, b) \in R \times R; \quad Ra + Rb = R, \quad a \equiv 1 \pmod{\mathfrak{q}}, \quad b \equiv 0 \pmod{\mathfrak{q}} \}.$$

Every pair (a, b) is the first column of a matrix $\begin{pmatrix} a & c \\ b & d \end{pmatrix} \in \mathrm{SL}(2, R)$. Multiplying M with a matrix of the type $\begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \in \mathrm{SL}(2, R)$ one can achieve $c \equiv 0 \pmod{\mathfrak{q}}$ and $d \equiv 1 \pmod{\mathfrak{q}}$.

5.1 Definition. *A Mennicke symbol mod \mathfrak{q} is a map*

$$\mathcal{C}(R, \mathfrak{q}) \longrightarrow G, \quad (a, b) \longmapsto \begin{bmatrix} b \\ a \end{bmatrix},$$

into some group G such that the following properties hold.

MS1 *It is invariant under the transformations $(a, b) \mapsto (a + xb, b)$ and $(a, b) \mapsto (a, b + qay)$ for integral x, y .*

MS2 *It satisfies the rule*

$$\begin{bmatrix} b_1 b_2 \\ a \end{bmatrix} = \begin{bmatrix} b_1 \\ a \end{bmatrix} \begin{bmatrix} b_2 \\ a \end{bmatrix}.$$

In our context, the group G will be the group of complex numbers of absolute value one. Mennicke symbols have been classified in [BSM] for Dedekind domains of arithmetic type. If R is the ring of algebraic integers in a number field that is not totally imaginary, then the Mennicke symbols are trivial. In the case of a totally imaginary field they can be described explicitly by means of power residue symbols.

The main result of this section is the following theorem.

5.2 Theorem. *Let v be a multiplier system of half integral weight on a subgroup of finite index of a Hermitian modular group of degree two. Then there exists a non-zero ideal $\mathfrak{q} \subset \mathfrak{o}$ with the following properties.*

1) $\Gamma_2[\mathfrak{q}] \subset \Gamma$.

2) *There exists a Mennicke symbol $[\cdot]$ for $(\mathfrak{o}, \mathfrak{q})$ such that for all $U \in \mathrm{SL}(2, \mathfrak{o})[\mathfrak{q}]$ one has*

$$\begin{bmatrix} c \\ a \end{bmatrix} = v \begin{pmatrix} \bar{U}'^{-1} & 0 \\ 0 & U \end{pmatrix}, \quad U = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

Proof. We mention that the kernel of v on $\mathrm{SL}(2, \mathfrak{o})[\mathfrak{q}]$ agrees for suitable \mathfrak{q} with a non congruence subgroup constructed by Kubota [Ku]. The proof of the theorem is given during the rest of this section.

We have to consider also the embeddings $\iota_1, \iota_2 : \Gamma_1[\mathfrak{q}] \longrightarrow \Gamma_2[\mathfrak{q}]$. As in the Hermitian case we have $v(i_1(M)) = v(i_2(M))$ and this depends only on the second row of $M \in \Gamma_1[\mathfrak{q}]$. Hence we can define

$$\begin{Bmatrix} c \\ d \end{Bmatrix} = v \begin{pmatrix} a & 0 & b & 0 \\ 0 & 1 & 0 & 0 \\ c & 0 & d & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}^{-1} = v \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & a & 0 & b \\ 0 & 0 & 1 & 0 \\ 0 & c & 0 & d \end{pmatrix}^{-1}.$$

The elements of $\mathcal{C}(\mathfrak{o}, \mathfrak{q})$ are the second columns of the matrices in $\mathrm{GL}(2, \mathfrak{o})[\mathfrak{q}]$. Hence

$$\begin{bmatrix} b \\ d \end{bmatrix} = \left(\iota \begin{pmatrix} a & b \\ c & d \end{pmatrix} \right)$$

is well-defined on $\mathcal{C}(\mathfrak{o}, \mathfrak{q})$. We claim that this symbol satisfies MS1. We notice that w is trivial on the image of ι . Hence v is a character on this group. The invariance under $(a, b) \mapsto (a, b + qay)$ follows from the equation

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & qy \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} a & b + qay \\ * & * \end{pmatrix}.$$

To prove the invariance under $(a, b) \mapsto (a + xb, b)$, we consider

$$\begin{pmatrix} 1 & 0 \\ -x & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix} = \begin{pmatrix} a + xb & b \\ * & * \end{pmatrix}.$$

Due to Lemma 3.2 we can assume that $v(\iota(M))$ is invariant under conjugation with $\iota \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix}$. This proves MS1.

We would like to have also MS2. To get a result in this direction, we make use of

$$v(\iota_\nu(M^{-1})) = v(\iota_\nu(M))^{-1}, \quad \nu = 1, 2.$$

This is true since in degree 1 one has $w(M, M^{-1}) = 0$. (This is a general rule for $c \neq 0$ and also for $c = 0$ and $a > 0$. But in our case $c = 0$ implies $a = 1$ since we assume $q > 2$.) From the analogue of Lemma 4.3 we get the general rule (compare Lemma 13.3 in [BMS].)

$$\begin{bmatrix} c_1 \\ a \end{bmatrix} \left\{ \begin{matrix} c_2 \\ a \end{matrix} \right\} = \left\{ \begin{matrix} c_1 \bar{c}_1 c_2 \\ a \end{matrix} \right\}.$$

We insert $c_2 = 1 - a$.

5.3 Lemma. *We have*

$$\left\{ \begin{matrix} 1 - a \\ a \end{matrix} \right\} = 1$$

for $a \equiv 1 \pmod{q}$.

Proof. We use

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ a-1 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 2-a & a-1 \\ 1-a & a \end{pmatrix}. \quad \square$$

Now we obtain

$$\begin{bmatrix} c \\ a \end{bmatrix} = \left\{ \begin{matrix} c\bar{c}(1-a) \\ a \end{matrix} \right\}.$$

Before we continue, we mention that $\{\cdot\}$ is not a Mennicke symbol. It does not satisfy MS1. Nevertheless it is closely related to $[\cdot]$.

5.4 Lemma. *We have*

$$\left\{ \begin{matrix} c \\ d \end{matrix} \right\} = \left\{ \begin{matrix} c \\ d + yc \end{matrix} \right\}$$

and

$$\left\{ \begin{matrix} c + xqd \\ d \end{matrix} \right\} = \left\{ \begin{matrix} c \\ d \end{matrix} \right\} e^{2\pi i r s} \quad \text{where } s = w\left(\begin{pmatrix} * & * \\ c & d \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ qx & 1 \end{pmatrix}\right).$$

Proof. The first relation can be derived from

$$\begin{pmatrix} 1 & -y \\ 0 & 1 \end{pmatrix} \begin{pmatrix} * & * \\ c & d \end{pmatrix} \begin{pmatrix} 1 & y \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} * & * \\ c & d + cy \end{pmatrix}.$$

To derive the second one we consider the relation

$$\begin{pmatrix} * & * \\ c & d \end{pmatrix} \begin{pmatrix} 1 & 0 \\ qx & 1 \end{pmatrix} = \begin{pmatrix} * & * \\ c + dxq & d \end{pmatrix}.$$

It shows

$$\left\{ \begin{matrix} c + dxq \\ b \end{matrix} \right\} = \left\{ \begin{matrix} c \\ d \end{matrix} \right\} e^{2\pi i r s}.$$

The w -value s is usually not zero. \square

But from the corollary of the table of Maass in the introduction we get

$$w\left(\begin{pmatrix} * & * \\ c\bar{c} & a \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ -qc\bar{c} & 1 \end{pmatrix}\right) = 0.$$

Using this, we get

$$\left\{ \begin{matrix} c\bar{c}(1-a) \\ a \end{matrix} \right\} = \left\{ \begin{matrix} c\bar{c} \\ a \end{matrix} \right\}.$$

So we obtain

$$\begin{bmatrix} c \\ a \end{bmatrix} = \left\{ \begin{matrix} c\bar{c} \\ a \end{matrix} \right\}$$

and moreover

$$\begin{bmatrix} c_1 c_2 \\ a \end{bmatrix} = \left\{ \begin{matrix} c_1 \bar{c}_1 c_2 \bar{c}_2 \\ a \end{matrix} \right\} = \begin{bmatrix} c_1 \\ a \end{bmatrix} \left\{ \begin{matrix} c_2 \bar{c}_2 \\ a \end{matrix} \right\} = \begin{bmatrix} c_1 \\ a \end{bmatrix} \begin{bmatrix} c_2 \\ a \end{bmatrix}.$$

This is part of the condition MS2. (We assume up to now $a \in \mathbb{Z}$).

5.5 Lemma. *Let (a, b) be two elements of \mathfrak{o} such that $(a, b) = \mathfrak{o}$. Then there exists $x \in \mathfrak{o}$ such that $a + xb$ is not divisible by any natural number > 1 .*

Proof. In the case $b = 0$ we can take $x = 0$. Hence we can assume that $b \neq 0$.

We write an element $a \in \mathfrak{o}$ in the form

$$a = \dot{a} + \ddot{a}\omega, \quad \omega = \frac{d + \sqrt{d}}{2}.$$

We will use

$$\omega^2 = -N(\omega) + d\omega.$$

From a solution $ax + by = 1$ we derive that the 4 integers

$$\dot{a}, \dot{b}, \ddot{a}N(\omega), \ddot{b}N(\omega)$$

are coprime. We have to find $x \in \mathfrak{o}$ such that

$$\dot{a} + \dot{x}\dot{b} - \ddot{x}\ddot{b}N(\omega), \quad \ddot{a} + \dot{x}\ddot{b} + \ddot{x}(\dot{b} + \ddot{b}d)$$

are coprime. We consider the greatest common divisors

$$g = \text{ggT}(\ddot{a}, \dot{b}, \ddot{b})$$

By Dirichlet's prime number theorem we can find $y \in \mathfrak{o}$ such that

$$\ddot{a} + \dot{y}\ddot{b} + \ddot{y}(\dot{b} + \ddot{b}d) = gp$$

where p is a prime number. There are infinitely many choices for p . Hence we can get that p is coprime to $\ddot{b}^2(d^2 - d)/4 + \dot{b}^2 + \dot{b}\ddot{b}d$. (This expression equals $N(\dot{b} + \ddot{b}\omega)$ which is positive.) Now we set

$$\dot{x} = \dot{y} + t(\dot{b} + \ddot{b}d), \quad \ddot{x} = \ddot{y} - t\ddot{b} \quad (t \in \mathbb{Z}).$$

Then we have still

$$\ddot{a} + \dot{x}\ddot{b} + \ddot{x}(\dot{b} + \ddot{b}d) = gp$$

but

$$\dot{a} + \dot{x}\dot{b} - \ddot{x}\ddot{b}N(\omega) = \dot{a} + \dot{y}\dot{b} - \ddot{y}\ddot{b}N(\omega) + t(\dot{b}^2 + \dot{b}\ddot{b}d + \ddot{b}^2N(\omega)).$$

Now we consider the greatest common divisor

$$g' = (\dot{a} + \dot{y}\dot{b} - \ddot{y}\ddot{b}N(\omega), \dot{b}^2 + \dot{b}\ddot{b}d + \ddot{b}^2N(\omega)).$$

We can choose t such that

$$\dot{a} + \dot{x}\dot{b} - \ddot{x}\ddot{b}N(\omega) = g'p'$$

where p' is a prime. We can choose p' coprime to gp . Our goal was to get gp and $g'p'$ coprime. This means that g, g' are coprime. But

$$\text{ggT}(g, g') = \text{ggT}(\ddot{a}, \dot{b}, \ddot{b}, \dot{a} + \dot{y}\dot{b} - \ddot{y}\ddot{b}N(\omega), \dot{b}^2 + \dot{b}\ddot{b}d + \ddot{b}^2N(\omega)) = \text{ggT}(\dot{a}, \dot{b}, \ddot{a}, \ddot{b}) = 1.$$

This proves Lemma 5.5. □

Now we come to the proof of MS2. We can write it in the form

$$\begin{bmatrix} q^2b_1b_2 \\ a \end{bmatrix} = \begin{bmatrix} qb_1 \\ a \end{bmatrix} \begin{bmatrix} qb_2 \\ a \end{bmatrix}$$

where a, b_1, b_2 are in \mathfrak{o} such that $a \equiv 1 \pmod{q}$ and such that $(a, b_1) = (a, b_2) = \mathfrak{o}$. This formula is invariant under the replacement $b_1 \mapsto b_1 + xa$. By Lemma 5.5 we can assume that b_1 is not divisible by any natural number. We also want to make an replacement for b_2 . For this we consider the ray class of the principal ideal (b_2) mod the ideal (a) . (Recall that two ideals $\mathfrak{b}_1, \mathfrak{b}_2$ are in the same ray class mod an ideal \mathfrak{a} if there exist $\beta_1 \equiv \beta_2 \equiv 1 \pmod{\mathfrak{a}}$ such that $\beta_1\mathfrak{b}_1 = \beta_2\mathfrak{b}_2$.) Our product formula does not change if one replaces b_2 by βb_2 for $\beta \equiv 1 \pmod{(a)}$. Hence we may replace (b_2) by any other (b'_2) in the same ray class. In each ray class there are infinitely many primes. Hence we can assume that b_2 is coprime to $N(b_1)$. Now we make the replacement $b_2 + xaN(b_1)$. Again we make use of Lemma 5.5 to reduce to the case where b_2 is not divisible by any natural number, and, in addition, is coprime to Nb_1 . We claim that then also b_1b_2 is not divisible by any natural number. We argue indirectly and assume that there is prime number p that divides b_1b_2 . Then p splits in \mathfrak{o} into two prime

ideals, $(p) = \bar{\mathfrak{p}}\mathfrak{p}$. Since p does not divide b_1 and b_2 , we can assume $\bar{\mathfrak{p}}|b_1$ and $\mathfrak{p}|b_2$. But then $\mathfrak{p}|\bar{b}_1$ which contradicts to the fact that \bar{b}_1 and b_2 are coprime.

Now we make the stronger assumption $a \equiv 1 \pmod{q^2}$. Then $\bar{a} \equiv 0 \pmod{q^2}$. We can make the replacement $a \mapsto a + xq^2b_1b_2$ without changing the product formula. This means that we replace $\bar{a} \mapsto q^2(\bar{a}/q^2 + \bar{y})$ where $y = xb_1b_2$. Since $b = b_1b_2$ is not divisible by any natural number, \bar{y} runs through all rational integers if x runs through all integers in \mathfrak{o} . This shows that can transform \bar{a} to 0. So we can assume $a \in \mathbb{Z}$. But in this case the product formula has been proved. Now we replace q by q^2 to obtain Theorem 5.2. \square

References

- [BMS] Bass, H., Milnor, J., Serre, J.P.: *Solution of the congruence subgroup problem for SL_n ($n \geq 3$) and Sp_{2n} ($n \geq 2$)*, Publications mathématiques l’I.H.É.S., tome **33**, p. 59–137 (1967)
- [Co] Cox, D.A.: *Primes of the Form $x^2 + ny^2$* , Wiley Series in Pure and Applied Mathematics (2013)
- [De1] Deligne, P.: *Extensions centrales non résiduellement finies de groupes arithmétiques*, C. R. Acad. Sci. Paris **287**, p. 203–208 (1978)
- [De2] Deligne, P.: *Extensions Centrales de groupes Algébriques Simplememt Connexes et Cohomology Galoisienne*, Publ. Math. I.H.E.S. **84**, 35–89 (1996)
- [Di] Dickson, L.E.: *Introduction to the Theory of Numbers*, Dover Publications, New York, Dover (1957).
- [FH] Freitag, E., Hauffe-Waschbüsch, A.: *Maultplier systems for Siegel modular groups*, arXiv:2009.06455 [math.NT] (2020)
- [Hi] Hill, R.: *Fractional weights and non-congruence subgroups*, Automorphic Forms and Representations of algebraic groups over local fields, Saito, H., Takahashi, T. (ed.) Surikenkoukyuroku series **1338**, 71–80 (2003)
- [Ku] Kubota, T.: *Ein arithmetischer Satz über eine Matrizengruppe*, J. reine angew. Math., **222**, 55–77 (1965)
- [Ma] Maass, H.: *Lectures on Modular Functions of One Complex Variable*, Notes by Sunder Lal, Tata Institute Of Fundamental Research, Bombay, Revised 1983 (1964)
- [Me] Mennicke, J.: *Zur Theorie der Siegelschen Modulgruppe*, Math. Annalen **159**, 115–129 (1965)
- [P] Prasad, P.: *Deligne’s topological central extension is universal*, Advances in Mathematics **181**, 160–164 (2004)
- [PR] Prasad, G., Rapinchuk, A.: *Computation of the metaplectic kernel*, Publications mathématiques de l’I.H.É.S. tome **84**, p.91–187 (1996)