

The construction of Green currents and singular theta lifts for unitary groups

Jens Funke, Eric Hofmann*

February 5, 2019

Contents

1. Introduction	2
2. The unitary group	6
2.1. The unitary symmetric space	6
2.2. The unitary Lie algebra	7
3. Schwartz forms	9
3.1. Weil representation	9
3.2. The Kudla-Millson form φ_{KM}	10
3.3. The Schwartz form ψ	11
4. A singular Schwartz form	13
4.1. The current equation	14
4.1.1. Local integrability	15
4.1.2. The current equation	17
5. The singular theta lift	18
5.1. Weak Maass forms	18
5.2. Regularized theta integral	20
5.3. Singularities and current equation	21
5.4. Adjointness to the Kudla-Millson lift	23
6. Comparison of the two Green currents	25
6.1. Green currents of Bruinier type	25
6.2. The Kudla type Green current as a theta lift	26

*Supported by DFG research fellowship HO 6123/1-1 in 2017/18

6.3. The difference of the two Green currents as a modular form	27
7. Poincaré series	28
7.1. An eigenvalue equation	28
7.2. Unfolding against the Poincaré series	30
A. Calculations in the Fock model	33
A.1. Proof of Proposition 3.2	33
A.2. Proof of Theorem 3.3	35
A.3. The auxiliary form $d^c\psi$	37
B. The Fock model for unitary dual pairs	37
B.1. The Fock model for the symplectic group	38
B.2. Unitary dual pairs	38

1. Introduction

Since its inception 20 years ago ([23]), the *Kudla program* has yielded many critical insights at the intersection of arithmetic geometry and automorphic forms. Roughly speaking, the Kudla program asserts the modularity of generating series of certain ‘special’ cycles in (integral models of) orthogonal and unitary Shimura varieties when viewed as elements in the arithmetic Chow group, in particular as derivatives of Eisenstein series. For an overview, see e.g. [25]. Note that this program can be viewed as the considerable refinement and extension of the work of Kudla and Millson in the 1980’s, see e.g. [28], which employed the theta correspondence and theta series to establish the modularity of the special cycles in the cohomology of locally symmetric spaces.

One important aspect in the Kudla program is the construction of appropriate Green currents for the complex points of the cycles which are then used to ‘upgrade’ the cycles to define elements in the arithmetic Chow group associated to the underlying Shimura variety.

For special (Heegner) divisors in Hermitian spaces to the orthogonal group $O(p, 2)$, Kudla [23, 24] constructed Green functions in terms of the logarithmic integral. Later Bruinier and the first named author of this paper employed Borchers’ singular theta lift [2] and its extension to harmonic weak Maass forms to construct another Green function for the divisors [4, 10, 6]. In [4] an adjointness result between the Borchers lift and the Kudla-Millson lift was established, and in this context also the relationship in terms of the construction of these two Green functions was clarified. We note that the underlying structure for all these constructions is the dual reductive pair $O(p, 2) \times \mathrm{SL}_2(\mathbb{R})$.

For the dual pair $U(p, 1) \times U(1, 1)$ the singular theta lift (for weakly holomorphic input) was first studied in detail by the second named author of this paper, see [19]. Its extension to harmonic weak Maass forms gives again Green functions for the special cycles which again are divisors. These have been utilized in [9, 7, 8].

The difference between the two Green functions and its consequences in the context of the Kudla program are fairly subtle. This was studied and clarified by Ehlen and

Sankaran [12]. They show in the case of $O(p, 2)$ that the difference of the generating series can be viewed as a smooth modular form of weight $\frac{p}{2} + 1$.

Bruinier [5] considered the situation in the Hilbert modular case. In this case, Bruinier manages to circumvent the problem of the non-existence of the harmonic weak Maass forms to define a singular theta lift for ‘Whittaker forms’ which then again gives rise to Green functions for the special divisors.

For cycles of higher codimension much less has been known until recently. In Kudla’s original work [23], Liu [29], and Bruinier and Yang [11] star products are used to construct Green forms for cycles of higher codimension for $O(p, 2)$ and $U(p, 1)$. In recent groundbreaking work, Garcia and Sankaran [17] employed Quillen’s theory of superconnections to construct Green forms in $O(p, 2)$ and $U(p, q)$ in any codimension.

In this paper, we consider the construction of Green currents, in fact Green forms, for the dual pair $U(p, q) \times U(1, 1)$. The associated Shimura varieties for $U(p, q)$ are very attractive and natural objects to study. Furthermore, the cycles in question are no longer divisors but have codimension q . On the other hand, as $SU(1, 1)$ is isomorphic to $SL_2(\mathbb{R})$ this case can be still approached via singular theta lifts of Borcherds type.

While this paper is certainly to a large degree written with applications in arithmetic geometry and the Kudla program in mind, we ignore this aspect in this paper completely and focus on the Archimedean side of the story.

Let V be an Hermitian space over an imaginary quadratic field of signature (p, q) . Then we can view the associated Hermitian domain \mathbb{D} as the Grassmannian of negative q -planes in $V(\mathbb{R})$. Let L be an even lattice in V and let Γ be a finite-index subgroup of the stabilizer of L in $U(V(\mathbb{R}))$ ¹. We then define $X = \Gamma \backslash \mathbb{D}$ which gives a quasi-projective variety of dimension pq .

To $x \in V$ with positive length we associate a subsymmetric space $\mathbb{D}(x) = \{z \in \mathbb{D}; z \perp x\}$. Let Γ_x be the stabilizer of x in Γ , and we define the cycle $Z(x)$ as the image of $\Gamma_x \backslash \mathbb{D}_x$ in X . Note that the cycles have codimension q in X and arise from a suitable embedding $U(p-1, q) \hookrightarrow U(p, q)$. Finally, for $m > 0$ we set

$$Z(m) = \sum_{\substack{x \in L, (x, x) = 2m \\ \text{mod } \Gamma}} Z(x) \in H^{q, q}(X)$$

We let $Z(0) = c_q$, the q -th Chern form on \mathbb{D} . Finally, we set $Z(m) = \emptyset$ for $m < 0$.

The starting point for our considerations is the Kudla-Millson Schwartz form

$$\varphi_{KM} \in [\mathcal{S}(V) \otimes \mathcal{A}^{q, q}(\mathbb{D})]^G,$$

which takes values in the closed differential (q, q) -forms in D . Under the action of the Weil representation of $SO(2) \subset SL_2(\mathbb{R}) \simeq SU(1, 1)$ it is an eigenfunction of weight

¹In the main text we allow for a coset condition and work in the context of vector-valued modular forms. The results of this paper of course also hold in an appropriate adelic setting.

$p + q$. Then the associated theta series $\theta(z, \tau, \varphi_{KM})$ to L ($\tau = u + iv \in \mathbb{H}$) is a (non-holomorphic) modular form of weight $p + q$ for a congruence subgroup $\Gamma' \subseteq \mathrm{SL}_2(\mathbb{Z})$ with values in the closed differential (q, q) -forms in X . Furthermore, in cohomology we have

$$[\theta(z, \tau, \varphi_{KM})] = \sum_{m \geq 0} [Z(m)] q^m. \quad (q = e^{2\pi i \tau})$$

The key observation for our construction is

Theorem 1.1. *There exists a Schwartz form*

$$\psi \in [\mathcal{S}(V) \otimes \mathcal{A}^{q-1, q-1}(\mathbb{D})]^G$$

such that

$$\omega(L) \varphi_{KM} = dd^c \psi. \quad (1.1)$$

Here $\omega(L)$ is the Weil-representation action of $L = \frac{1}{2} \begin{pmatrix} 1 & -i \\ -i & -1 \end{pmatrix} \in \mathfrak{sl}_2(\mathbb{C}) \simeq \mathfrak{su}(1, 1)(\mathbb{C})$ which corresponds to Maass lowering operator $L = \bar{L}_{p+q}$ for forms on the upper half plane and d and d^c are the standard exterior derivatives acting on $\mathcal{A}^\bullet(\mathbb{D})$. Furthermore, ψ has weight $p + q - 2$ under the action of $\mathrm{SO}(2)$.

Note that the solution to the equation $\omega(L)\varphi_{KM} = d\psi'$ was already constructed in [28], in fact, more generally for the dual pairs $\mathrm{O}(p, q) \times \mathrm{Sp}(n)$ and $\mathrm{U}(p, q) \times \mathrm{U}(n, n)$. In the same way our form ψ can be used to solve the higher rank equations for $\mathrm{U}(p, q) \times \mathrm{U}(n, n)$. We explicitly construct ψ and establish its properties using the Fock model of the Weil representation, see Appendix A. For convenience and future use we develop the formulas for the Weil representation much more generally for the dual pair $\mathrm{U}(p, q) \times \mathrm{U}(r, s)$.

We then define the *Green current of Kudla type* by setting

$$\Psi^0(x, z) := - \int_1^\infty \psi(\sqrt{t}x, z) e^{\pi t(x, x)} \frac{dt}{t}$$

for nonzero x and then for $m \in \mathbb{Q}$ and $w > 0$,

$$\Xi^K(m, w)(z) := \sum_{\substack{\lambda \in L, \lambda \neq 0 \\ (\lambda, \lambda) = 2m}} \Psi^0(\sqrt{w}\lambda, z) e^{-2\pi m w},$$

which defines a $(q - 1, q - 1)$ -form on X with singularities along the cycles $Z(m)$ for $m > 0$. For $m \leq 0$, the forms are smooth.

The principle of this construction and its properties were already outlined in [16] for $\mathrm{O}(p, q) \times \mathrm{SL}_2(\mathbb{R})$ for the form ψ' mentioned above and was also implicit in [6] for the Hermitian case $\mathrm{O}(p, 2)$. Garcia and Sankaran [17] also follow these lines but use superconnections to solve (1.1). We have not checked the details but it seems likely that for $n = 1$ their form ν is equal to our form ψ . Garcia and Sankaran then succeed to construct Green forms for $n > 1$ by realizing that an analogous integral will work.

On the other hand, we define a singular theta lift (of Borchers type) using the theta series $\theta(z, \tau, \psi)$ as integral kernel. Namely, for f , a harmonic Maass form of weight $k = 2 - p - q$, we set

$$\Phi(z, f) := \int_{\Gamma' \setminus \mathbb{H}}^{reg} f(z) \theta(z, \tau, \psi) d\mu(\tau).$$

Here the regularization follows the by now standard procedure introduced by Harvey and Moore [18] and Borchers [2]. We then define for $m > 0$ the *Green current of Bruinier type* by

$$\mathcal{G}^B(m)(z) := \Phi(z, F_m).$$

Here $F_m(\tau)$ denotes the Hejhal Poincaré series of weight k which has principal part q^{-m} and ‘shadow’ $\xi_k(F_m) = P_m$, the holomorphic Poincaré series for Γ' of index m and weight $2 - k = p + q$. Here $\xi_k = v^{k-2} \frac{\partial}{\partial \tau} = v^{k-2} \overline{L}_k$ is the differential operator mapping forms of weight k to weight $2 - k$. For $m \leq 0$, we set $\mathcal{G}^B(m)(z) = 0$. We show

Theorem 1.2. *The forms $\Xi^K(m, w)$ and $\mathcal{G}^B(m)$ both define Green currents for the cycle $Z(m)$. More precisely, as currents we have*

$$\begin{aligned} dd^c[\Xi^K(m, w)] + (-i)^q \delta_{Z(m)} &= [\varphi_{KM}(m, w) e^{2\pi m w}], \\ dd^c[\mathcal{G}^B(m)] + (-i)^q \delta_{Z(m)} &= [dd^c \Phi(F_m)]. \end{aligned}$$

Here $\varphi_{KM}(m, w) = \sum_{\lambda \in L, (\lambda, \lambda) = 2m} \varphi_{KM}(\sqrt{w}x)$.

The proof employs the same Lie-theoretic set-up as in [6] and [16] for the orthogonal case. We first consider the analogous question for $\Psi^0(x)$, and as a corollary we obtain the Green property for $\Xi^K(m, w)$. We then show that $\mathcal{G}^B(m)$ has the same singularities as $\Xi^K(m, w)$, hence yielding the same residue.

We can identify the term $dd^c \Phi(z, f)$ in the previous theorem explicitly as follows:

Theorem 1.3. *Let f be a harmonic weak Maass form for Γ' of weight $k = 2 - p - q$ with holomorphic constant term a_0^+ , and let $\xi_k(f)$ be its shadow, a cusp form of weight $p + q$. Then*

$$dd^c \Phi(z, f) = (\Theta(\cdot, z, \varphi_{KM}), \xi_k(f))_{p+q} + a^+(0, 0) c_q$$

as differential (q, q) -forms on X . Here $(\alpha, \beta)_\ell$ denotes the Petersson inner product in weight ℓ . In particular, $dd^c \Phi(z, f)$ extends to a smooth closed (q, q) -form of moderate growth and $dd^c \Phi(z, f) = a^+(0, 0) c_q$ for f weakly holomorphic.

This can be viewed as an adjointness result between the Kudla-Millson lift and the singular theta lift associated to ψ . It is the analogue of the main result in [6], and the proof follows along the same lines.

Following ideas of Ehlen and Sankaran [12] we then compare the two Green currents in a different way. We show

Theorem 1.4. *For each $z \in \mathbb{D}$, the generating series*

$$F(\tau) = -\log(v) \psi(0)(z) - \sum_{m \in \mathbb{Q}} (\Xi^K(m, v) - \mathcal{G}^B(m))(z) q^m$$

transforms like a smooth modular form of weight $p + q$. In addition, F is orthogonal to cusp forms and satisfies $L_{p+q}F(\tau) = -\theta(\tau, \psi)$.

Finally, we construct for $m > 0$ a different Green object $\mathcal{G}_s^B(m)(z)$ depending on a complex parameter s . It is given essentially² as $\Phi(F_m(s), z)$, where $F_m(\tau, s)$ is the Hejhal Poincaré series of weight k with complex parameter s (at $s = s_0 = 1 - k/2$ this is the weak Maass form F_m introduced above). We show

Theorem 1.5. *Let Δ be the Laplace operator acting on differential forms on X . Then*

$$\Delta \mathcal{G}_s^B(m) = ((2s - 1)^2 - (2s_0 - 1)^2) \mathcal{G}_s^B(m).$$

Furthermore, $\mathcal{G}^B(m, s)$ agrees (up to a multiplicative constant) with the Green current constructed for $m > 0$ by Oda and Tsuzuki [30].

In view of applications in the Kudla program but also in its own right it will among other things be interesting to consider suitable integrals of the singular theta lift $\Phi(z, f)$, say along the lines of [24] and [10], and also to compute the Fourier-Jacobi expansion of the singular theta lift $\Phi(z, f)$ and to analyze the growth at the boundary components at suitable toroidal compactifications of X . We hope to come back to these questions in the near future.

We thank Jan Bruinier, Stephan Ehlen, and Steve Kudla for very valuable discussions, suggestions, and help. Hofmann would like to thank the Department of Mathematical sciences at Durham University for its hospitality during the academic year 2017/18. His stay was supported by a research fellowship (*Forschungsstipendium*) of the DFG.

2. The unitary group

2.1. The unitary symmetric space

We let $(V, (\cdot, \cdot))$ be a complex vector space of dimension m with a non-degenerate Hermitian form (\cdot, \cdot) of signature (p, q) . We assume that (\cdot, \cdot) is \mathbb{C} -linear in the second and \mathbb{C} -antilinear in the first variable. We pick standard orthogonal basis elements v_α ($\alpha = 1, \dots, p$) and v_μ ($\mu = p + 1, \dots, m$) of length 1 and -1 respectively.³ We let z_α and z_μ be the corresponding coordinate functions so that

$$(x, x) = \sum_{\alpha=1}^p |z_\alpha|^2 - \sum_{\mu=p+1}^m |z_\mu|^2,$$

for $x = \sum_{\alpha} z_\alpha v_\alpha + \sum_{\mu} z_\mu v_\mu \in V$. The choice of basis also gives a decomposition $V = V_+ \oplus V_-$ into definite subspaces. We let $G = \mathrm{U}(V)$ be the unitary group of V ,

²Due to slightly different regularization process $\mathcal{G}_{s_0}^B(m)(z)$ differs from $\Phi(z, F_m)$ by a smooth form.

³Throughout the paper we will follow [28] and use ‘early’ Greek letters to indices ranging from 1 to p and ‘late’ for indices from $p + 1$ to m .

and let $\mathbb{D} = G/K$ be the associated symmetric space of complex dimension pq . Here $K \simeq U(p) \times U(q)$ is the maximal compact subgroup corresponding to the basis of V chosen above. We realize the symmetric space as the Grassmannian of negative definite q -planes in V :

$$\mathbb{D} \simeq \{z \subset V : \dim(z) = q, (x, x)|_z < 0\}.$$

For $z \in \mathbb{D}$ and a vector $x \in V$, we denote by x_z the orthogonal projection of x onto the subspace z , and define

$$R(x, z) := -(x_z, x_z).$$

Note that $R(x, z) \geq 0$ with $R(x, z) = 0$ if and only if $x \in z^\perp$. When x has positive norm, let $\mathbb{D}(x)$ denote the codimension q sub-Grassmannian

$$\mathbb{D}(x) := \{z \in \mathbb{D} : z \perp x\} = \{z \in \mathbb{D} : R(x, z) = 0\}.$$

Also, for convenience, if x is non-positive, set $\mathbb{D}(x) = \emptyset$.

Let $L \subset V$ be an even hermitian lattice, i.e. a projective module over the ring of integers $\mathcal{O}_{\mathbb{F}}$ of an imaginary quadratic number field \mathbb{F} , on which the restriction of (\cdot, \cdot) is $\mathcal{O}_{\mathbb{F}}$ -valued. We fix an embedding of \mathbb{F} into \mathbb{C} . Denote by $\mathcal{D}_{\mathbb{F}}^{-1}$ the inverse different ideal of \mathbb{F} . The dual lattice L^\sharp is given by

$$L^\sharp = \{x \in V : (x, \lambda) \in \mathcal{D}_{\mathbb{F}}^{-1}, \forall \lambda \in L\} = \{x \in V : \text{trace}_{\mathbb{F}/\mathbb{Q}}(x, \lambda) \in \mathbb{Z}, \forall \lambda \in L\}.$$

Note that $L \subset L^\sharp$. The quotient L^\sharp/L is called the discriminant group of L .

For $m \in \mathbb{Q}$ and $h \in L^\sharp/L$, we define the special cycle $\mathbb{D}(m, h)$ in \mathbb{D} by

$$\mathbb{D}(m, h) = \sum_{\substack{\lambda \in L+h \\ (\lambda, \lambda) = m}} \mathbb{D}(\lambda).$$

Note that $\mathbb{D}(m, h)$ is locally finite. We let $\Gamma = \text{Fix}(L^\sharp/L) \subset G$ and write

$$X = \Gamma \backslash \mathbb{D}$$

for the resulting quasi-projective variety. Further, we let $Z(x)$ respectively $Z(m, h)$ be the image of $\mathbb{D}(\lambda)$ respectively $\mathbb{D}(m, h)$ in X .

2.2. The unitary Lie algebra

We let $\mathfrak{g}_0 = \mathfrak{u}(V)$ be the Lie algebra of G . We define the \mathbb{R} -linear surjective map

$$\phi_V : \bigwedge_{\mathbb{R}}^2 V \longrightarrow \mathfrak{u}(V)$$

by

$$\phi_V(v \wedge \tilde{v})(x) = (v, x)\tilde{v} - (\tilde{v}, x)v$$

Note that we have

$$\phi_V(iv \wedge \tilde{v}) = \phi_V(v \wedge -i\tilde{v}).$$

In the following we will abuse notation and drop ϕ_V and just write $v \wedge \tilde{v} \in \mathfrak{u}(V)$. Note that in this way we realize $\mathfrak{u}(V)$ as a quotient of $\bigwedge_{\mathbb{R}}^2 V$. We have

$$\mathfrak{g}_0 = \text{span}_{\mathbb{R}}\{v_r \wedge v_s, iv_r \wedge v_s\}.$$

We put

$$X_{rs} = v_r \wedge v_s \quad \text{and} \quad Y_{rs} = iv_r \wedge v_s.$$

In the Cartan decomposition $\mathfrak{g}_0 = \mathfrak{k}_0 \oplus \mathfrak{p}_0$ with $\mathfrak{k}_0 = \text{Lie}(K) = \mathfrak{u}(p) \times \mathfrak{u}(q)$, we note that

$$\mathfrak{p}_0 = \text{span}_{\mathbb{R}}\{X_{\alpha\mu}, Y_{\alpha\mu}; 1 \leq \alpha \leq p, p+1 \leq \mu \leq m\}.$$

We let $\{\omega_{\alpha\mu}, \omega'_{\beta\nu}\}$ be the corresponding dual basis for \mathfrak{p}_0^* . Furthermore, the natural complex structure on \mathfrak{p}_0 is given by $X_{\alpha\mu} \mapsto Y_{\alpha\mu}$; $Y_{\alpha\mu} \mapsto -X_{\alpha\mu}$.

We let $\mathfrak{g} = \mathfrak{g}_0 \otimes \mathbb{C}$ be the complexification of \mathfrak{g}_0 , which we view as a right \mathbb{C} vector space. We define

$$Z'_{rs} = \frac{1}{2}(X_{rs} - Y_{rs}i) \quad \text{and} \quad Z''_{rs} = \frac{1}{2}(X_{rs} + Y_{rs}i).$$

Note that $Z''_{rs} = -Z'_{sr}$. In the Harish-Chandra decomposition

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}^+ \oplus \mathfrak{p}^-,$$

we see that

$$\mathfrak{k} = \text{span}_{\mathbb{C}}\{Z'_{\alpha\beta}, Z'_{\mu\nu}\}, \quad \mathfrak{p}^+ = \text{span}_{\mathbb{C}}\{Z'_{\alpha\mu}\}, \quad \mathfrak{p}^- = \text{span}_{\mathbb{C}}\{Z''_{\alpha\mu}\}.$$

We let $\{\xi'_{\alpha\mu}\}$ and $\{\xi''_{\alpha\mu}\}$ be the corresponding dual basis of \mathfrak{p}^+ and \mathfrak{p}^- .

We let $V_{\mathbb{C}} = V \otimes_{\mathbb{R}} \mathbb{C}$ and view $V_{\mathbb{C}}$ as a right complex vector space of dimension $2m$. We decompose $V_{\mathbb{C}} = V' \oplus V''$ into the $+i$ and $-i$ eigenspaces under left multiplication by i . The maps

$$v \longmapsto v - ivi \quad \text{and} \quad v \longmapsto v + ivi$$

realize a \mathbb{C} -linear isomorphism of (the left \mathbb{C} -vector space) V with (the right \mathbb{C} -vector space) V' and a \mathbb{C} -anti-linear isomorphism with V and V'' . Hence we can view $V'' \simeq V^*$ as \mathbb{C} -vector spaces. We denote the natural bases of V' and V'' by

$$v'_r := v_r - iv_r i \quad \text{and} \quad v''_r := v_r + iv_r i,$$

respectively. Furthermore, we obtain decompositions $V' = V'_+ \oplus V'_-$ and $V'' = V''_+ \oplus V''_-$ in the natural way. We have

$$Z'_{rs}(v'_t) = -(v_s, v_t)v'_r \quad \text{and} \quad Z'_{rs}(v''_t) = (v_r, v_t)v''_s,$$

and we note that this realizes the isomorphism $\mathfrak{g} \simeq \mathfrak{gl}_m(\mathbb{C})$ by the action of \mathfrak{g} on V' . More precisely, we obtain

$$\mathfrak{k} \simeq \text{Hom}(V'_+, V'_+) \oplus \text{Hom}(V'_-, V'_-), \quad \mathfrak{p}^+ \simeq \text{Hom}(V'_-, V'_+), \quad \mathfrak{p}^- \simeq \text{Hom}(V''_+, V''_-).$$

Correspondingly, the action \mathfrak{g} on V'' realizes the dual representation to the standard representation of \mathfrak{g} .

We write $V_{\mathbb{R}}$ for V considered as a real quadratic space together with the quadratic form $(\cdot, \cdot)_{\mathbb{R}} = \operatorname{Re}(\cdot, \cdot)$. Then $\{v_{\alpha}, iv_{\alpha}, v_{\mu}, iv_{\mu}\}$ forms an orthogonal basis of $V_{\mathbb{R}}$.

We let $\mathfrak{o}_{V_{\mathbb{R}}}$ be the Lie algebra of the orthogonal group $O(V_{\mathbb{R}})$. We now have the isomorphism

$$\phi_{V_{\mathbb{R}}} : \bigwedge^2 V_{\mathbb{R}} \simeq \mathfrak{o}(V_{\mathbb{R}})$$

given by

$$\phi_{V_{\mathbb{R}}}(v_1 \wedge v_2)(x) = (v_1, x)_{\mathbb{R}} v_2 - (v_2, x)_{\mathbb{R}} v_1.$$

We let $\iota : \mathfrak{g}_0 = \mathfrak{u}(V) \mapsto \mathfrak{o}(V_{\mathbb{R}})$ be the natural embedding. We easily see

$$\iota(\phi_V(v \wedge \tilde{v})) = \phi_{V_{\mathbb{R}}}(v \wedge \tilde{v}) + \phi_{V_{\mathbb{R}}}(iv \wedge i\tilde{v}).$$

Note this realizes $\mathfrak{u}(V)$ as the subspace of $\bigwedge^2_{\mathbb{R}} V$ which is fixed by (left)-multiplication with i in both factors.

3. Schwartz forms

3.1. Weil representation

Let $\mathcal{S}(V)$ be the Schwartz space of V . We consider the Weil representation for the dual reductive pair $U(1, 1) \times U(V)$, associated to the additive character $e^{2\pi it}$ acting in the Schrödinger model on $\mathcal{S}(V)$. The setup of the Weil representation in the polynomial Fock model is explained in detail in the Appendix B. We note that $U(V)$ acts linearly, with $\omega(g)\phi(x) = \phi(g^{-1}x)$, for any Schwartz form $\phi(x)$. For matrices in $\operatorname{SL}_2(\mathbb{R}) \simeq \operatorname{SU}(1, 1)$ it is given as follows, see e.g. [22, Sec. 1] for details (see also Shintani [31]):

$$\begin{aligned} \omega\left(\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}\right)\phi(x) &= e^{2\pi ib(x,x)}\phi(x), \\ \omega\left(\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}\right)\phi(x) &= a^m\phi(ax), \\ \omega\left(\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}\right)\phi(x) &= i^{q-p}\widehat{\phi}(-x), \end{aligned}$$

where $\widehat{\phi}(x) = \int_V \phi(y)e(2(y,x)_{\mathbb{R}})dy$ denotes the Fourier transform of $\phi(x)$.

For a Schwartz form ϕ , we say that ϕ has weight $r \in \mathbb{Z}$ if $\omega(k'_{\theta})\phi = e^{ri\theta}\phi$ for $k'_{\theta} = \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix}$ in $K' = U(1) \simeq \operatorname{SO}(2)$, the maximal compact subgroup of $\operatorname{SL}_2(\mathbb{R}) \simeq \operatorname{SU}(1, 1)$ ⁴ Now, for $\tau = u + iv \in \mathbb{H}$ let g'_{τ} be any element of $\operatorname{SL}_2(\mathbb{R})$, mapping i to τ . Then we set

$$\phi(x, \tau) := v^{-\frac{r}{2}}\omega(g'_{\tau})\phi(x) = v^{-\frac{r}{2} + \frac{p+q}{2}}\phi^0(\sqrt{v}x)e^{\pi i(x,x)\tau},$$

where $\phi^0(x) = e^{\pi(x,x)}\phi(x)$.

⁴We could also work with $U(1) \times U(1)$ inside $U(1, 1)$, but we won't need this for our purposes.

3.2. The Kudla-Millson form φ_{KM}

We now consider the complex $[\mathcal{S}(V) \otimes \mathcal{A}^\bullet(\mathbb{D})]^G$ of G -invariant Schwartz functions on V with values in the differential forms on \mathbb{D} . Note that evaluation at the base point z_0 yields an isomorphism

$$[\mathcal{S}(V) \otimes \mathcal{A}^\bullet(\mathbb{D})]^G \simeq [\mathcal{S}(V) \otimes \bigwedge^\bullet(\mathfrak{p}^*)]^K.$$

We use the same symbol for corresponding objects. For example, given $z \in \mathbb{D}$, the standard majorant $(x, x)_z$ is given by

$$(x, x)_z = (x_{z^\perp}, x_{z^\perp}) - (x_z, x_z),$$

where $x = x_z + x_{z^\perp}$ using the orthogonal decomposition $V = z \oplus z^\perp$. Then the standard Gaussian on V is given by

$$\varphi_0(x, z) = e^{-\pi(x, x)_z} \in [\mathcal{S}(V) \otimes \mathcal{A}^0(\mathbb{D})]^G,$$

and evaluation at the base point gives $\varphi_0(x) = \varphi_0(x, z_0) = e^{-\pi \sum_{i=1}^m |z_i|^2} \in \mathcal{S}(V)^K$.

Following [26, Proposition 5.2] and [28, Section 5], we define the differential operator

$$\mathcal{D} = \frac{1}{2^{2q}} \prod_{\mu=p+1}^m \left\{ \sum_{\alpha=1}^p \left(\bar{z}_\alpha - \frac{1}{\pi} \frac{\partial}{\partial z_\alpha} \right) \otimes A'_{\alpha\mu} \right\} = \frac{1}{2^{2q}} \prod_{\mu=p+1}^m \left\{ \sum_{\alpha=1}^p \mathcal{D}_\alpha \otimes A'_{\alpha\mu} \right\},$$

where $\frac{\partial}{\partial z_\alpha} = \frac{1}{2} \left(\frac{\partial}{\partial x_\alpha} - i \frac{\partial}{\partial y_\alpha} \right)$, and $A'_{\alpha\mu}$ denotes the left multiplication with $\xi'_{\alpha\mu}$. Also, we have set $\mathcal{D}_\alpha := \left(\bar{z}_\alpha - \frac{1}{\pi} \frac{\partial}{\partial z_\alpha} \right)$.

Following Kudla and Millson [26], we then define

$$\varphi_{KM} := \mathcal{D} \bar{\mathcal{D}} \varphi_0 \in [\mathcal{S}(V) \otimes \bigwedge^{q,q}(\mathfrak{p}^*)]^K \simeq [\mathcal{S}(V) \otimes \mathcal{A}^{q,q}(\mathbb{D})]^G.$$

Thus, using multi-index notation with $\underline{\alpha} = \{\alpha_1, \dots, \alpha_q\}$ and $\underline{\beta} = \{\beta_1, \dots, \beta_q\}$,

$$\varphi_{KM} = \frac{1}{2^{2q}} \sum_{\underline{\alpha}, \underline{\beta}} \mathcal{D}_{\underline{\alpha}} \bar{\mathcal{D}}_{\underline{\beta}} \varphi_0 \otimes \Omega_q(\underline{\alpha}; \underline{\beta}),$$

where $\mathcal{D}_{\underline{\alpha}} = \prod_{j=1}^q \mathcal{D}_{\alpha_j}$ and

$$\begin{aligned} \Omega_q(\underline{\alpha}; \underline{\beta}) &= \xi'_{\alpha_1 p+1} \wedge \cdots \wedge \xi'_{\alpha_q p+q} \wedge \xi''_{\beta_1 p+1} \cdots \wedge \xi''_{\beta_q p+q} \\ &= (-1)^{q(q-1)/2} \xi'_{\alpha_1 p+1} \wedge \xi''_{\beta_1 p+1} \wedge \cdots \wedge \xi'_{\alpha_q p+q} \wedge \xi''_{\beta_q p+q}. \end{aligned}$$

The properties of the Schwartz form φ_{KM} are summarized in the following theorem

Theorem 3.1 (Kudla-Millson). *The Schwarz form φ_{KM} has the following properties:*

1. φ_{KM} is an eigenfunction of weight $p+q$ under the operation of K' [see 26].

2. As a differential form, $\varphi_{KM}(x, z)$ is closed for every $x \in V$ [see 26, Section 4].
3. The Thom Lemma holds for φ_{KM} [see 27, Theorem 4.1], i.e.,

$$\int_{\Gamma_x \setminus \mathbb{D}} \eta \wedge \varphi_{KM}(x) = i^{-q} \int_{\Gamma_x \setminus \mathbb{D}(x)} \eta$$

for any compactly supported closed differential $(2p - 1)q$ form η on $\Gamma_x \setminus \mathbb{D}$.

3.3. The Schwartz form ψ

We define another Schwartz form ψ by setting

$$\psi := \frac{2i(-1)^{q-1}}{2^{2(q-1)}} \sum_{\substack{\underline{\alpha} = \{\alpha_1, \dots, \alpha_{q-1}\} \\ \underline{\beta} = \{\beta_1, \dots, \beta_{q-1}\}}} \mathcal{D}_{\underline{\alpha}} \bar{\mathcal{D}}_{\underline{\beta}} \varphi_0 \otimes \Omega_{q-1}(\underline{\alpha}; \underline{\beta}) \in [\mathcal{S}(V) \otimes \bigwedge^{q-1, q-1}(\mathfrak{p}^*)],$$

where

$$\begin{aligned} & \Omega_{q-1}(\underline{\alpha}; \underline{\beta}) \\ &= (-1)^{q(q-1)/2} \sum_{j=1}^q \xi'_{\alpha_{1p+1}} \wedge \xi''_{\beta_{1p+1}} \wedge \cdots \wedge \xi'_{\alpha_{(p+j)p+1}} \wedge \widehat{\xi''_{\beta_{(p+j)p+1}}} \wedge \cdots \wedge \xi'_{\alpha_{(q-1)p+q}} \wedge \xi''_{\beta_{(q-1)p+q}}. \end{aligned}$$

In Appendix A we will employ the Fock model of the Weil representation to show

Proposition 3.2. (i) The Schwartz form ψ is invariant under the operation of K , that is,

$$\psi \in [\mathcal{S}(V) \otimes \bigwedge^{q-1, q-1}(\mathfrak{p}^*)]^K \simeq [\mathcal{S}(V) \otimes \mathcal{A}^{q-1, q-1}(\mathbb{D})]^G.$$

(ii) The Schwartz form ψ is an eigenfunction of weight $p + q - 2$ under the operation of K' .

The main property linking φ_{KM} and ψ is the following.

Theorem 3.3. Let $d = \frac{1}{2}(\partial + \bar{\partial})$ and $d^c = \frac{1}{4\pi i}(\partial - \bar{\partial})$ be the standard exterior derivatives acting on $\mathcal{A}^\bullet(\mathbb{D})$, and let $L_\kappa = -2iv^2 \frac{\partial}{\partial \bar{\tau}}$ be the Maass lowering operator of weight κ acting on functions on the upper half plane. Then

$$L_{p+q} \varphi_{KM}^0(x, \tau, z) = dd^c \psi^0(x, \tau, z).$$

This implies

$$v \frac{\partial}{\partial v} \varphi_{KM}^0(\sqrt{v}x) = dd^c \psi^0(\sqrt{v}x).$$

Proof. This is carried out in Appendix A, again using the Fock model. \square

In order to derive a more explicit description of the Schwartz form ψ , when evaluated at the base point z_0 , we examine the properties of the differential operators \mathcal{D}_α and $\bar{\mathcal{D}}_\alpha$ for $\alpha \in \{1, \dots, p\}$. First, we note that all the differential operators commute, i.e. $\mathcal{D}_\alpha \mathcal{D}_\beta = \mathcal{D}_\beta \mathcal{D}_\alpha$, $\bar{\mathcal{D}}_\alpha \bar{\mathcal{D}}_\beta = \bar{\mathcal{D}}_\beta \bar{\mathcal{D}}_\alpha$ and $\mathcal{D}_\alpha \bar{\mathcal{D}}_\beta = \bar{\mathcal{D}}_\alpha \mathcal{D}_\beta$ for all $\alpha, \beta \in \{1, \dots, p\}$.

Further, by direct calculation, we get

$$\mathcal{D}_\alpha \varphi_0 = 2\bar{z}_\alpha \varphi_0, \quad \bar{\mathcal{D}}_\alpha \varphi_0 = 2z_\alpha \varphi_0 \quad \text{and} \quad \mathcal{D}_\alpha \bar{\mathcal{D}}_\alpha \varphi_0 = \left(4|z_\alpha|^2 - \frac{2}{\pi}\right) \varphi_0.$$

In fact (see e.g., [27, p. 303 (6.41)]),

$$\mathcal{D}_\alpha^k \bar{\mathcal{D}}_\alpha^k \varphi_0 = (\mathcal{D}_\alpha \bar{\mathcal{D}}_\alpha)^k \varphi_0 = \left(\frac{1}{\pi}\right)^k 2^k k! L_k(2\pi|z_\alpha|^2) \varphi_0, \quad (3.1)$$

where $L_k(t) = \frac{e^t}{k!} \left(\frac{d}{dt}\right)^k (e^{-t} t^k)$ is the k -th Laguerre polynomial. More generally, we get

$$\mathcal{D}_\alpha^l \bar{\mathcal{D}}_\alpha^k \varphi_0 = 2^k \sum_{m=0}^l \binom{l}{m} \sum_{n=0}^{\min(m,k)} \bar{z}_\alpha^{l-n} z_\alpha^{k-n} \binom{m}{n} \frac{k!}{(k-n)!} \left(\frac{-1}{\pi}\right)^n \varphi_0. \quad (3.2)$$

Hence the Schwartz form ψ can be expressed using (in general non-homogeneous) polynomials $P_{\underline{\alpha}, \underline{\beta}}^{2q-2} \in \mathcal{P}(V)$ as follows:

$$\psi(x, z_0) = \frac{2i(-1)^{q-1}}{2^{2(q-1)}} \sum_{\underline{\alpha}, \underline{\beta}} P_{\underline{\alpha}, \underline{\beta}}^{2q-2}(x) \varphi_0(x) \otimes \Omega_{q-1}(\underline{\alpha}; \underline{\beta}), \quad (3.3)$$

$$\psi^0(x, z_0) = \frac{2i(-1)^{q-1}}{2^{2(q-1)}} \sum_{\underline{\alpha}, \underline{\beta}} P_{\underline{\alpha}, \underline{\beta}}^{2q-2}(x) e^{-2\pi R(x, z_0)} \otimes \Omega_{q-1}(\underline{\alpha}; \underline{\beta}). \quad (3.4)$$

The following lemma is easily obtained.

Lemma 3.4. *For any pair of multi-indices $\underline{\alpha}, \underline{\beta} \in \{1, \dots, p\}^{q-1}$, the attached polynomial $P_{\underline{\alpha}, \underline{\beta}}^{2q-2}(x)$ has the following properties:*

1. *It has degree $2q - 2$ and depends only on V_+ .*
2. *The leading term is given by*

$$2^{2(q-1)} \prod_{l=1}^{q-1} \bar{z}_{\alpha_l} \prod_{k=1}^{q-1} z_{\beta_k}.$$

3. *All monomials occurring in $P_{\underline{\alpha}, \underline{\beta}}^{2q-2}(x)$ have even degree.*
4. *The constant term is non-zero if and only if for every $\alpha \in \{1, \dots, p\}$ the multiplicity of α in the multi-indices $\underline{\alpha}$ and $\underline{\beta}$ is the same. In which case, $P_{\underline{\alpha}, \underline{\beta}}^{2q-2}(x)$ is a product of Laguerre functions, and the constant term is given by*

$$2^{q-1} \left(\frac{-1}{\pi}\right)^{q-1} \prod_{\alpha \in \underline{\alpha}} m(\alpha)!,$$

where $m(\alpha)$ is the multiplicity of α .

In particular, the situation in part 4 of the lemma occurs when $x = z_\alpha v_\alpha$, and only the terms with $\underline{\alpha} = \underline{\beta} = (\alpha, \alpha, \dots, \alpha)$ are non-zero.

It will often be convenient to write $P_{\underline{\alpha}, \underline{\beta}}^{2q-2}$ as a sum of its homogeneous components

$$P_{\underline{\alpha}, \underline{\beta}}^{2q-2}(x) = \sum_{\ell=0}^{q-1} P_{\underline{\alpha}, \underline{\beta}; 2\ell}^{2q-2}(x),$$

with 2ℓ the respective weight. Note that $P_{\underline{\alpha}, \underline{\beta}; 2\ell}^{2q-2}(wx) = |w|^{2\ell} P_{\underline{\alpha}, \underline{\beta}; 2\ell}^{2q-2}(x)$ for any $w \in \mathbb{C}$.

Remark 3.5. We mention that besides (3.2) the polynomials $P_{\underline{\alpha}, \underline{\beta}}^{2q-2}(x)$ can also be expressed using derivatives of Laguerre functions by (3.1) or, alternatively through Hermite functions in the real and imaginary parts of the z_α 's as indeterminates.

4. A singular Schwartz form

Analogously to Kudla [25] for $O(p, 2)$, we define for $x \neq 0$ the singular Schwartz form

$$\Psi^0(x, z) := - \int_1^\infty \psi^0(\sqrt{t}x, z) \frac{dt}{t}.$$

The form Ψ^0 has its singularities where $R(x, z) = 0$, i.e., precisely along the cycles $\mathbb{D}(x)$. Thus, in particular, $\Psi^0(x, z)$ is smooth for $(x, x) \leq 0$. We also set

$$\Psi(x, z) = \Psi^0(x, z) e^{-\pi(x, x)}.$$

Recall the definition of the incomplete Γ -function, $\Gamma(s, a) = \int_a^\infty t^{s-1} e^{-t} dt$. The following lemma is obtained by a straightforward calculation.

Lemma 4.1. *At the base point $z = z_0$, the singular Schwartz form Ψ^0 is given by*

$$\Psi^0(x, z_0) = \frac{2i(-1)^{q-1}}{2^{2(q-1)}} \sum_{\underline{\alpha}, \underline{\beta}} \left[\sum_{\ell=0}^{q-1} P_{\underline{\alpha}, \underline{\beta}; 2\ell}^{2q-2}(x) (2\pi R(x, z_0))^{-\ell} \Gamma(\ell, 2\pi R(x, z_0)) \right] \otimes \Omega_{q-1}(\underline{\alpha}; \underline{\beta}).$$

We conclude that $R(x, z)^{q-1} \Psi^0(x, z)$ extends to a smooth differential $(q-1, q-1)$ -form on \mathbb{D} .

While it should be emphasized that Ψ is not a Schwartz function on V , we nonetheless define (as if Ψ had weight $p+q$)

$$\Psi(x, \tau, z) = \Psi^0(\sqrt{v}x, z) e^{\pi i(x, x)\tau} \quad (\tau \in \mathbb{H}).$$

This is motivated by the second statement in the Proposition below. Note that

$$\Psi(x, \tau, z) = - \left(\int_v^\infty \psi^0(\sqrt{t}x, z) \frac{dt}{t} \right) e^{\pi i(x, x)\tau}. \quad (4.1)$$

From the definition of Ψ and the properties of ψ , we get

Proposition 4.2. *Outside the singularities, $\Psi(x, \tau, z)$ has the following properties:*

1. *For d and d^c the standard exterior differentials on $\mathcal{A}^\bullet(\mathbb{D})$, we have outside $\mathbb{D}(x)$*

$$dd^c \Psi(x, \tau, z) = \varphi_{KM}(x, \tau, z).$$

2. *We have*

$$L_{p+q} \Psi(x, \tau, z) = \psi(x, \tau, z),$$

with the Maass lowering operator L_{p+q} as before.

Proof. 1. This follows from Theorem 3.3 and the rapid decay of the Schwartz form φ_{KM} :

$$\begin{aligned} dd^c \Psi(x, \tau, z) &= - \left(\int_v^\infty dd^c \psi(\sqrt{t}x, z) \frac{dt}{t} \right) e^{\pi i(x,x)\tau} \\ &= - \left(\int_v^\infty \frac{\partial}{\partial t} \varphi_{KM}^0(\sqrt{t}x, z) dt \right) e^{\pi i(x,x)\tau} = \varphi_{KM}(x, \tau, z). \end{aligned}$$

2. Immediately from the definition,

$$\begin{aligned} L_{p+q} \Psi(x, \tau, z) &= 2iv \frac{\partial}{\partial \bar{\tau}} \left(\int_v^\infty \psi^0(\sqrt{t}x, z) \frac{dt}{t} \right) e^{\pi i(x,x)\tau} \\ &= v \left(\frac{\partial}{\partial v} \int_v^\infty \psi^0(\sqrt{t}x, z) \frac{dt}{t} \right) e^{\pi i(x,x)\tau} = \psi^0(\sqrt{v}x, z) e^{\pi i(x,x)\tau} = \psi(x, \tau, z), \end{aligned}$$

again by rapid decay. □

4.1. The current equation

We denote by $\mathcal{A}_c^k(\mathbb{D})$ the space of compactly supported differential forms on \mathbb{D} of degree k . Recall that a locally integrable degree k -form ω on \mathbb{D} defines a current, i.e., a (continuous) linear functional on the compactly supported forms of complementary degree, via

$$[\omega](\eta) := \int_{\mathbb{D}} \eta \wedge \omega \quad (\eta \in \mathcal{A}_c^{2pq-k}(\mathbb{D})).$$

Furthermore, for the exterior derivatives of a current $[\omega]$ we have

$$dd^c[\omega](\eta) := [\omega](dd^c \eta).$$

The goal of this section is to prove the following generalization of the Thom Lemma, see Theorem 3.13.

Theorem 4.3. *Let $x \in V$ and let $\delta_{Z(x)}$ denote the delta current for the special cycle $Z(x)$. Then*

$$dd^c[\Psi^0(x)] + (-i)^q \delta_{Z(x)} = [\varphi_{KM}^0(x)]$$

as currents on $\Gamma_x \backslash \mathbb{D}$. In other words, we have

$$\int_{\Gamma_x \backslash \mathbb{D}} dd^c \eta \wedge \Psi^0(x) + (-i)^q \int_{Z(x)} \eta = \int_{\Gamma_x \backslash \mathbb{D}} \eta \wedge \varphi_{KM}^0(x)$$

for any $\eta \in \mathcal{A}_c^{2(p-1)q}(\Gamma_x \backslash \mathbb{D})$.

We prove the theorem in the next two subsections following the same method employed in [6] and [16].

With this we can now define a Green current for the special cycles $Z(h, m) \subset X$. Namely, for $m \in \mathbb{Q}$, $h \in L^\# / L$ satisfying $m \equiv \frac{1}{2}(h, h) \pmod{\mathbb{Z}}$ and a real parameter $w > 0$ we introduce the Green current of Kudla type on X by setting

$$\Xi^K(m, w, h)(z) := \sum_{\substack{\lambda \in L+h \\ (\lambda, \lambda) = 2m \\ \lambda \neq 0}} \Psi^0(\sqrt{w}\lambda, z). \quad (4.2)$$

Then by Theorem 4.3 we immediately obtain

Corollary 4.4. *The singular differential $(q-1, q-1)$ -form $\Xi^K(m, w, h)$ defines a Green current for the cycle $Z(m, h)$ on X .*

4.1.1. Local integrability

Proposition 4.5. *Let $x \in V$. Then $\Psi^0(x)$ and $d^c \Psi^0(x)$ are locally integrable differential forms on \mathbb{D} .*

Proof. We view a top-degree differential form $\phi \in \mathcal{A}^{2pq}(\mathbb{D})$ via the Hodge $*$ -operator as a (K -invariant) function on G . We pick suitable coordinates on \mathbb{D} , using the decomposition $G = HAK$, where H is the stabilizer of the first basis vector v_1 of V , A is a one parameter subgroup $A = \{a_t = \exp(tX_{1p+q}); t \in \mathbb{R}\}$. Set $A_0 = \{a_t : t \geq 0\}$. Then, see [14, Sec. 2] or [30, Section 2] for details,

$$\int_{\mathbb{D}} \phi = \int_G \phi(g) dg = C \int_{A_0} \int_H \phi(ha_t) \sinh(t)^{2q-1} \cosh(t)^{2p-1} dh dt, \quad (4.3)$$

with C a positive constant, depending on the normalization of the invariant measures.

Now $\Psi^0(x)$ is smooth unless $(x, x) > 0$. In that case we may assume that $x = \sqrt{m}v_1$, for some $m > 0$. Then for $\eta \in \mathcal{A}_c^{2(pq-(q-1))}(\mathbb{D})$. We set $\phi = \eta \wedge \Psi(x)$ and see

$$\phi(ha_t) = \eta(ha_t) \wedge \Psi^0(a_t^{-1}h^{-1}\sqrt{m}v_1),$$

wherein

$$a_t^{-1}h^{-1}\sqrt{m}v_1 = \cosh(t)\sqrt{m}v_1 - \sinh(t)\sqrt{m}v_{p+q}.$$

Hence,

$$(a_t^{-1}h^{-1}\sqrt{m}v_1)_{z_0} = -\sinh(t)\sqrt{m}v_{p+q} \quad \text{and} \quad (a_t^{-1}h^{-1}\sqrt{m}v_1)_{z_0^\perp} = \cosh(t)\sqrt{m}v_1. \quad (4.4)$$

Thus, we have (see Lemma 4.1),

$$\begin{aligned} \Psi^0(a_t^{-1}h^{-1}\sqrt{m}v_1) &= \frac{2i(-1)^{q-1}}{2^{2(q-1)}} \left[\sum_{\ell=0}^{q-1} (2\pi m \sinh^2(t))^{-\ell} \Gamma(\ell, 2\pi m \sinh^2(t)) \right. \\ &\quad \left. \cdot \sum_{\underline{\alpha}, \underline{\beta}} P_{\underline{\alpha}, \underline{\beta}; 2\ell}^{2q-2}(\kappa\sqrt{m} \cosh(t)v_1) \right] \otimes \Omega_{q-1}(\underline{\alpha}; \underline{\beta}). \end{aligned}$$

We conclude that the integrand of (4.3), i.e.,

$$\eta(ha_t) \wedge \Psi^0(a_t^{-1}h^{-1}\sqrt{m}v_1) \sinh(t)^{2q-1} \cosh(t)^{2p-1}$$

is bounded, in fact, vanishes, as $t \rightarrow 0$. Further, as η has compact support, the integral is convergent.

For the local integrability of $d^c\Psi(x)$ the reasoning is similar, but a bit more tedious. Again, we may assume that $x = \sqrt{m}v_1$, with $m > 0$. Further, note that we only need to consider highest-degree terms.

Note $d^c\Psi^0(x) = -\int_1^\infty d^c\psi^0(\sqrt{s}x) \frac{ds}{s}$, which can be evaluated similarly to Lemma 4.1. By (A.1), $d^c\psi$ consists of two parts. Both involve polynomials of degree $2q-1$ which depend on the positive coordinates of x (note that there is no constant part). If by (4.4), we set $x = \cosh(t)\sqrt{m}v_1$, only the polynomials which depend exclusively on the first vector can contribute to $d^c\Psi^0(x)$. From their highest-degree terms, we get

$$2^{-(2q-1)} \cosh(t)^{2q-1} m^{q-\frac{1}{2}} \sqrt{s}^{2q-1}.$$

Also, in (A.1) there are linear homogenous polynomials in the negative coordinates, $Q'_{\alpha'_q, \underline{\alpha}'_{(q-1)}}$ and $Q'_{\alpha_q, \underline{\alpha}'_{(q-1)}}$. From them, again by (4.4) we have contributions of

$$-\sqrt{s} \sqrt{m} \sinh(t)$$

Hence, gathering the contributions of the non-vanishing highest-degree terms, we still have the integral

$$\begin{aligned} \int_1^\infty s^{q-1} e^{2\pi R(x, z_0)} ds &= (2\pi R(x, z_0))^{-q} \Gamma(q, 2\pi R(x, z_0)) \\ &= (2\pi \sinh^2(t))^{-q} \Gamma(q, 2\pi \sinh^2(t)). \end{aligned}$$

Thus, up to sign, for $t \rightarrow 0$ the behaviour of $d^c\Psi(a_t^{-1}h^{-1}\sqrt{m}v_1)$ is dominated by terms of the form

$$\frac{(-1)^{q-1}\pi}{2^{2q-1}} \sinh(t) \cosh(t)^{2q-1} (\sinh^2(t))^{-q} \Gamma(q, 2\pi \sinh^2(t)). \quad (4.5)$$

In particular, it follows that the integrand in

$$\int_{A_0} \int_H \eta(ha_t) \wedge (d^c\Psi(a_t^{-1}h^{-1}\sqrt{m}v_1)) \sinh(t)^{2q-1} \cosh(t)^{2p-1} dh dt,$$

remains bounded as $t \rightarrow 0$, and hence the integral converges. \square

4.1.2. The current equation

Proof. Let $\eta \in \mathcal{A}_c^{2(p-1)q}(\Gamma_x \setminus \mathbb{D})$, not necessarily closed. First note using $(dd^c \eta) \wedge \Psi^0(x) = (d\eta) \wedge d^c \Psi^0(x) - d^c(d\eta \wedge \Psi^0(x))$ and Stokes' theorem

$$\int_{\Gamma_x \setminus \mathbb{D}} (dd^c \eta) \wedge \Psi^0(x) = - \int_{\Gamma_x \setminus \mathbb{D}} (d\eta) \wedge d^c \Psi^0(x) + \lim_{\epsilon \rightarrow 0} \int_{\Gamma_x \setminus \partial(\mathbb{D} - U_\epsilon(x))} (d\eta) \wedge d^c \Psi^0(x),$$

where U_ϵ , ($\epsilon > 0$) denotes an open neighbourhood of the cycle $\mathbb{D}(x)$. Next we show that the limit on the right hand side vanishes. We may again assume $x = \sqrt{m}v_1$, with $m > 0$ and use the *HAK* coordinates introduced on p. 14. Then for $\epsilon > 0$, an open neighborhood of $\mathbb{D}(v_1)$ is defined by

$$U_\epsilon = \mathbb{D} - (H \times A_\epsilon), \quad (4.6)$$

with $A_\epsilon = \{a_t : t \geq \epsilon\}$. With the analog of the integral formula from (4.3), the limit can be written as

$$C \lim_{\epsilon \rightarrow 0} \int_{\Gamma_{v_1} \setminus H} \eta(ha_\epsilon) \wedge \Psi^0(a_\epsilon^{-1}h^{-1}\sqrt{m}v_1) \sinh(\epsilon)^{2q-1} \cosh(\epsilon)^{2p-1} dh$$

for some constant C . Only the highest degree term of $\Psi(a_t^{-1}h^{-1}\sqrt{m}v_1)$ (see Lemma 4.1) can contribute. Further, note that, since $(a_t^{-1}h^{-1}\sqrt{m}v_1)_{z_0^\perp} = \cosh(t)\sqrt{m}v_1$ by (4.4), we have $P_{\underline{\alpha}, \underline{\beta}; 2q-2}^{2q-2}(\sqrt{m} \cosh(t)v_1) \neq 0$ only for $\underline{\alpha} = \underline{\beta} = (1, \dots, 1)$, thus, up to constants, the highest degree term is given by

$$(m \sinh^2(t))^{-(q-1)} \Gamma(q-1, 2\pi m \sinh^2(t)) (2\sqrt{m} \cosh(t))^{2q-1}.$$

Hence, comparing powers of $\sinh(t)$ we see that the integrand goes to zero for $t = \epsilon \rightarrow 0$, and the limit vanishes as claimed.

Now, since $dd^c \Psi(x) = \varphi_{KM}(x)$, we have

$$\begin{aligned} - \int_{\Gamma_x \setminus \mathbb{D}} (d\eta) \wedge d^c \Psi^0(x) &= \int_{\Gamma_x \setminus \mathbb{D}} \eta \wedge dd^c \Psi^0(x) + \int_{\Gamma_x \setminus \mathbb{D}} d(\eta \wedge d^c \Psi^0(x)) \\ &= \int_{\Gamma_x \setminus \mathbb{D}} \eta \wedge \varphi_{KM}^0(x) + \lim_{\epsilon \rightarrow 0} \int_{\Gamma_x \setminus \partial(\mathbb{D} - U_\epsilon(x))} \eta \wedge d^c \Psi^0(x), \end{aligned}$$

again by applying Stokes' theorem. Thus it remains to show that

$$\lim_{\epsilon \rightarrow 0} \int_{\Gamma_x \setminus \partial(\mathbb{D} - U_\epsilon(x))} \eta \wedge d^c \Psi^0(x) = \int_{Z(x)} \eta.$$

We have to consider the limit of the same integral as in the proof of second part of Proposition 4.5:

$$C \lim_{\epsilon \rightarrow 0} \int_{\Gamma_{v_1} \setminus H} \eta(ha_\epsilon) \wedge d^c \Psi^0(a_{-\epsilon}h^{-1}\sqrt{m}v_1) \cosh(\epsilon)^{2p-1} \sinh(\epsilon)^{2q-1} dh, \quad (4.7)$$

with a non-zero constant C , independent of η . With (4.5) we see that for both parts of $d^c\Psi^0(x)$, the integral is bounded as $t = \epsilon \rightarrow 0$. We have

$$\begin{aligned} C \lim_{\epsilon \rightarrow 0} \int_H \eta(ha_\epsilon) \wedge (d^c\Psi^0(a_{-\epsilon}h^{-1}\sqrt{m}v_1)) \cosh(\epsilon)^{2p-1} \sinh(\epsilon)^{2q-1} dh \\ = \tilde{C} \int_H \eta(h) dh, \end{aligned}$$

with a constant \tilde{C} independent of η . By Kudla-Millson theory [see 27, Theorem 6.4], we see that $\tilde{C} = i^{-q}$ for η closed, see Theorem 3.1 3.

To summarize, we have showed that for all $\eta \in \mathcal{A}_c^{2(p-1)q}(\Gamma_x \backslash \mathbb{D})$,

$$\int_{\Gamma_x \backslash \mathbb{D}} (dd^c\eta) \wedge \Psi^0(x) = \int_{\Gamma_x \backslash \mathbb{D}} \eta \wedge \varphi_{KM}^0(x) - \int_{Z(x)} \eta,$$

as claimed. \square

5. The singular theta lift

5.1. Weak Maass forms

We let L be an even hermitian lattice with the hermitian form (\cdot, \cdot) . Further, we denote by L^- the same $\mathcal{O}_{\mathbb{F}}$ module L but with the hermitian form $-(\cdot, \cdot)$.

We denote the standard basis elements of $\mathbb{C}[L^\sharp/L]$ by \mathbf{e}_h ($h \in L^\sharp/L$) and introduce the Hermitian pairing

$$\langle \cdot, \cdot \rangle_L : \mathbb{C}[L^\sharp/L] \times \mathbb{C}[L^\sharp/L] \rightarrow \mathbb{C}, \quad \text{by setting } \langle \mathbf{e}_\mu, \mathbf{e}_\nu \rangle_L = \delta_{\mu, \nu} \quad (\mu, \nu \in L^\sharp/L).$$

Similar definitions are made for the lattice L^- .

Now, recall that from the Weil representation ω of $\mathrm{SL}_2(\mathbb{Z})$ one obtains a unitary representation of $\mathrm{SL}_2(\mathbb{Z})$ on $\mathbb{C}[L^\sharp/L]$, which we denote by ω_L (see [6, p. 52]). It is most easily described through the action of the generators $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $T = \begin{pmatrix} 1 & \\ & 1 \end{pmatrix}$:

$$\omega_L(T)\mathbf{e}_h = e((h, h))\mathbf{e}_h, \quad \omega_L(S)\mathbf{e}_h = \frac{i^{p-q}}{\sqrt{|L^\sharp/L|}} \sum_{\mu \in L^\sharp/L} e(-2(\mu, h)_{\mathbb{R}})\mathbf{e}_\mu.$$

We denote by $\omega_L^\vee = \bar{\omega}_L$ the dual representation. Note that $\omega_L^\vee = \omega_{L^-}$. For $k \in \mathbb{Z}$ and $\gamma \in \mathrm{SL}_2(\mathbb{Z})$, define the weight k -slash operation on functions $\mathbb{C}[L^\sharp/L] \rightarrow \mathbb{C}$ as

$$f|_{k,L} \gamma = (c\tau + d)^{-k} \omega_L(\gamma)f.$$

The slash-operation for the dual representation is defined similarly.

Following [6] and [12] we now define several spaces of modular forms.

Definition 5.1 ([see 6, Section 3]). For $k \in \mathbb{Z}$, let $\mathbf{H}_{k,L}$ be the space of twice continuously differentiable functions $f : \mathbb{H} \rightarrow \mathbb{C}[L^\sharp/L]$, which satisfy

1. $f|_{k,L}(\gamma) = f$ for all $\gamma \in \mathrm{SL}_2(\mathbb{Z})$.
2. There exists a constant $C > 0$ such that $f(\tau) = \mathbf{O}(e^{Cv})$ as $v \rightarrow \infty$.
3. $\Delta_k f = 0$.

The elements of $\mathbf{H}_{k,L}$ are called harmonic weak Maass forms. Any such form f has a decomposition $f(\tau) = f^+(\tau) + f^-(\tau)$ into a holomorphic and a non-holomorphic part, where the Fourier expansion of the holomorphic part is

$$f^+(\tau) = \sum_{h \in L^\sharp/L} \sum_{n \in \mathbb{Q}} a^+(h, n) e(n\tau) \mathbf{e}_h,$$

whilst that of the non-holomorphic part is

$$f^-(\tau) = \sum_{h \in L^\sharp/L} \left(a^-(h, 0) v^{1-k} + \sum_{\substack{n \in \mathbb{Q} \\ n \neq 0}} a^-(h, n) \Gamma(1-k, 4\pi m v) e(nu) \right) \mathbf{e}_h.$$

We denote by $P(f)$ the principal part of f , i.e. the Fourier polynomial

$$P(f)(\tau) = P(f^+)(\tau) = \sum_{h \in L^\sharp/L} \sum_{\substack{n \in \mathbb{Q} \\ n < 0}} a^+(h, n) e(n\tau) \mathbf{e}_h.$$

Note that $\mathbf{H}_{k,L}$ contains the spaces of weakly holomorphic modular forms $\mathbf{M}_{k,L}^!$ and holomorphic modular forms $\mathbf{M}_{k,L}$, with $\mathbf{H}_{k,L} \supset \mathbf{M}_{k,L}^! \supset \mathbf{M}_{k,L}$.

The operator ξ_k affords an antilinear mapping given by

$$\xi_k : \mathbf{H}_{k,L} \longrightarrow \mathbf{M}_{2-k,L}^!, \quad f(\tau) \longmapsto v^{k-2} \overline{\frac{\partial f(\tau)}{\partial \tau}} = v^{k-2} \overline{L_k f(\tau)}.$$

Now, the space $\mathbf{H}_{k,L}^+$ is defined as the inverse image of the cusp forms \mathbf{S}_{2-k,L^-} . It follows immediately from this definition that for $f \in \mathbf{H}_{k,L}^+$,

$$f(\tau) - P(f)(\tau) = \mathbf{O}(e^{-Cv}),$$

as $v \rightarrow \infty$ for some constant $C > 0$.

Further, by [6, Corollary 3.8], there are exact sequences

$$0 \longrightarrow \mathbf{M}_{k,L}^! \longrightarrow \mathbf{H}_{k,L} \xrightarrow{\xi_k} \mathbf{M}_{2-k,L}^! \longrightarrow 0,$$

and

$$0 \longrightarrow \mathbf{M}_{k,L}^! \longrightarrow \mathbf{H}_{k,L}^+ \xrightarrow{\xi_k} \mathbf{S}_{2-k,L^-} \longrightarrow 0.$$

Following Ehlen and Sankaran [12], we generalize this this setup by introducing two further spaces of modular forms, \mathbf{A}_{k,L^-}^{mod} and $\mathbf{A}_{k,L^-}^!$. For the former space, we use the following, slightly modified definition from [21, Definition 3.2]:

Definition 5.2. Let $A_k^{mod}(\omega_L^\vee) = A_{k,L^-}^{mod}$ denote the space of \mathcal{C}^∞ -functions $f : \mathbb{H} \rightarrow \mathbb{C}[L^\sharp/L]$ satisfying

1. $f|_{k,L^-}(\gamma) = f$ for all $\gamma \in \mathrm{SL}_2(\mathbb{Z})$.
2. For all $a, b \in \mathbb{Z}_{\geq 0}$, there is an $r \in \mathbb{Z}$ such that $\frac{\partial^a}{\partial u^a} \frac{\partial^b}{\partial v^b} f(\tau) = \mathbf{O}(v^r)$ as $v \rightarrow \infty$.
3. If $f = \sum_{m \in \mathbb{Q}} c(m, v) e(m\tau)$ denotes the Fourier expansion of f , then the integral

$$\int_1^\infty c(0, t) t^{-2-s} dv,$$

has a meromorphic continuation to a half-plane $\mathrm{Re}(s) > -\epsilon$ for some $\epsilon > 0$. (The integral converges for sufficiently large $\mathrm{Re}(s) \gg 0$, since by 2., f is of polynomial growth as $v \rightarrow \infty$.)

Definition 5.3 ([see 12, Definition 2.8]). Denote by $A_k^!(\omega_L^\vee) = A_{k,L^-}^!$ the space of \mathcal{C}^∞ functions $f : \mathbb{H} \rightarrow \mathbb{C}[L^\sharp/L]^\vee$ satisfying

1. $f|_{k,L^-}(\gamma) = f$ for all $\gamma \in \mathrm{SL}_2(\mathbb{Z})$.
2. There exists a constant $C > 0$ such that $f(\tau) = \mathbf{O}(e^{Cv})$ as $v \rightarrow \infty$.
3. $L_k(f) \in A_{k-2}^{mod}(\omega_L^\vee)$.

5.2. Regularized theta integral

In the following, we set $\kappa = p + q - 2$ and $k = -\kappa = -(p + q) + 2$. For $h \in L^\sharp/L$ we define the theta function component

$$\theta(\tau, z, \psi)_h = \sum_{\lambda \in L+h} \psi(\lambda, \tau, z). \quad (\tau \in \mathbb{H}, z \in \mathbb{D}).$$

The theta function $\Theta(\tau, z)$ is defined as the vector

$$\Theta(\tau, z) = \Theta(\tau, z, \psi) := (\theta(\tau, z, \psi)_h)_{h \in L^\sharp/L} = \sum_{h \in L^\sharp/L} \theta(\tau, z, \psi)_h \mathbf{e}_h.$$

The theta function transforms as a vector-valued modular form for $\mathrm{SL}_2(\mathbb{Z})$ with weight κ . Explicitly, we can write

$$\Theta(\tau, z) = v \sum_{h \in L^\sharp/L} \sum_{\lambda \in L+h} P_\psi(\sqrt{v}\lambda, z) e^{-\pi v(\lambda, \lambda)_z + i\pi u(\lambda, \lambda)} \mathbf{e}_h, \quad (5.1)$$

with a polynomial $P_\psi(x, z) \in [\mathcal{P}(V) \otimes \mathcal{A}^\bullet(\mathbb{D})]^K$ given, see (3.3), at the base point $z = z_0$ by

$$P_\psi(x, z_0) = \frac{2i(-1)^{q-1}}{2^{2(q-1)}} \sum_{\ell=1}^{q-1} \sum_{\underline{\alpha}, \underline{\beta}} P_{\underline{\alpha}, \underline{\beta}; 2\ell}^{2q-2}(x) \otimes \Omega_{q-1}(\underline{\alpha}; \underline{\beta}).$$

Following Borchers [2], Bruinier [4], and Bruinier and Funke [6], for a weak harmonic Maass form $f \in H_{k,L}^+$, we consider the regularized theta integral

$$\int_{\mathrm{SL}_2(\mathbb{Z}) \backslash \mathbb{H}}^{reg} \left\langle f(\tau), \overline{\Theta(\tau, z)} \right\rangle_L d\mu.$$

Due to invariance under $\mathrm{SL}_2(\mathbb{Z})$, the following regularization recipe (due to Harvey and Moore [18]) can be used: For $t \in \mathbb{R}_{>0}$ denote by \mathcal{F}_t the truncated fundamental domain given by

$$\mathcal{F}_t := \left\{ \tau = u + iv; |\tau| > 1, -\frac{1}{2} < u < \frac{1}{2}, 0 < v \leq t \right\},$$

and define

$$\Phi(z, f, \psi) := \int_{\mathcal{F}}^{reg} \left\langle f, \overline{\Theta(\tau, z)} \right\rangle_L d\mu := \mathrm{CT}_{s=0} \left[\lim_{t \rightarrow \infty} \int_{\mathcal{F}_t} \left\langle f, \overline{\Theta(\tau, z)} \right\rangle_L v^{-s} d\mu \right], \quad (5.2)$$

where the notation $\mathrm{CT}_{s=0}$ denotes the constant term at $s = 0$ of the meromorphic continuation of the limit.

More generally, we introduce a regularized pairing as follows. For $f \in H_{k,L}^+$ and g transforming as a modular form of weight κ under ω_L set

$$(f, g)_{L^-}^{reg} = \mathrm{CT}_{s=0} \left[\lim_{t \rightarrow \infty} \int_{\mathcal{F}_t} \langle f, \bar{g} \rangle_L v^{-s} d\mu \right]. \quad (5.3)$$

We say that *the pairing exists* if for sufficiently large $\mathrm{Re}(s)$ the limit $t \rightarrow \infty$ defines a holomorphic function in s for which a meromorphic continuation to some $\mathrm{Re}(s) < 0$ exists, so that constant of the Laurent expansion around $s = 0$ can be evaluated.

5.3. Singularities and current equation

Let f be a harmonic weak Maass form with holomorphic Fourier coefficients $a^+(h, n)$, $h \in L^\sharp/L$, $n \in \mathbb{Q}_{<0}$. We define a locally finite cycle $\mathbb{D}(f)$ on \mathbb{D} by

$$\mathbb{D}(f) := \sum_{h \in L^\sharp/L} \sum_{n \in \mathbb{Q}_{<0}} a^+(h, n) \mathbb{D}(h.n)$$

and denote by $Z(f)$ the image of $\mathbb{D}(f)$ on X .

Proposition 5.4. *The regularized lift $\Phi(z, f, \psi)$ converges to a smooth differential form on \mathbb{D} with singularities along the cycle $\mathbb{D}(f)$. In a small neighbourhood of $w \in \mathbb{D}$, the singularities are of type*

$$- \sum_{h \in L^\sharp/L} \sum_{\substack{n \in \mathbb{Q} \\ n < 0}} a^+(h, n) \sum_{\substack{\lambda \in L+h \\ (\lambda, \lambda) = -2n \\ \lambda \in w^\perp}} \Psi^0(\lambda, \tau, z),$$

i.e., the difference of $\Phi(z, f, \psi)$ and this sum extends to a smooth form.

Proof. The argument closely follows [6, Sec. 5]. It suffices to consider the integral up to smooth functions. Due to the rapid decay of the non-holomorphic part of f , the integral converges for f^- to a smooth form, and we only need to consider

$$\sum_h \lim_{t \rightarrow \infty} \int_{\mathcal{F}_t}^{reg} f_h^+(\tau) \theta(\tau, z, \psi)_h v^{-s} d\mu.$$

Also, since the integral over \mathcal{F}_1 is smooth, it suffices to consider the integral over $v > 1$:

$$\sum_h \lim_{t \rightarrow \infty} \int_1^t \int_{-\frac{1}{2}}^{\frac{1}{2}} f_h^+(\tau) \theta(\tau, z, \psi)_h v^{-s-2} du dv.$$

Now, the integration over u picks out the constant term in the Fourier expansion of the integrand, which is given by

$$v \sum_h \sum_{\lambda \in L+h} a^+(h, -\frac{1}{2}(\lambda, \lambda)) P_\psi(\lambda\sqrt{v}, z) e^{2\pi v(\lambda_z, \lambda_z)},$$

with the polynomial $P_\psi(x, z) \in [\mathcal{P}(V) \otimes \mathcal{A}(\mathbb{D})]^G$ from (5.1). We have

$$\sum_{\lambda \in L^\sharp} a^+(\lambda, -\frac{(\lambda, \lambda)}{2}) \int_1^\infty P_\psi(\lambda\sqrt{v}, z) e^{2\pi v(\lambda_z, \lambda_z)} v^{-s-1} dv. \quad (5.4)$$

For a relatively compact open neighbourhood $U \subset \mathbb{D}$, define the set

$$S_f(U, \epsilon) = \{ \lambda \in L^\sharp; a^+(\lambda, -\frac{1}{2}(\lambda, \lambda)) \neq 0 \text{ and } \frac{1}{2}|(\lambda_z, \lambda_z)| < \epsilon \text{ for some } z \in U \}.$$

By reduction theory, this set is finite, as f^+ has only finitely many non-vanishing Fourier coefficients in its principal part.

Using standard arguments, like in [6], one finds that in (5.4) the sum of all terms with $\lambda \in L^\sharp - S_f(U, \epsilon)$ is majorized by a convergent sum, $\sum_{\lambda \in L^\sharp} \exp(-\pi(\lambda, \lambda)_z)$, and hence converges. Further, in (5.4), the term with $\lambda = 0$ is given by $a^+(0, 0) P_\psi(0, z) \int_1^\infty \frac{1}{v^{s+1}} dv$. It falls out after regularization, since it has vanishing constant term at $s = 0$.

Finally, all that remains of (5.4) is the following finite sum, which dictates the singularities in U :

$$\sum_{0 \neq \lambda \in S_f(U, \epsilon)} a^+(\lambda, -\frac{(\lambda, \lambda)}{2}) \int_1^\infty P_\psi(\lambda\sqrt{v}, z) e^{2\pi v(\lambda_z, \lambda_z)} v^{-s-1} dv.$$

Clearly, the integral has meromorphic continuation to the entire s -plane, and for $s = 0$ is equal to $-\Psi^0(\lambda, \tau, z)$, cf. the definition in section 4. Hence, the singularity for $z \in U$ is dictated by

$$- \sum_{\substack{\lambda \in S_f(U, \epsilon) \\ \lambda \neq 0}} a^+(\lambda, -\frac{(\lambda, \lambda)}{2}) \Psi^0(\lambda, \tau, z).$$

In particular, z is a singular point precisely if $R(\lambda, z) = -(\lambda_z, \lambda_z) = 0$ for some $\lambda \in S_f(U, \epsilon) - \{0\}$. \square

The singular theta lift as a current Using the relationship between the singular theta lift and the singular Schwartz form Ψ , already seen in the proof of Proposition 5.4, we derive a current equation for $\Phi(f, \psi)$. The role of φ_{KM} in Theorem 4.3 is now played by

$$\Lambda_\psi(f) := dd^c \Phi(z, f, \psi), \quad (5.5)$$

where $f \in \mathbf{H}_{k, L^-}^+$.

Theorem 5.5. *The singular theta lift $\Phi(z, f, \psi)$ and the lifting $\Lambda_\psi(f)$ satisfy the following current equation on X :*

$$dd^c[\Phi(f, \psi)] + (-i)^q \delta_{Z(f)} = [\Lambda_\psi(f)].$$

Proof. This follows directly from Theorem 4.3. For $x \in V$, we have

$$dd^c[\Psi^0(x)] + (-i)^q \delta_{\Gamma(x) \setminus \mathbb{D}(x)} = [\varphi_{KM}^0(x)]. \quad (5.6)$$

As usual, denote the Fourier coefficients of f^+ by $a^+(\lambda, n)$ for $\lambda \in L^\sharp$, $n \in \mathbb{Q}$. For any relatively compact open neighbourhood $U \subset \mathbb{D}$ and any $\epsilon > 0$, we consider the set $S_f(U, \epsilon)$ from p. 22. Then, from the left hand side of (5.6), we get

$$dd^c \sum_{\substack{\lambda \in S_f(U, \epsilon) \\ \lambda \neq 0}} a^+(\lambda, -\frac{(\lambda, \lambda)}{2}) [\Psi^0(\lambda)] + (-i)^q \sum_{\substack{\lambda \in S_f(U, \epsilon) \\ \lambda \neq 0}} a^+(\lambda, -\frac{(\lambda, \lambda)}{2}) \delta_{Z(\lambda)}.$$

Now, by Proposition 5.4, and after taking the (locally finite) union over neighbourhoods U containing singular points, we get the current associated to (the singular part of) $\Phi(z, f, \psi)$ plus the delta current for the cycle $Z(f)$:

$$dd^c[\Phi(f, \psi)] + (-i)^q \delta_{Z(f)}.$$

(Note that, through Stokes' theorem, the current is determined by the singular part.)

Repeating the same steps on the right hand side of (5.6), by using the identity $dd^c \Psi(x, \tau, z) = \varphi_{KM}(x, \tau, z)$ (see Proposition 4.2), we recover the current

$$[dd^c \Phi(f, \psi)] = [\Lambda_\psi(f)],$$

as claimed. □

5.4. Adjointness to the Kudla-Millson lift

We now show an adjointness result analogous to [6, Theorem 6.1, Theorem 6.2].

Denote by $\Theta(\tau, z, \varphi_{KM})$ the theta function for the Schwartz form φ_{KM} from Section 3 (see, [26, 27, 28]). By Proposition 3.3 it is a closed differential (q, q) -form (in z), which has weight $p + q$ as a modular form (in τ). The Kudla-Millson lift Λ_{KM} is now defined for any rapidly decreasing $2(p - 1)q$ -form η through the assignment

$$\eta \longmapsto \Lambda_{KM}(\eta) := \int_X \eta \wedge \Theta(\tau, z, \varphi_{KM}).$$

This map factors through the de Rham cohomology with compact supports on X . By [28, Theorem 2] if η is closed, $\Lambda(\tau, \eta)$ is a holomorphic modular form.

To facilitate notation, we introduce a pairing $\{\cdot, \cdot\}'$ between the spaces M_{k,L^-} and $H_{k,L}^+$ see [6, (3.15) on p. 62]. Let $f \in H_{k,L}^+$ with $f^+ = \sum_{h,n} a^+(h,n)e(n\tau)\mathbf{e}_h$ and $h \in M_{k,L^-}$ with q -expansion $h = \sum_{h,n} b(h,n)e(n\tau)\mathbf{e}_h$. We set

$$\{h, f\}' := (h, \xi_k(f))_{2-k,L} - \sum_{h \in L^\sharp/L} a^+(h,0)b(h,0) = \sum_{h \in L^\sharp/L} \sum_{\substack{n \in \mathbb{Q} \\ n < 0}} a^+(h,n)b(h,-n).$$

Theorem 5.6. *The lift Λ_ψ has the following properties:*

1. *Let $f \in H_{k,L^-}^+$. Then*

$$(\Theta(\cdot, z, \varphi_{KM}), \xi_k(f))_{2-k,L} + a^+(0,0)\varphi_{KM}(0) = \Lambda_\psi(f)$$

as differential forms on X . In particular, $\Lambda_\psi(f)$ extends to a smooth closed (q, q) -form of moderate growth.

2. *The Kudla-Millson lift Λ_{KM} and Λ_ψ are adjoint in the sense that*

$$(\eta, \Lambda_\psi(f))_X = \{\Lambda_{KM}(\eta), f\}'$$

for any $f \in H_{k,L^-}^+$ and any rapidly decreasing closed $2(p-1)q$ -form η .

We note that, in particular, if $f \in M_{k,L^-}^!$, we have $\Lambda_\psi(f) = a^+(0,0)\varphi_{KM}(0)$.

Corollary 5.7. *For any rapidly decreasing closed $2(p-1)q$ -form η and any $f \in H_{k,\omega_L}$, we have*

$$(\eta, \Lambda_\psi(f))_X = \int_{Z(f)} \eta.$$

Proof of the Theorem. 1. We have

$$L_{2-k}\Theta(\tau, z, \varphi_{KM}) = \Theta(\tau, z, dd^c\psi),$$

since $L\varphi_{KM}(0) = dd^c\psi(0)$, Hence, we have

$$\begin{aligned} \lim_{t \rightarrow \infty} \int_{\mathcal{F}_t} \langle L_{2-k}\Theta(\tau, z, \varphi_{KM}), \bar{f} \rangle d\mu &= \int_{\mathcal{F}}^{reg} \langle L_{2-k}\Theta(\tau, z, \varphi_{KM}), \bar{f} \rangle d\mu \\ &= \int_{\mathcal{F}}^{reg} \langle \Theta(\tau, z, dd^c\psi), \bar{f} \rangle d\mu, \end{aligned}$$

and this quantity defines a smooth form on $\mathbb{D} - \mathbb{D}(f)$, which extends smoothly to \mathbb{D} . With [6, Lemmas 6.6, 6.7] we get the following identity, valid outside $\mathbb{D}(f)$:

$$(\Theta(z, \varphi_{KM}), \xi_k(f))_{2-k,L} = \int_{\mathcal{F}}^{reg} \langle \Theta(\tau, z, dd^c\psi), \bar{f} \rangle d\mu + a^+(0,0)\varphi_{KM}(0).$$

Now, the statement follows by showing that

$$\int_{\mathcal{F}}^{\text{reg}} \langle \Theta(\tau, z, dd^c \psi), \bar{f} \rangle d\mu = dd^c \int_{\mathcal{F}}^{\text{reg}} \langle \Theta(\tau, z, \psi), \bar{f} \rangle d\mu. \quad (5.7)$$

First, note that

$$\int_{\mathcal{F}}^{\text{reg}} \langle \Theta(\tau, z, \psi), \bar{f} \rangle d\mu = \lim_{t \rightarrow \infty} \int_{\mathcal{F}_t} (\langle \Theta(\tau, z, \psi), \bar{f} \rangle - a^+(0, 0)v) d\mu + Ca^+(0, 0), \quad (5.8)$$

with a constant C , coming from the regularisation of the constant term. Arguing along the same lines as in the proof of Proposition 5.4, we see that in the integrand, the sum over $\lambda \in L^\sharp - S_f(U, \epsilon)$ (see p. 22) converges uniformly for any relatively compact open neighbourhood $U \subset \mathbb{D}$ and any $\epsilon > 0$. For the remaining terms, with $\lambda \in S_f(U, \epsilon)$ the integrand decays exponentially.

Thus, switching the order of differentiation from the right hand side of (5.7) and the limit from (5.8) is justified, which completes the proof.

2. The second statement follows from the first, the proof is exactly like the one of [6, Theorem 6.3], which we briefly reproduce here. Denote by $(\cdot, \cdot)_X$ the natural pairing between closed forms of complementary degree (where one is rapidly decreasing and the other of moderate growth). We have

$$\begin{aligned} (\eta, \Lambda_\psi(f))_X &= (\eta, (\Theta(\cdot, z, \varphi_{KM}), \xi_k(f))_{k,L})_X \\ &= ((\eta, \Theta(\cdot, z, \varphi_{KM}))_X, \xi_k(f))_{k,L} = \{\Lambda_{KM}(\eta), f\}. \end{aligned}$$

Note only that the order of integration can be switched by absolute convergence. \square

6. Comparison of the two Green currents

In this section, we want to compare the Green current of Kudla type $\mathcal{G}^K(m, w, h)$, for $m \in \mathbb{Q}$, $h \in L^\sharp/L$ and $w \in \mathbb{R}_{>0}$, and that of Bruinier type $\mathcal{G}^B(m, h)$ (see below). The aim is to transfer some of the results of Ehlen and Sankaran from [12] to the present setting.

6.1. Green currents of Bruinier type

We first introduce the Green currents of Bruinier type.

The Hejhal Poincaré series (also known as Maass-Poincaré series) of weight k of index (m, h) , $h \in L^\sharp/L$, $m \in \mathbb{Z}$ is defined as (for $\tau \in \mathbb{H}$, $s \in \mathbb{C}$ with $\sigma = \text{Re}(s) > 1$)

$$F_{m,h}(\tau, s) = \frac{1}{4\Gamma(2s)} \sum_{A \in \Gamma_\infty \backslash \text{SL}_2(\mathbb{Z})} \mathcal{M}_s(4\pi|m|v) e^{2\pi i m u} \mathbf{e}_h |_{k,L^-} A, \quad (6.1)$$

where $\mathcal{M}_s(t) = t^{-\frac{k}{2}} M_{-\frac{k}{2}, s-\frac{1}{2}}(t)$, with the M-Whittaker function $M_{\kappa, \mu}(t)$. Note that our definition of $F_{m,h}(\tau, s)$ differs from [4, Definition 1.8] by a factor of $\frac{1}{2}$.

Set $s_0 = 1 - \frac{k}{2}$. For fixed $s = s_0$, the Poincaré series $F_{m,h}(\tau, s_0)$ have principal part $q^m \mathbf{e}_h$ and form a basis of H_{k, L^-}^+ , [see 4, Proposition 1.12]. Note further that by [6, Remark 3.10] $\xi_k(F_{m,h}(\tau, s_0))$ is a holomorphic, cuspidal Poincaré series of index $(-m, h)$.

We now introduce two Green currents through the regularised pairing (see p. 21) of the Hejhal Poincaré series with $\Theta(\tau, z)$. First, we define the Bruinier type Green current $\mathcal{G}^B(m, h)$ by setting

$$\mathcal{G}^B(m, h)(z) := (F_{m,h}(\tau, s_0), \Theta(\cdot, z))_{L^-}^{reg}, \quad (6.2)$$

i.e., the regularised theta lift of the weak Maass form $F_{m,h}(\tau, s_0)$. By Theorem 5.5 $\mathcal{G}^B(m, h)$ is thus a Green current for the cycle $Z(m, h)$.

6.2. The Kudla type Green current as a theta lift

Following [12, Section 2.4], we introduce truncated Poincaré series $P_{m,w,h}$ with $m \in \mathbb{Z}$, $w \in \mathbb{R}_{>0}$ and $h \in L^\sharp/L$, of weight $k = 2 - (p + q)$:

$$P_{m,w,h}(\tau) = \frac{1}{2} \sum_{A \in \Gamma_\infty \backslash \mathrm{SL}_2(\mathbb{Z})} [\sigma_w(\tau) q^{-m} \mathbf{e}_h] |_{k, L^-} A,$$

$$\text{where } \sigma_w(\tau) = \begin{cases} 1 & \text{if } v \geq w \\ 0 & \text{if } v < w. \end{cases}$$

Further, if $m \notin \frac{1}{2}(h, h) + \mathbb{Z}$ we set $P_{m,w,h} = 0$.

Proposition 6.1. *The regularised pairing $(P_{m,w,h}, \Theta(\cdot, z))_{L^-}^{reg}$ exists. On $\mathbb{D} \setminus \mathbb{D}(h, m)$, it satisfies the identity*

$$(P_{m,w,h}, \Theta(\cdot, z))_{L^-}^{reg} = -\Xi^K(m, w, h) - \delta_{m,0} \delta_{h,0} \psi(0) \log(w).$$

The Kudla type Green current $\Xi^K(m, w, h)$ can thus be expressed as a regularized theta lifting. This also affords an (albeit discontinuous) extension of $\Xi^K(m, w, h)$ to all \mathbb{D} .

Proof. Assume that $z \notin \mathbb{D}(m, h)$. We evaluate the regularized pairing by unfolding and see

$$\begin{aligned} (P_{m,w,h}, \Theta(\cdot, z))_{L^-}^{reg} &= \mathrm{CT} \lim_{s=0} \lim_{t \rightarrow \infty} \int_{\mathcal{F}_t - \mathcal{F}_w} \sum_{\substack{\lambda \in L+h \\ (\lambda, \lambda) = 2m}} q^{-m} \psi(\lambda) v^{-s} d\mu \\ &= \mathrm{CT} \int_w^\infty \sum_{\substack{\lambda \in L+h \\ (\lambda, \lambda) = 2m}} \psi^0(\sqrt{v}\lambda, z) v^{-s-1} dv. \end{aligned}$$

Now, for $m \neq 0$ this extends smoothly to the entire s -plane and for $s = 0$, we obtain

$$\sum_{\substack{\lambda \in L+h \\ (\lambda, \lambda) = 2m}} \Psi^0(\sqrt{v}\lambda, z) = -\Xi^K(m, w, h).$$

Similarly, for $m = 0$ we obtain $-\Xi^K(m, w, h)$ from the sum over $\lambda \neq 0$. The term for $\lambda = 0$ contributes

$$\psi(0) \text{CT} \lim_{s=0} \int_w^t v^{s-1} dv = -\psi(0) \text{CT} \lim_{s=0} \frac{1}{s} (t^{-s} - w^{-s}) = -\psi(0) \log(w). \quad \square$$

6.3. The difference of the two Green currents as a modular form

Now, with the results of [12], we can show that the difference of $\mathcal{G}^K(m, v)$ and $\mathcal{G}^B(m)$ is, essentially a modular form.

Lemma 6.2. *The difference*

$$(P_{m,w,h}, \Theta(\cdot, z))_{L^-}^{reg} - (F_{m,h}, \Theta(\cdot, z))_{L^-}^{reg}$$

extends to a smooth differential $(q-1, q-1)$ -form on \mathbb{D} .

Proof. Since the principal part of $F_{m,h}$ is given by $q^{-m} \mathbf{e}_h$ this is immediate from Proposition 5.4 and Proposition 6.1. \square

Ehlen and Sankaran have shown [12, Proposition 2.12] that for every $f = \sum_{h \in L^\sharp/L} f_h \mathbf{e}_h$ in $A_{\kappa-2, L}^{mod}$, there is unique element of $A_{\kappa, L}^!$ denoted $L^\sharp(f)$, with the following properties (we are mainly interested in the first two):

1. $L^\sharp(f)$ is a preimage of f under the lowering operator, i.e. $L_\kappa(L^\sharp(f)) = f$.
2. It has trivial cuspidal holomorphic projection, i.e. for every cusp form G in $S_{\kappa, L}$, the (regularised) Petersson product $\langle L^\sharp(f), G \rangle^{reg}$ vanishes.
3. Its principal part is in the image of η , a splitting [see 12, Section 2.3] of the map $\text{Sing}_{\kappa, L^-} \rightarrow M_{2-\kappa, L}^\vee$ introduced by Borcherds [3, Theorem 3.1].

Now, with the following result from [12] we will be able show that the difference of the two Green functions is a modular form contained in $A_{p+q, L}^!$ and which is equal to $L^\sharp(\Theta)$.

Theorem ([12] Theorem 2.14). *Suppose $f = \sum_{h \in L^\sharp/L} f_h \mathbf{e}_h \in A_{\kappa-2, L}^{mod}$ and $F = L^\sharp(f) \in A_{\kappa, L}^!$. Then for any $h \in L^\sharp/L$ and $m \in \mathbb{Q}$,*

$$(P_{m,v,h} - F_{m,h}, f)_{L^-}^{reg} = -c_F(m, h).$$

Theorem 6.3. *Fix $z \in \mathbb{D}$. The generating series*

$$F(\tau, z) = -\log(v) \psi(0) \mathbf{e}_0 - \sum_{m \in \mathbb{Q}} (\Xi^K(m, v) - \mathcal{G}^B(m)) (z) q^m$$

is an element of $A_{p+q, L}^!$. Furthermore, F satisfies $L_{p+q}(F)(\tau, z) = -\Theta(\tau, z)$ and is orthogonal to cusp forms.

Proof. We observe that $\Theta(\tau, z; \psi)$, as a function on \mathbb{H} is contained in the space $A_{(p+q-2), L}^{mod}$ see Definition 5.2. Clearly by Proposition 6.1 and using the above theorem from [12] the above generating series can be written as

$$\sum_{m \in \mathbb{Q}} \sum_{h \in L^\sharp/L} (P_{m,v,h} - F_{m,h}, \Theta_h(\cdot, z))_{L^-}^{reg} q^m \mathbf{e}_h = - \sum_{m \in \mathbb{Q}} \sum_{h \in L^\sharp/L} c_{L^\sharp(\Theta)}(m, h) q^m \mathbf{e}_h.$$

Hence, the generating series, as a function on \mathbb{H} , is the q -expansion of the modular form $F = L^\sharp(\Theta) \in A_{p+q, L}^!$. The statement on lowering and the orthogonality to cusp forms thus follow from the properties of $L^\sharp(\Theta)$ in [12, Proposition 2.12], see above. \square

7. Poincaré series

In this section we introduce and study the current $\mathcal{G}_s^B(m, h)$ depending on a complex parameter s and identify it with the green current onstructed by Oda-Tsuzuki [30].

Namely, for $s \in \mathbb{C}$ with $\sigma > 1$, we define

$$\mathcal{G}_s^B(m, h)(z) := \lim_{t \rightarrow \infty} \int_{\mathcal{F}_t} \langle F_{m,h}(\tau, s), \Theta(\tau, z) \rangle_L d\mu$$

Similar to Section 2.2 in [4] it can be seen that the (regularized) integral converges for σ sufficiently large and can be analytically continued to the region $\sigma > 1$ with $s \neq s_0$.

Remark 7.1. We can also define $\mathcal{G}_{s_0}^B(m, h)(z)$ for $s = s_0$ as the constant term of the Laurent expansion of $\mathcal{G}_s^B(m, h)(z)$ at $s = s_0$. We note that $\mathcal{G}^B(m, h)$, see (6.2) and $\mathcal{G}_{s_0}^B(m, h)$ are not quite identical; due to the different regularization procedures, they differ by a smooth term. See [4, Proposition 2.11] for further details in the orthogonal case.

To ease the comparison with the work of Oda-Tsuzuki, we use the identification of differential forms on \mathbb{D} with K -invariant functions on G with values in $\bigwedge^\bullet \mathfrak{p}^*$. In our situation, this means to consider $\mathcal{G}_s^B(m, h)$ as a function on G with values in $\bigwedge^{q-1, q-1} \mathfrak{p}^*$ by first setting $\psi(x, g) := \psi(g^{-1}x, z_0)$ for $g \in G$ and then defining

$$\mathcal{G}_s^B(m, h)(g) := \lim_{t \rightarrow \infty} \int_{\mathcal{F}_t} \langle F_{m,h}(\tau, s), \Theta(\tau, g) \rangle_L d\mu.$$

It is then clear that $\mathcal{G}_s^B(m, h)$ is holomorphic in s in the convergent range.

7.1. An eigenvalue equation

Now, we show that the Green current $\mathcal{G}_s^B(m, h)$ satisfies an eigenvalue equation under the action of the Casimir element for $U(p, q)$ as the one in [30], Theorem 18 (iii) (with a different normalization of the holomorphic parameter s). The overall strategy follows of [4, Chapter 4.1] using results of Shintani [31] and additionally Hufner [20]. We denote

by C_{SL_2} , $C_{\mathrm{U}(p,q)}$ and $C_{\mathrm{O}(2p,2q)}$ the respective Casimir elements of $\mathrm{SL}_2(\mathbb{R})$, $\mathrm{U}(p, q)$ and $\mathrm{O}(2p, 2q)$ in the universal enveloping algebra.

Let $\phi = \phi(x, \tau, z_0)$ be a Schwartz form and κ the weight of $\phi(\tau)$ under the Weil representation. As ϕ satisfies condition (1.19) of [31] with $m = 2\kappa$, by [31, Lemma 1.4] we have

$$\begin{aligned}\omega(g'_\tau)C_{\mathrm{SL}_2}\phi(x) &= 4\left[v^2\left(\frac{\partial^2}{\partial^2u} + \frac{\partial^2}{\partial^2v}\right) - \kappa iv\frac{\partial}{\partial u}\right]\omega(g_\tau)\phi(x) \\ &= -4\left[\Delta_\kappa - v\kappa\frac{\partial}{\partial v}\right]\omega(g_\tau)\phi(x),\end{aligned}$$

wherein $g'_\tau = \begin{pmatrix} \sqrt{v} & u\sqrt{v}^{-1} \\ & \sqrt{v}^{-1} \end{pmatrix}$. By a brief calculation we thus have

$$4\Delta_\kappa\phi(x, \tau) = \kappa(\kappa - 2)\phi(x, \tau) - v^{-\frac{\kappa}{2}}\omega(g'_\tau)C_{\mathrm{SL}_2}\phi(x).$$

Now, by [31, Lemma 1.5] we have with $m = \dim_{\mathbb{C}}(V) = p + q$

$$C_{\mathrm{SL}_2}\phi(x) = [C_{\mathrm{O}(2p,2q)} + m(m - 2)]\phi(x).$$

We note that the operation of $\mathrm{SL}_2(\mathbb{R})$ through the Weil representation commutes with $C_{\mathrm{O}(2p,2q)}$. Hence, we get

$$4\Delta_\kappa\phi(x, \tau) = [\kappa(\kappa - 2) - m(m - 2)]\phi(x, \tau) - C_{\mathrm{O}(2p,2q)}\phi(x, \tau).$$

Now, by a result of Hufler [see 20, Satz 6.10], who in his thesis carries out the analogous computations for the Schwartz form φ_0 ,

$$C_{\mathrm{U}(p,q)}\phi(x) = C_{\mathrm{O}(2p,2q)}\phi(x) - 2\left(\Im\left(\sum_{j=1}^m z_j\frac{\partial}{\partial z_j}\right)\right)^2\phi(x). \quad (7.1)$$

Now set $\phi = \psi$. The second term on the right hand side of (7.1) vanishes for ψ and with $\kappa = p + q - 2 = m - 2$, we get

$$4\Delta_\kappa\psi = -4\kappa\psi - C_{\mathrm{U}(p,q)}\psi.$$

The following Lemma is an immediate consequence.

Lemma 7.2. *The theta function $\Theta(\tau, z)$, considered as a function on \mathbb{H} , satisfies the following differential equation:*

$$4\Delta_\kappa\Theta(\tau, z_0) = [-4\kappa - C_{\mathrm{U}(p,q)}]\Theta(\tau, z_0).$$

Noting that the Poincaré series $F_{m,h}$ is an eigenfunction of Δ_k with eigenvalue $\frac{\kappa^2}{4} + \frac{\kappa}{2} + s(1 - s)$ [see 4, p. 29], we have the following analogue of [4, Lemma 4.4], the proof of which is quite similar:

Lemma 7.3. *The regularised pairing of $\Theta(\tau, z)$ and the Maass Poincaré series $F_{m,h}$ of weight $-\kappa$, we have*

$$\begin{aligned} (F_{m,h}, \Delta_\kappa \Theta(\cdot, z))_{L^-}^{reg} &= (\Delta_{-\kappa} F_{h,m}, \Theta(\cdot, z))_{L^-}^{reg} - \kappa (F_{h,m}, \Theta(\cdot, z))_{L^-}^{reg} \\ &= \left(\frac{\kappa^2}{4} - \frac{\kappa}{2} + s(1-s) \right) (F_{h,m}, \Theta(\cdot, z))_{L^-}^{reg}. \end{aligned}$$

By combining the two Lemmas we get

Theorem 7.4. *Recall $\kappa = p + q - 2$. The Green current $\mathcal{G}_s^B(h, m)$ is an eigenfunction of the Casimir operator $C_{U(p,q)}$, with*

$$C_{U(p,q)} \mathcal{G}_s^B(m, h) = ((2s-1)^2 - (\kappa+1)^2) \mathcal{G}_s^B(m, h). \quad (7.2)$$

Proof. Due to locally uniform convergence of the regularized lift and all partial derivatives, we have

$$\begin{aligned} C_{U(p,q)} (F_{m,h}(\cdot, s), \Theta(\cdot, z))_{L^-}^{reg} &= (F_{m,h}(\cdot, s), C_{U(p,q)} \Theta(\cdot, z))_{L^-}^{reg} \\ &= -4 (F_{m,h}(\cdot, s), (\Delta_\kappa \Theta)(\cdot, z))_{L^-}^{reg} - 4k (F_{m,h}(\cdot, s), \Theta(\cdot, z))_{L^-}^{reg}, \end{aligned}$$

by Lemma 7.2. The statement then follows by Lemma 7.3. \square

7.2. Unfolding against the Poincaré series

In this section, we calculate $\mathcal{G}^B(m, h)(z_0)$ by unfolding the theta integral against the Poincaré series $F_{m,h}(\tau, s)$.

To facilitate notation, recall that we have defined

$$\mathbf{P}_{2\ell}^\psi(\lambda) := \frac{2i(-1)^{q-1}}{2^{2(q-1)}} \sum_{\alpha, \beta} P_{\alpha, \beta; 2\ell}^{2q-2}(\lambda) \otimes \Omega_{q-1}(\underline{\alpha}; \underline{\beta}). \quad (7.3)$$

Theorem 7.5. *We have*

$$\begin{aligned} \mathcal{G}_s^B(m, h) &= \frac{(4\pi|m|)^{s-\frac{k}{2}}}{2\Gamma(2s)} \\ &\times \sum_{\substack{\lambda \in h+L \\ (\lambda, \lambda) = 2m}} \sum_{\ell=0}^{q-1} \mathbf{P}_{2\ell}^\psi(\lambda) \frac{\Gamma(s - \frac{k}{2} + \ell)}{\left(2\pi(\lambda_{z_0^\perp}, \lambda_{z_0^\perp})\right)^{s-\frac{k}{2}+\ell}} {}_2F_1\left(s - \frac{k}{2} + \ell, s + \frac{k}{2}; 2s; \frac{2|m|}{(\lambda_{z_0^\perp}, \lambda_{z_0^\perp})}\right). \end{aligned}$$

Here ${}_2F_1$ denotes the standard Gaussian hypergeometric function.

Proof. From the definition of $F_{m,h}$ (6.1) and the Petersson operation, we have

$$\begin{aligned} \mathcal{G}_s^B(m, h) &= \\ &= \frac{1}{4\Gamma(2s)} \int_{\mathcal{F}}^{reg} \sum_{A \in \Gamma_\infty \backslash \mathrm{SL}_2(\mathbb{Z})} \mathcal{M}_s(4\pi|m|\Im(A\tau)) e^{2\pi i m \operatorname{Re}(A\tau)} j(A, \tau)^{-k} \theta_h(\tau, z_0) d\mu \\ &= \frac{1}{4\Gamma(2s)} \int_{\mathcal{F}}^{reg} \sum_{A \in \Gamma_\infty \backslash \mathrm{SL}_2(\mathbb{Z})} \mathcal{M}_s(4\pi|m|\Im(A\tau)) e^{2\pi i m \operatorname{Re}(A\tau)} \theta_h(A\tau, z_0) d\mu, \end{aligned}$$

since $\theta_h(\tau, z)$ is modular of weight $-k$ for $\mathrm{SL}_2(\mathbb{Z})$. Now, arguing exactly as in [4, p.55f], the unfolding (justified by absolute convergence for $\sigma = \mathrm{Re}(s) > 1 + \frac{p}{2} + \frac{q}{2}$) is completed and we obtain

$$\mathcal{G}_s^B(m, h) = \frac{2}{4\Gamma(2s)} \int_{v=0}^{\infty} \int_{u=0}^1 \mathcal{M}_s(4\pi|m|v) e^{2\pi i m u} \theta_h(\tau, z_0) v^{-2} du dv.$$

Inserting the Fourier expansion of $\theta_h(\tau, z)$ and integrating over u one sees

$$\begin{aligned} & \frac{(4\pi|m|)^{-\frac{k}{2}}}{2\Gamma(2s)} \int_{v=0}^{\infty} \sum_{\substack{\lambda \in h+L \\ (\lambda, \lambda) = -2m}} M_{-\frac{k}{2}, s-\frac{1}{2}}(4\pi|m|v) e^{2\pi(\lambda_{z_0}, \lambda_{z_0})v - \pi(\lambda, \lambda)v} v^{-\frac{k}{2}-1} \sum_{\ell=0}^{q-1} v^\ell \mathbf{P}_{2\ell}^\psi(\lambda) \\ &= \frac{(4\pi|m|)^{-\frac{k}{2}}}{2\Gamma(2s)} \sum_{\substack{\lambda \in h+L \\ (\lambda, \lambda) = -2m}} \sum_{\ell=0}^{q-1} \mathbf{P}_{2\ell}^\psi(\lambda) \int_{v=0}^{\infty} v^{-\frac{k}{2}+\ell-1} M_{-\frac{k}{2}, s-\frac{1}{2}}(4\pi|m|v) e^{-\pi v(\lambda, \lambda)_{z_0}} dv. \end{aligned}$$

The integrals are Laplace transforms, which can be evaluated as usual [see 13, p. 215]. We get for each term

$$\frac{(4\pi|m|)^s}{\left(2\pi(\lambda_{z_0^\perp}, \lambda_{z_0^\perp})\right)^{s-\frac{k}{2}+\ell}} \Gamma\left(s - \frac{k}{2} + \ell\right) {}_2F_1\left(s - \frac{k}{2} + \ell, s + \frac{k}{2}; 2s; \frac{2|m|}{(\lambda_{z_0^\perp}, \lambda_{z_0^\perp})}\right). \quad \square$$

We denote the individual summands for $\mathcal{G}_s^B(h, m)(g)$ in Theorem 7.5 by $\phi_s(\lambda, g)$.

Proposition 7.6. *Assume $m > 0$. Let H be the stabilizer of λ in G . Then*

(i)

$$\phi_s(\lambda) \in C^\infty\left((G - HK)/K; \bigwedge^{(q-1), (q-1)} \mathfrak{p}^*\right)$$

(ii) $\phi_s(\lambda)$ is holomorphic in s .

(iii) Let $\lambda = \sqrt{2m}v_1$ and consider $g = a_t = \exp(tX_{1p+q})$ as in the proof of Proposition 4.5. Then there exists a non-zero constant C such that

$$\lim_{t \rightarrow 0} t^{2(q-1)} \phi_s(\lambda, a_t) = C \Omega_{q-1}(\underline{1}, \underline{1}).$$

(iv) With the hypothesis as in (iv) we have

$$\phi_s(\lambda, a_t) = O(e^{-(2\mathrm{Re}(s)+p+q)t})$$

as $t \rightarrow \infty$.

Proof. (i) and (ii) are clear. Now assume $\lambda = \sqrt{2mv_1}$ and take $g = a_t = \exp(tX_{1p+q})$. Then $a_t^{-1}\lambda_{z_0} = \cosh(t)\sqrt{2mv_1}$, and we calculate

$$\begin{aligned} \phi_s(\lambda, a_t) &= \frac{1}{2\Gamma(2s)} \sum_{\ell=0}^{q-1} \mathbf{P}_{2\ell}^\psi(\cosh(t)\sqrt{2mv_1}) \\ &\quad \cdot \frac{\Gamma(s - \frac{k}{2} + \ell)}{(4\pi m)^\ell (\cosh^2 t)^{2s-k+2\ell}} {}_2F_1\left(s - \frac{k}{2} + \ell, s + \frac{k}{2}; 2s; \frac{1}{\cosh^2 t}\right) \\ &= \frac{1}{2\Gamma(2s)} \sum_{\ell=0}^{q-1} \mathbf{P}_{2\ell}^\psi(v_1) \frac{\Gamma(s - \frac{k}{2} + \ell)}{(2\pi)^\ell (\cosh t)^{2s-k}} {}_2F_1\left(s - \frac{k}{2} + \ell, s + \frac{k}{2}; 2s; \frac{1}{\cosh^2 t}\right) \\ &= \frac{1}{2\Gamma(2s)} \sum_{\ell=0}^{q-1} \mathbf{P}_{2\ell}^\psi(v_1) \frac{\Gamma(s - \frac{k}{2} + \ell)}{(2\pi)^\ell (\cosh t)^{2s-k}} \left(\frac{\sinh t}{\cosh t}\right)^{-2\ell} {}_2F_1\left(s + \frac{k}{2} - \ell, s - \frac{k}{2}; 2s; \frac{1}{\cosh^2 t}\right). \end{aligned}$$

Here we used ${}_2F_1(a, b; c, z) = (1-z)^{c-a-b} {}_2F_1(c-a, b-a; c, z)$. Then (iii) follows from the second line of the previous equation, while (iv) from the third line, properties of $\mathbf{P}_{2\ell}^\psi(v_1)$ and ${}_2F_1(s + \frac{k}{2} - (q-1), s - \frac{k}{2}; 2s; 1) = \Gamma(2s)\Gamma(q-1)/\Gamma(s - \frac{k}{2} + q - 1)\Gamma(s + \frac{k}{2})$. \square

Oda and Tsuzuki in [30], Theorem 18, show that the properties (i)-(iv) in Theorem 7.6 together with the Casimir equation uniquely determine the function. Using Theorem 7.4 we conclude

Corollary 7.7. *The Green forms $\mathcal{G}_s^B(m, h)$ agree (up to a constant) with the (global) Green forms constructed by Oda and Tsuzuki in [30].*

Remark 7.8. Similarly one can evaluate the regularized pairing of $\Theta(\tau, z)$ with the non-holomorphic Eisenstein series

$$E_h(\tau, s) = \sum_{A \in \Gamma_\infty \backslash \mathrm{SL}_2(\mathbb{Z})} v^s \mathbf{e}_h |_{k, L^-} A,$$

corresponding to $\mathcal{G}_s^B(0, h)$. After unfolding, and integration one has

$$(E_h(\cdot, s), \Theta(\cdot, z))_{L^-}^{reg} |_{z=z_0} = 2 \sum_{\ell=0}^{q-1} \frac{\Gamma(s+\ell)}{(2\pi)^{s+\ell}} \sum_{\substack{\lambda \in L+h \\ (\lambda, \lambda)=0}} (\lambda_{z_0}, \lambda_{z_0})^{-s-\ell} \mathbf{P}_{2\ell}^\psi(\lambda).$$

This expression can be written in terms Eisenstein series for the discriminant kernel $G(L)$ in $U(V)$. After setting

$$\zeta_{h, \lambda}(s) := \sum_{\substack{a \in \mathcal{O}_F^\times \\ a\lambda \in L+h}} N_{F/\mathbb{Q}}(a)^{-s}, \quad P(L) = \{\lambda \in L^\sharp; \lambda \text{ primitive}, (\lambda, \lambda) = 0\},$$

one obtains

$$2 \sum_{\ell=0}^{q-1} \frac{\Gamma(s+\ell)}{|\mathcal{O}_F^\times| (2\pi)^{s+\ell}} \sum_{\lambda \in G(L) \setminus P(L)} \zeta_{h, \lambda}(s) \mathbf{P}_{2\ell}^\psi(\lambda) \sum_{\gamma \in G(L)_\lambda \setminus G(L)} (\lambda_{\gamma z_0}, \lambda)^{-s-\ell}.$$

A. Calculations in the Fock model

In this section, we prove the main properties of the Schwartz functions from section 3. We use the polynomial Fock model for the Weil representation, the setup of which is reviewed in section B. We use the intertwining map $\iota : \mathcal{S}(V) \rightarrow \mathcal{P}(\mathbb{C}^{2(p+q)})$ between the Schrödinger model and the space of complex polynomials in $2(p+q)$ variables, on which action of the Weil representation ω is given by the Fock model. Note that $\iota(\varphi_0) = 1$. Further main properties of the intertwining operator are summarized in Lemma B.3.

We abbreviate the variables in Fock model for $U(p, q) \times U(1, 1)$ by $z''_\alpha = z''_{\alpha 1}$; $z'_\alpha = z'_{\alpha 2}$; $z'_\mu = z'_{\mu 1}$; $z''_\mu = z''_{\mu 2}$. We then have (see Lemma B.3):

$$\mathcal{D} = \frac{1}{2^{2q}} \left(\frac{-i}{\sqrt{2\pi}} \right)^q \prod_{\mu} \sum_{\alpha=1}^p z''_{\alpha} \otimes A_{\alpha\mu} \quad \text{and} \quad \bar{\mathcal{D}} = \frac{1}{2^{2q}} \left(\frac{-i}{\sqrt{2\pi}} \right)^q \prod_{\mu} \sum_{\beta=1}^p z'_{\beta} \otimes \bar{A}_{\beta\mu}.$$

By applying this to $1 \otimes 1 = \iota(\varphi_0 \otimes 1)$, we see that φ_{KM} is given by

$$\varphi_{KM} = \frac{(-1)^q}{2^{3q}\pi^{2q}} \sum_{\substack{\alpha_1, \dots, \alpha_q \\ \beta_1, \dots, \beta_q}} z''_{\alpha_1} \cdots z''_{\alpha_q} z'_{\beta_1} \cdots z'_{\beta_q} \otimes \Omega_q(\alpha_1, \dots, \alpha_q; \beta_1, \dots, \beta_q),$$

while the form ψ is given by

$$\psi = \frac{2i}{2^{3(q-1)}\pi^{2(q-1)}} \sum_{\substack{\alpha_1, \dots, \alpha_{q-1} \\ \beta_1, \dots, \beta_{q-1}}} z''_{\alpha_1} \cdots z''_{\alpha_{q-1}} z'_{\beta_1} \cdots z'_{\beta_{q-1}} \otimes \Omega_{q-1}(\alpha_1, \dots, \alpha_{q-1}; \beta_1, \dots, \beta_{q-1}).$$

A.1. Proof of Proposition 3.2

We first verify that ψ has the correct transformation behavior under the operation of $\mathfrak{k}' \simeq \mathfrak{so}_2(\mathbb{R})$.

Lemma A.1. *Under the operation of \mathfrak{k}' , the form ψ has weight $p+q-2$. That is,*

$$\omega \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \psi = i(p+q-2)\psi.$$

Proof. We use the formula for the operation of the generators of \mathfrak{k}' through the Weil representation from Lemma B.2 on p. 41, setting $r = s = 1$:

$$\begin{aligned} \omega(w_1 \circ w_1 + iw_1 \circ w_1 i) &= 2i \left[\sum_{\alpha=1}^p z''_{\alpha} \frac{\partial}{\partial z''_{\alpha}} - \sum_{\mu'=p+1}^{p+q} z'_{\mu'} \frac{\partial}{\partial z'_{\mu'}} \right] + i(p-q) \\ \text{and } \omega(w_2 \circ w_2 - iw_2 \circ w_2 i) &= 2i \left[\sum_{\alpha'=1}^p z'_{\alpha'} \frac{\partial}{\partial z'_{\alpha'}} - \sum_{\mu=p+1}^{p+q} z''_{\mu} \frac{\partial}{\partial z''_{\mu}} \right] + i(p-q). \end{aligned}$$

Note that, since $\Phi_W(iw \circ w) = 0$, this is actually the same as $\omega(w_1 \circ w_1)$ and $\omega(w_2 \circ w_2)$, respectively.

As $\mathfrak{su}(W) \simeq \mathfrak{sl}_2(\mathbb{R})$, we are mainly interested in the behaviour of ψ under the operation of $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ (while of course, $\begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix}$ generates the center). We have

$$\begin{aligned} \omega \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} &= \omega \left(\frac{1}{2} (w_1 \circ w_1 + w_2 \circ w_2) \right) \\ &= i \left[\sum_{\alpha=1}^p z''_{\alpha} \frac{\partial}{\partial z''_{\alpha}} + \sum_{\alpha'=1}^p z'_{\alpha'} \frac{\partial}{\partial z'_{\alpha'}} - \sum_{\mu'=p+1}^{p+q} z'_{\mu'} \frac{\partial}{\partial z'_{\mu'}} - \sum_{\mu=p+1}^{p+q} z''_{\mu} \frac{\partial}{\partial z''_{\mu}} \right] + i(p-q). \end{aligned}$$

Bearing in mind that ψ doesn't depend on $z'_{\mu'}$ and z''_{μ} the claim now follows from

$$\sum_{\alpha=1}^p z''_{\alpha} \frac{\partial}{\partial z''_{\alpha}} \psi = \sum_{\alpha'=1}^p z'_{\alpha'} \frac{\partial}{\partial z'_{\alpha'}} \psi = (q-1)\psi,$$

which is easily checked. \square

Lemma A.2. *The Schwartz form ψ is invariant under the operation of \mathfrak{k} .*

Proof. We need to show $Z(\psi) = 0$ for all $Z \in \mathfrak{k}$. Using the explicit formula for ψ given above (and ignoring constants), this means, using that Z acts as derivation,

$$\begin{aligned} 0 &= \sum_{\substack{\alpha_1, \dots, \alpha_{q-1} \\ \beta_1, \dots, \beta_{q-1}}} \omega(Z) \left(z''_{\alpha_1} \cdots z''_{\alpha_{q-1}} z'_{\beta_1} \cdots z'_{\beta_{q-1}} \right) \otimes \Omega_{q-1}(\alpha_1, \dots, \alpha_{q-1}; \beta_1, \dots, \beta_{q-1}) \\ &+ \sum_{\substack{\alpha_1, \dots, \alpha_{q-1} \\ \beta_1, \dots, \beta_{q-1}}} z''_{\alpha_1} \cdots z''_{\alpha_{q-1}} z'_{\beta_1} \cdots z'_{\beta_{q-1}} \otimes Z. (\Omega_{q-1}(\alpha_1, \dots, \alpha_{q-1}; \beta_1, \dots, \beta_{q-1})). \end{aligned}$$

Now let $Z = Z'_{\alpha\beta} \in \text{Hom}(V'_+, V'_+)$. Then the Weil representation action gives

$$\begin{aligned} &\omega(Z'_{\alpha\beta}) \left(z''_{\alpha_1} \cdots z''_{\alpha_{q-1}} z'_{\beta_1} \cdots z'_{\beta_{q-1}} \right) \otimes \Omega_{q-1}(\alpha_1, \dots, \alpha_{q-1}; \beta_1, \dots, \beta_{q-1}) \\ &= - \sum_{j=1}^{q-1} z''_{\alpha} z''_{\alpha_1} \cdots \widehat{z''_{\alpha_j}} \cdots z''_{\alpha_{q-1}} z'_{\beta_1} \cdots z'_{\beta_{q-1}} \otimes \Omega_{q-1}(\alpha_1, \dots, \beta, \dots, \alpha_{q-1}; \beta_1, \dots, \beta_{q-1}) \\ &+ \sum_{j=1}^{q-1} z''_{\alpha_1} \cdots z''_{\alpha_{q-1}} z'_{\beta} z'_{\beta_1} \cdots \widehat{z'_{\beta_j}} \cdots z'_{\beta_{q-1}} \otimes \Omega_{q-1}(\alpha_1, \dots, \alpha_{q-1}; \beta_1, \dots, \alpha, \dots, \beta_{q-1}). \end{aligned}$$

Now $\mathfrak{k} \simeq \text{Hom}(V'_+, V'_+)$ acts on $\mathfrak{p}_+ \simeq \text{Hom}(V_-, V_+)$ by composition. We obtain

$$Z'_{\alpha\beta} \cdot Z'_{\alpha_j\mu} = -\delta_{\beta\alpha_j} Z'_{\alpha\mu},$$

and hence for the dual action we see

$$Z'_{\alpha\beta} \cdot \xi'_{\alpha_j\mu} = \delta_{\alpha\alpha_j} \xi'_{\beta\mu}.$$

In the same way we see

$$Z'_{\alpha\beta} \cdot \xi''_{\beta_j\mu} = -\delta_{\beta\beta_j} \xi''_{\alpha\mu}.$$

This gives

$$\begin{aligned}
& z''_{\alpha_1} \cdots z''_{\alpha_{q-1}} z'_{\beta_1} \cdots z'_{\beta_{q-1}} \otimes Z'_{\alpha\beta} \Omega_{q-1}(\alpha_1, \dots, \alpha_{q-1}; \beta_1, \dots, \beta_{q-1}) \\
&= \sum_{j=1}^{q-1} z''_{\alpha_1} \cdots z''_{\alpha_j} \cdots z''_{\alpha_{q-1}} z'_{\beta_1} \cdots z'_{\beta_{q-1}} \Omega_{q-1}(\alpha_1, \dots, \beta, \dots, \alpha_{q-1}; \beta_1, \dots, \beta_{q-1}) \\
&= - \sum_{j=1}^{q-1} z''_{\alpha_1} \cdots z''_{\alpha_{q-1}} z'_{\alpha'_1} \cdots z'_{\beta} \cdots z'_{\alpha'_{q-1}} \Omega_{q-1}(\alpha_1, \dots, \alpha_{q-1}; \beta_1, \dots, \alpha, \dots, \beta_{q-1}).
\end{aligned}$$

Combining all this shows $Z'_{\alpha\beta}\psi = 0$, as desired.

We now consider the action of $Z'_{\mu\nu} \in \text{Hom}(V'_-, V'_-)$. The Weil representation action on ψ clearly vanishes. Now the action on \mathfrak{p}^+ is given by $Z'_{\mu\nu} \cdot Z'_{\alpha\nu} = \delta_{\mu\nu} Z'_{\alpha\nu}$ and hence

$$Z'_{\mu\nu} \xi'_{\alpha_j \mu'} = -\delta_{\nu\mu'} \xi'_{\alpha_j \mu} \quad \text{and} \quad Z'_{\mu\nu} \xi''_{\beta_j \mu'} = \delta_{\mu\mu'} \xi''_{\beta_j \nu}.$$

From this it is easy to see that

$$Z'_{\mu\nu} \Omega_{q-1}(\alpha_1, \dots, \alpha_{q-1}; \beta_1, \dots, \beta_{q-1}) = 0.$$

for all $\underline{\alpha}, \underline{\beta}$. □

A.2. Proof of Theorem 3.3

Recall

$$d = \frac{1}{2} (\partial + \bar{\partial}), \quad d^c = \frac{(\partial - \bar{\partial})}{4\pi i}, \quad dd^c = -\frac{1}{4\pi i} \partial \bar{\partial}.$$

In the Fock model, the differential operators $\partial, \bar{\partial}$ are given by (see Lemmas B.1, B.2)

$$\partial = \sum_{\alpha, \mu} \left[\frac{1}{4\pi} z''_{\alpha} z'_{\mu} - 4\pi \frac{\partial^2}{\partial z'_{\alpha} \partial z''_{\mu}} \right] \otimes A'_{\alpha\mu}, \quad \bar{\partial} = \sum_{\beta, \nu} \left[\frac{1}{4\pi} z'_{\beta} z''_{\nu} - 4\pi \frac{\partial^2}{\partial z''_{\beta} \partial z'_{\nu}} \right] \otimes A''_{\beta\nu}.$$

For the lowering operator $L = -\frac{i}{2} \omega(w_1 \circ w_2 + iw_1 \circ w_2 i)$, we have

$$L = -4\pi \sum_{\gamma} \frac{\partial^2}{\partial z''_{\gamma} \partial z'_{\gamma}} + \frac{1}{4\pi} \sum_{\mu} z''_{\mu} z'_{\mu}.$$

For simplicity we drop all constants and consider

$$\begin{aligned}
\varphi'_{KM} &= \sum_{\substack{\alpha_1, \dots, \alpha_q \\ \beta_1, \dots, \beta_q}} z''_{\alpha_1} \cdots z''_{\alpha_q} z'_{\beta_1} \cdots z'_{\beta_q} \otimes \xi'_{\alpha_1 p+1} \wedge \cdots \wedge \xi'_{\alpha_q p+q} \wedge \xi''_{\beta_1 p+1} \wedge \cdots \wedge \xi''_{\beta_q p+q}, \\
\psi' &= \sum_{\substack{\alpha_1, \dots, \alpha_{q-1} \\ \beta_1, \dots, \beta_{q-1}}} z''_{\alpha_1} \cdots z''_{\alpha_{q-1}} z'_{\beta_1} \cdots z'_{\beta_{q-1}} \\
&\quad \otimes \sum_{j=1}^q \xi'_{\alpha_1 p+1} \wedge \cdots \wedge \widehat{\xi'_{p+j}} \cdots \wedge \xi'_{\alpha_{q-1} p+q} \wedge \xi''_{\beta_1 p+1} \wedge \cdots \wedge \widehat{\xi''_{p+j}} \wedge \cdots \wedge \xi''_{\beta_{q-1} p+q}.
\end{aligned}$$

Then the claim is equivalent to

$$L\varphi'_{KM} = (-1)^{q-1}4\pi\partial\bar{\partial}\psi',$$

which we show by a direct calculation of both sides. We have

$$\begin{aligned} L\varphi'_{KM} &= \frac{1}{4\pi} \left(\sum_{\mu} z''_{\mu} z'_{\mu} \right) \varphi'_{KM} \\ &\quad - 4\pi \sum_{\underline{\alpha}, \underline{\beta}} \sum_{j,k=1}^q \delta_{\alpha_j \beta_k} z''_{\alpha_1} \cdots \widehat{z''_{\alpha_j}} \cdots z''_{\alpha_{q-1}} z'_{\beta_1} \cdots \widehat{z'_{\beta_k}} \cdots z'_{\beta_{q-1}} \\ &\quad \otimes \xi'_{\alpha_1 p+1} \wedge \cdots \wedge \xi'_{\alpha_j p+j} \wedge \cdots \wedge \xi'_{\alpha_q p+q} \wedge \xi''_{\beta_1 p+1} \wedge \cdots \wedge \xi''_{\beta_k p+k} \wedge \cdots \wedge \xi''_{\beta_q p+q}. \end{aligned}$$

On the other hand,

$$\begin{aligned} \partial\bar{\partial}\psi' &= \frac{1}{16\pi^2} \sum_{\alpha, \beta, \mu, \nu} (z''_{\alpha} z'_{\beta} z'_{\nu} z''_{\mu} \otimes \xi'_{\alpha\nu} \wedge \xi''_{\beta\mu}) \psi' \\ &\quad - \sum_{\substack{\alpha_1, \dots, \alpha_{q-1} \\ \beta_1, \dots, \beta_{q-1} \\ \alpha, \beta, \mu}} z''_{\alpha_1} \cdots z''_{\alpha_{q-1}} \frac{\partial}{\partial z'_{\alpha}} \left(z'_{\beta} z'_{\beta_1} \cdots z'_{\beta_{q-1}} \right) \\ &\quad \otimes \xi'_{\alpha\mu} \wedge \xi''_{\beta\mu} \wedge \sum_{j=1}^q \xi'_{\alpha_1 p+1} \wedge \cdots \wedge \widehat{\xi'_{p+j}} \cdots \wedge \xi'_{\alpha_{q-1} p+q} \wedge \xi''_{\beta_1 p+1} \wedge \cdots \wedge \widehat{\xi''_{p+j}} \cdots \wedge \xi''_{\beta_{q-1} p+q}. \end{aligned}$$

For the first term, it is easy to see that only the terms $\mu = \nu$ contribute and one obtains

$$(-1)^{q-1} \frac{1}{16\pi^2} \left(\sum_{\mu} z''_{\mu} z'_{\mu} \right) \varphi'_{KM}.$$

For the second, only terms $\mu = p + j$ contribute and one obtains

$$\begin{aligned} (-1)^q \sum_{\substack{\alpha_1, \dots, \alpha_{q-1} \\ \beta_1, \dots, \beta_{q-1} \\ \alpha_0, \beta_0}} z''_{\alpha_1} \cdots z''_{\alpha_{q-1}} \sum_{k=0}^{q-1} \delta_{\alpha_0 \beta_k} z'_{\beta_0} z'_{\beta_1} \cdots \widehat{z'_{\beta_k}} \cdots z'_{\beta_{q-1}} \\ \otimes \sum_{j=1}^q \xi'_{\alpha_1 p+1} \wedge \cdots \wedge \xi'_{\alpha_0 p+j} \cdots \wedge \xi'_{\alpha_{q-1} p+q} \wedge \xi''_{\beta_1 p+1} \wedge \cdots \wedge \xi''_{\beta_0 p+j} \wedge \cdots \wedge \xi''_{\beta_{q-1} p+q}. \end{aligned}$$

Now comparing the formulas for $L\varphi'_{KM}$ and $\partial\bar{\partial}\psi'$ now gives the claim.

A.3. The auxiliary form $d^c\psi$

We now give a more explicit description of $d^c\psi$. We have

$$\begin{aligned} \frac{1}{4\pi}\partial\psi &= \frac{i}{2^{3(q-1)}\pi^{2(q-1)}} \frac{1}{2\pi} \sum_{\substack{\underline{\alpha}, \underline{\beta} \\ \gamma, \underline{\mu}}} z''_{\gamma} z'_{\mu} z''_{\alpha} z'_{\beta} \otimes \xi'_{\gamma\mu} \wedge \Omega_{q-1}(\underline{\alpha}; \underline{\beta}) \\ &= \frac{i}{2^{3q-2}\pi^{2q-1}} \sum_{\substack{\gamma, \alpha_1, \dots, \alpha_{q-1} \\ \beta_1, \dots, \beta_{q-1}}} z''_{\gamma} z''_{\alpha_1} \cdots z''_{\alpha_{q-1}} z'_{\beta} \sum_{j=1}^q (-1)^{j-1} z'_{p+j} \\ &\quad \otimes \xi'_{\alpha_1 p+1} \wedge \cdots \wedge \xi'_{\gamma p+j} \cdots \wedge \xi'_{\alpha_{q-1} p+q} \wedge \xi''_{\beta_1 p+1} \wedge \cdots \wedge \widehat{\xi''_{p+j}} \cdots \wedge \xi''_{\beta_{q-1} p+q}. \end{aligned}$$

Similarly,

$$\begin{aligned} \frac{1}{4\pi}\bar{\partial}\psi &= \frac{i}{2^{3q-2}\pi^{2q-1}} \sum_{\substack{\alpha_1, \dots, \alpha_{q-1} \\ \gamma, \beta_1, \dots, \beta_{q-1}}} z''_{\alpha} z'_{\gamma} z'_{\beta_1} \cdots z'_{\beta_{q-1}} \sum_{j=1}^q (-1)^{q+j} z''_{p+j} \\ &\quad \otimes \xi'_{\alpha_1 p+1} \wedge \cdots \wedge \widehat{\xi'_{p+j}} \cdots \wedge \xi'_{\alpha_{q-1} p+q} \wedge \xi''_{\beta_1 p+1} \wedge \cdots \wedge \xi''_{\gamma p+j} \cdots \wedge \xi''_{\beta_{q-1} p+q}. \end{aligned}$$

Now, ψ' is the difference of these two terms.

Finally, we want give an explicit form of $d^c\psi$ in the Schrödinger model. (Note that $\mathcal{D}_{\mu}\varphi_0 = 2\bar{z}_{\mu}\varphi_0$.) We have

$$d^c\psi(x) = \frac{1}{2^{3q-1}\pi^{2q-1}} \left[\sum_{\substack{\underline{\alpha}, \underline{\beta} \\ \gamma}} \mathcal{D}_{\underline{\alpha}} \mathcal{D}_{\gamma} \bar{\mathcal{D}}_{\underline{\beta}} \varphi_0(x) \otimes Q'_{\underline{\alpha}, \gamma; \underline{\beta}}(x) - \sum_{\substack{\underline{\alpha}, \underline{\beta} \\ \gamma}} \mathcal{D}_{\underline{\alpha}} \bar{\mathcal{D}}_{\gamma} \bar{\mathcal{D}}_{\underline{\beta}} \varphi_0(x) \otimes Q''_{\underline{\alpha}; \underline{\beta}, \gamma}(x) \right]. \quad (\text{A.1})$$

Here $Q'_{\underline{\alpha}, \gamma; \underline{\beta}}(x)$ and $Q''_{\underline{\alpha}; \underline{\beta}, \gamma}(x)$ are given by

$$\begin{aligned} Q'_{\underline{\alpha}, \gamma; \underline{\beta}}(x) &= \sum_{j=1}^q (-1)^{j-1} z_{p+j} \otimes \xi'_{\alpha_1 p+1} \wedge \cdots \wedge \xi'_{\gamma p+j} \cdots \wedge \xi'_{\alpha_{q-1} p+q} \wedge \xi''_{\beta_1 p+1} \wedge \cdots \wedge \widehat{\xi''_{p+j}} \cdots \wedge \xi''_{\beta_{q-1} p+q} \\ Q''_{\underline{\alpha}; \underline{\beta}, \gamma}(x) &= \sum_{j=1}^q (-1)^{q+j} \bar{z}_{p+j} \otimes \xi'_{\alpha_1 p+1} \wedge \cdots \wedge \widehat{\xi'_{p+j}} \cdots \wedge \xi'_{\alpha_{q-1} p+q} \wedge \xi''_{\beta_1 p+1} \wedge \cdots \wedge \xi''_{\gamma p+j} \cdots \wedge \xi''_{\beta_{q-1} p+q}. \end{aligned}$$

B. The Fock model for unitary dual pairs

We review the Fock model of the Weil representation for the dual pair $U(p, q) \times U(r, s)$. We follow [1, 28, 26], see also [15].

B.1. The Fock model for the symplectic group

Let $(\mathbb{W}, \langle \cdot, \cdot \rangle)$ be a non-degenerate real symplectic space of dimension $2N$ and let J be a positive definite complex structure on W , i.e., the bilinear form given by $\langle w_1, Jw_2 \rangle$ is positive definite. Let $e_1, \dots, e_N; f_1, \dots, f_N$ be a standard symplectic basis of W so that $Je_j = f_j$ and $Jf_j = -e_j$. We decompose

$$\mathbb{W} \otimes \mathbb{C} = \mathbb{W}' \oplus \mathbb{W}''$$

into the $+i$ and $-i$ eigenspaces under J . Then $w'_j = e_j - f_j i$ and $w''_j = e_j + f_j i$ form a (symplectic) basis for \mathbb{W}' and \mathbb{W}'' respectively with $\langle w'_j, w''_k \rangle = 2i\delta_{jk}$.

We identify $\text{Sym}^\bullet(\mathbb{W}'')$ with the polynomial functions $\mathcal{P}(\mathbb{C}^N) = \mathbb{C}[z_1, \dots, z_N]$ on \mathbb{W}'' via $z_j(w''_k) = \langle w'_j, w''_k \rangle = 2i\delta_{jk}$. For $\lambda \in \mathbb{C}^*$, we define an action ρ_λ of \mathbb{W} on $\mathcal{P}(\mathbb{C}^N)$ by

$$\rho_\lambda(w''_j) = z_j \quad \text{and} \quad \rho_\lambda(w'_j) = 2i\lambda \frac{\partial}{\partial z_j},$$

which induces an action of the associated quantum algebra \mathbb{W}_λ . We identify $\text{Sym}^2(\mathbb{W})$ with $\mathfrak{sp}(\mathbb{W})$ via

$$(x \circ y)(z) = \langle x, z \rangle y + \langle y, z \rangle x.$$

Then the action ω_λ of $\mathfrak{sp}(\mathbb{W}) \otimes \mathbb{C}$ on $\mathcal{P}(\mathbb{C}^N)$ is given by

$$\omega_\lambda(x \circ y) = \frac{1}{2\lambda} (\rho_\lambda(x)\rho_\lambda(y) + \rho_\lambda(y)\rho_\lambda(x)). \quad (\text{B.1})$$

This is the Fock model of the Weil representation with central character λ .

B.2. Unitary dual pairs

Let $(V, (\cdot, \cdot))$ be a complex vector space of dimension $m = p + q$ with a non-degenerate Hermitian form (\cdot, \cdot) of signature (p, q) . Recall that we assume that (\cdot, \cdot) is \mathbb{C} -linear in the second and \mathbb{C} -antilinear in the first variable. We pick standard orthogonal basis elements v_α ($\alpha = 1, \dots, p$) and v_μ ($\mu = p + 1, \dots, m$) of length 1 and -1 respectively. We let θ be the Cartan involution with respect to this chosen basis of V and obtain a decomposition $V = V_+ \oplus V_-$. Let $G = \text{U}(V) \simeq \text{U}(p, q)$ and let $\mathfrak{g}_0 = \mathfrak{u}(V) \simeq \mathfrak{u}(p, q)$. We write $\mathfrak{g} = \mathfrak{g}_0 \otimes \mathbb{C}$ for the complexification of \mathfrak{g}_0 , viewed as a right \mathbb{C} vector space.

We let $(W, \langle \cdot, \cdot \rangle)$ be a complex vector space with a non-degenerate skew-Hermitian form $\langle \cdot, \cdot \rangle$ of signature (r, s) , again \mathbb{C} -linear in the second and \mathbb{C} -antilinear in the first variable. We pick an ‘‘orthogonal’’ basis w_a ($a = 1, \dots, r$) and w_u ($u = r + 1, \dots, r + s$) such that $\langle w_a, w_a \rangle = i$ and $\langle w_u, w_u \rangle = -i$. We obtain a decomposition $W = W_+ \oplus W_-$. We let J_0 be a positive definite complex structure with respect to this decomposition of W ; that is, J_0 acts by multiplication with $-i$ on W_+ and with i on W_- . We let $G' = \text{U}(W) \simeq \text{U}(r, s)$ and let $\mathfrak{g}'_0 = \mathfrak{u}(W) \simeq \mathfrak{u}(r, s)$. The two factors of the maximal compact subgroup $K' \simeq \text{U}(r) \times \text{U}(s)$ of G' act on the subspaces W_+ and W_- respectively. We let $W_{\mathbb{C}} = W \otimes_{\mathbb{R}} \mathbb{C}$ be the complexification of W , which we again view as a right

\mathbb{C} -vectorspace. We write $\mathfrak{g}' = \mathfrak{g}'_0 \otimes \mathbb{C}$. Then the $+i$ -eigenspace W' and the $-i$ eigenspace W'' of J_0 are spanned by

$$\begin{aligned} w'_a &:= w_a + iw_a i & w'_u &:= w_u - iw_u i, \\ w''_a &:= w_a - iw_a i & w''_u &:= w_u + iw_u i. \end{aligned}$$

Similarly as for V , we define the \mathbb{R} -linear surjective map by

$$\phi_W(w \circ \tilde{w})(x) = \langle w, x \rangle \tilde{w} + \langle \tilde{w}, x \rangle w.$$

Note that we have $\phi_W(iw \circ \tilde{w}) = \phi_W(w \circ -i\tilde{w})$. As for V , we will drop ϕ_W and just write $w \circ \tilde{w} \in \mathfrak{u}(W)$. In the standard decomposition

$$\mathfrak{g}' = \mathfrak{k}' \oplus \mathfrak{p}'^+ \oplus \mathfrak{p}'^-$$

we have

$$\begin{aligned} \mathfrak{k}' &= \text{span}_{\mathbb{C}}\{w_a \circ w_b + iw_a \circ w_b i\} \oplus \text{span}_{\mathbb{C}}\{w_u \circ w_v + iw_u \circ w_v i\}, \\ \mathfrak{p}'^+ &= \text{span}_{\mathbb{C}}\{w_a \circ w_u - iw_a \circ w_u i\}, \\ \mathfrak{p}'^- &= \text{span}_{\mathbb{C}}\{w_a \circ w_u + iw_a \circ w_u i\}. \end{aligned}$$

Note that for $r = s = n$, W is the split skew-Hermitian space of dimension $2n$, and we obtain a (symplectic) basis $e_j := w_j + w_{n+j}$, $f_j := -i(w_j - w_{n+j})$ of W . (Note that $\langle e_j, f_j \rangle = 2$). Then the positive definite almost complex structure is the usual one given by $J_0 e_j = f_j$ and $J_0 f_j = -e_j$. For $n = 1$, we have $\mathfrak{su}(W) \simeq \mathfrak{sl}_2(\mathbb{R})$, and the isomorphism is realized by switching to the symplectic basis above. Then for \mathfrak{k}' we see

$$\frac{1}{2}(w_1 \circ w_1 + w_2 \circ w_2) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad \text{and} \quad \frac{1}{2}(w_1 \circ w_1 - w_2 \circ w_2) = \begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix}.$$

Note that $\phi_W(iw \circ w) = 0$. Furthermore,

$$\begin{aligned} L &:= \frac{1}{2} \begin{pmatrix} 1 & -i \\ -i & -1 \end{pmatrix} = \frac{-i}{2}[w_1 \circ w_2 + iw_1 \circ w_2 i] \in \mathfrak{p}'^-, \\ R &:= \frac{1}{2} \begin{pmatrix} 1 & i \\ i & -1 \end{pmatrix} = \frac{i}{2}[w_1 \circ w_2 - iw_1 \circ w_2 i] \in \mathfrak{p}'^+ \end{aligned}$$

give rise to the classical Maass lowering and raising operators for SL_2 .

Pairing up We define $\mathbb{W} = V \otimes_{\mathbb{C}} W$, which we consider as a real vector space of dimension $2(r + s)m$. We define a symplectic form on \mathbb{W} by

$$\langle\langle v \otimes w, \tilde{v} \otimes \tilde{w} \rangle\rangle = \text{Re}(v, \tilde{v}) \langle w, \tilde{w} \rangle.$$

We note that $v_\alpha \otimes w_a, v_\mu \otimes w_a, v_\alpha \otimes w_u, v_\mu \otimes w_u$ and $v_\alpha \otimes -iw_a, v_\mu \otimes iw_a, v_\alpha \otimes iw_u, v_\mu \otimes -iw_u$ span Langrangian subspaces and give rise to a symplectic basis of \mathbb{W} . (Note that $\langle\langle v_\alpha \otimes w_a, v_\alpha \otimes -iw_a \rangle\rangle = 1$). Now $J = \theta \otimes J_0$ defines a positive definite complex structure on \mathbb{W} . We let $\mathbb{W}_{\mathbb{C}} = \mathbb{W} \otimes_{\mathbb{R}} \mathbb{C}$ be the complexification of \mathbb{W} , which again we view as a

right \mathbb{C} -vector space, and we extend $\langle\langle \cdot, \cdot \rangle\rangle$ \mathbb{C} -linearly. Then for the $+i$ -eigenspace \mathbb{W}' and the $-i$ eigenspace \mathbb{W}'' of J , we have

$$\begin{aligned}\mathbb{W}' &= \text{span}_{\mathbb{C}}\{v_{\alpha} \otimes w'_a, v_{\mu} \otimes w''_a, v_{\alpha} \otimes w'_u, v_{\mu} \otimes w''_u\}, \\ \mathbb{W}'' &= \text{span}_{\mathbb{C}}\{v_{\alpha} \otimes w''_a, v_{\mu} \otimes w'_a, v_{\alpha} \otimes w''_u, v_{\mu} \otimes w'_u\}.\end{aligned}$$

Note $\langle\langle v_{\alpha} \otimes w'_a, v_{\alpha} \otimes w''_a \rangle\rangle = 2i$. In the Fock model, $\mathfrak{sp}(\mathbb{W})$ acts on $\text{Sym}^{\bullet}(\mathbb{W}'')$, which we identify $\text{Sym}^{\bullet}(\mathbb{W}'')$ with $\mathcal{P}(\mathbb{C}^{(r+s)m})$ as follows. We denote the variables in $\mathcal{P}(\mathbb{C}^{(r+s)m})$ by $z''_{\alpha a}$ corresponding to $v_{\alpha} \otimes w''_a$, $z'_{\mu a}$ corresponding to $v_{\mu} \otimes w'_a$, $z'_{\alpha u}$ corresponding to $v_{\alpha} \otimes w'_u$, and $z''_{\mu u}$ corresponding to $v_{\mu} \otimes w'_u$. Thus we have

$$\begin{aligned}\rho_{\lambda}(v_{\alpha} \otimes w'_a) &= 2i\lambda \frac{\partial}{\partial z''_{\alpha a}}, & \rho_{\lambda}(v_{\alpha} \otimes w'_u) &= 2i\lambda \frac{\partial}{\partial z'_{\alpha u}}, \\ \rho_{\lambda}(v_{\alpha} \otimes w''_a) &= z''_{\alpha a}, & \rho_{\lambda}(v_{\alpha} \otimes w''_u) &= z'_{\alpha u}, \\ \rho_{\lambda}(v_{\mu} \otimes w''_a) &= 2i\lambda \frac{\partial}{\partial z'_{\mu a}}, & \rho_{\lambda}(v_{\mu} \otimes w''_u) &= 2i\lambda \frac{\partial}{\partial z''_{\mu u}}, \\ \rho_{\lambda}(v_{\mu} \otimes w'_a) &= z'_{\mu a}, & \rho_{\lambda}(v_{\mu} \otimes w'_u) &= z''_{\mu u}.\end{aligned}$$

Weil representation We naturally have $\mathfrak{u}(V) \times \mathfrak{u}(W) \subset \mathfrak{sp}(V \otimes W)$, and one easily checks that the inclusions $j_1 : \mathfrak{u}(V) \rightarrow \mathfrak{sp}(V \otimes W) \simeq \text{Sym}_{\mathbb{R}}^2(V \otimes W)$ and $j_2 : \mathfrak{u}(W) \rightarrow \mathfrak{sp}(V \otimes W) \simeq \text{Sym}_{\mathbb{R}}^2(V \otimes W)$ are given by

$$\begin{aligned}j_1(v \wedge \tilde{v}) &= \sum_{a=1}^r \left[(v \otimes iw_a) \circ (\tilde{v} \otimes w_a) - (v \otimes w_a) \circ (\tilde{v} \otimes iw_a) \right] \\ &\quad - \sum_{u=r+1}^{r+s} \left[(v \otimes iw_u) \circ (\tilde{v} \otimes w_u) - (v \otimes w_u) \circ (\tilde{v} \otimes iw_u) \right]\end{aligned}$$

and

$$\begin{aligned}j_2(w \circ \tilde{w}) &= \sum_{\alpha=1}^p \left[(v_{\alpha} \otimes w) \circ (v_{\alpha} \otimes \tilde{w}) + (iv_{\alpha} \otimes w) \circ (iv_{\alpha} \otimes \tilde{w}) \right] \\ &\quad - \sum_{\mu=p+1}^{p+q} \left[(v_{\mu} \otimes w) \circ (v_{\mu} \otimes \tilde{w}) + (iv_{\mu} \otimes w) \circ (iv_{\mu} \otimes \tilde{w}) \right].\end{aligned}$$

with $v, \tilde{v} \in V$ and $w, \tilde{w} \in W$. From this, we see

$$\begin{aligned}j_1((v \wedge \tilde{v}) + (iv \wedge \tilde{v})i) &= \frac{1}{i} \sum_{a=1}^r [v \otimes w'_a] \circ [\tilde{v} \otimes w''_a] - \frac{1}{i} \sum_{u=r+1}^{r+s} [v \otimes w''_u] \circ [\tilde{v} \otimes w'_u], \\ j_1((v \wedge \tilde{v}) - (iv \wedge \tilde{v})i) &= -\frac{1}{i} \sum_{a=1}^r [v \otimes w''_a] \circ [\tilde{v} \otimes w'_a] + \frac{1}{i} \sum_{u=r+1}^{r+s} [v \otimes w'_u] \circ [\tilde{v} \otimes w''_u]\end{aligned}$$

and

$$j_2(w \circ \tilde{w} \pm (iw \circ \tilde{w})i) = \sum_{\alpha=1}^p [v_\alpha \otimes (w \pm iwi)] \circ [v_\alpha \otimes (\tilde{w} \mp i\tilde{w}i)] \\ - \sum_{\mu=p+1}^{p+q} [v_\mu \otimes (w \pm iwi)] \circ [v_\mu \otimes (\tilde{w} \mp i\tilde{w}i)].$$

With this we obtain the formulas for the Weil representation (see p. 38).

Lemma B.1. *For the action of $\mathfrak{g} \simeq \mathfrak{u}(p, q)(\mathbb{C})$ on $\mathcal{P}(\mathbb{C}^{2mn})$, we have the following:*

(i) *The elements $Z'_{\alpha\beta}$, $Z''_{\alpha\beta}$ and $Z'_{\mu\nu}$, $Z''_{\mu\nu}$ in \mathfrak{k} act by*

$$\omega_\lambda(Z'_{\alpha\beta}) = -\omega_\lambda(Z''_{\beta\alpha}) = -\sum_{a=1}^r z''_{\alpha a} \frac{\partial}{\partial z''_{\beta a}} + \sum_{u=r+1}^{r+s} z'_{\beta u} \frac{\partial}{\partial z'_{\alpha u}} - \frac{r-s}{2} \delta_{\alpha\beta},$$

$$\omega_\lambda(Z'_{\mu\nu}) = -\omega_\lambda(Z''_{\nu\mu}) = -\sum_{a=1}^r z'_{\nu a} \frac{\partial}{\partial z'_{\mu a}} + \sum_{u=r+1}^{r+s} z''_{\mu u} \frac{\partial}{\partial z''_{\nu u}} - \frac{r-s}{2} \delta_{\mu\nu}.$$

(ii) *The elements $Z'_{\alpha\mu}$ of \mathfrak{p}^+ and $Z''_{\alpha\mu}$ of \mathfrak{p}^- act by*

$$\omega_\lambda(Z'_{\alpha\mu}) = -\frac{1}{2i\lambda} \sum_{a=1}^r z''_{\alpha a} z'_{\mu a} + 2i\lambda \sum_{u=r+1}^{r+s} \frac{\partial^2}{\partial z'_{\alpha u} \partial z''_{\mu u}}, \\ \omega_\lambda(Z''_{\alpha\mu}) = 2i\lambda \sum_{a=1}^r \frac{\partial^2}{\partial z''_{\alpha a} \partial z'_{\mu a}} - \frac{1}{2i\lambda} \sum_{u=r+1}^{r+s} z'_{\alpha u} z''_{\mu u}.$$

Lemma B.2. *For the action of $\mathfrak{g}' \simeq \mathfrak{u}(r, s)(\mathbb{C})$ on $\mathcal{P}(\mathbb{C}^{2mn})$, we have the following:*

(i) *For \mathfrak{k}' we have*

$$\omega_\lambda(w_a \circ w_b + iw_a \circ w_b i) = 2i \left[\sum_{\alpha=1}^p z''_{\alpha b} \frac{\partial}{\partial z''_{\alpha a}} - \sum_{\mu=p+1}^{p+q} z'_{\mu a} \frac{\partial}{\partial z'_{\mu b}} \right] + i(p-q)\delta_{ab}, \\ \omega_\lambda(w_u \circ w_v + iw_u \circ w_v i) = 2i \left[\sum_{\alpha=1}^p z'_{\alpha v} \frac{\partial}{\partial z'_{\alpha u}} - \sum_{\mu=p+1}^{p+q} z''_{\mu v} \frac{\partial}{\partial z''_{\mu u}} \right] + i(p-q)\delta_{uv}.$$

(ii) *For \mathfrak{p}'^\pm we have*

$$\omega_\lambda(w_a \circ w_u - iw_a \circ w_u i) = \frac{1}{\lambda} \sum_{\alpha=1}^p z''_{\alpha a} z'_{\alpha u} + 4\lambda \sum_{\mu=p+1}^{p+q} \frac{\partial^2}{\partial z'_{\mu a} \partial z''_{\mu u}}, \\ \omega_\lambda(w_a \circ w_u + iw_a \circ w_u i) = -4\lambda \sum_{\alpha=1}^p \frac{\partial^2}{\partial z''_{\alpha a} \partial z'_{\alpha u}} - \frac{1}{\lambda} \sum_{\mu=p+1}^{p+q} z'_{\mu a} z''_{\mu u}.$$

Intertwining We now give the intertwiner of the Fock model for $\lambda = 2\pi i$ with the Schrödinger model in the case when $r = s = n$. In that case, the Schrödinger model is given by the space of Schwartz functions $\mathcal{S}(V^n)$ on V^n .

The K' -finite vectors form the polynomial Fock space $S(V^n) \subset \mathcal{S}(V^n)$ which consists of functions on V^n of the form $p(\mathbf{z})\varphi_0(\mathbf{z})$, where $p(\mathbf{z})$ is a polynomial function on V^n and $\varphi_0(\mathbf{z})$ is the standard Gaussian on V^n . Here we use complex coordinates $\mathbf{z} = (z_1, \dots, z_n)$ with $z_i = {}^t(z_{1i}, \dots, z_{mi})$ in V relative to the basis $\{v_\alpha, v_\mu\}$. The Weil representation action of $\mathfrak{sp}(V \otimes W)$ now arises by the following action of the quantum algebra \mathbb{W}_λ :

$$\begin{aligned} \omega(v_\alpha \otimes w_j'') &= \sqrt{2}\pi i \left(\bar{z}_{\alpha j} - \frac{1}{\pi} \frac{\partial}{\partial z_{\alpha j}} \right), & \omega(v_\alpha \otimes w_{n+j}') &= \sqrt{2}\pi i \left(\bar{z}_{\alpha j} + \frac{1}{\pi} \frac{\partial}{\partial z_{\alpha j}} \right), \\ \omega(v_\alpha \otimes w_{n+j}'') &= \sqrt{2}\pi i \left(z_{\alpha j} - \frac{1}{\pi} \frac{\partial}{\partial \bar{z}_{\alpha j}} \right), & \omega(v_\alpha \otimes w_j') &= \sqrt{2}\pi i \left(z_{\alpha j} + \frac{1}{\pi} \frac{\partial}{\partial \bar{z}_{\alpha j}} \right), \\ \omega(v_\mu \otimes w_{n+j}') &= -\sqrt{2}\pi i \left(\bar{z}_{\mu j} - \frac{1}{\pi} \frac{\partial}{\partial z_{\mu j}} \right), & \omega(v_\mu \otimes w_j'') &= -\sqrt{2}\pi i \left(\bar{z}_{\mu j} + \frac{1}{\pi} \frac{\partial}{\partial z_{\mu j}} \right), \\ \omega(v_\mu \otimes w_j') &= -\sqrt{2}\pi i \left(z_{\mu j} - \frac{1}{\pi} \frac{\partial}{\partial \bar{z}_{\mu j}} \right), & \omega(v_\mu \otimes w_{n+j}'') &= -\sqrt{2}\pi i \left(z_{\mu j} + \frac{1}{\pi} \frac{\partial}{\partial \bar{z}_{\mu j}} \right). \end{aligned}$$

Here $1 \leq j \leq n$. We then have a unique \mathbb{W}_λ -intertwining operator $\iota : S(V^n) \rightarrow \mathcal{P}(\mathbb{C}^{2nm})$ satisfying $\iota(\varphi_0) = 1$, and we have

Lemma B.3. *The intertwining operator between the Schrödinger and the Fock model satisfies*

$$\begin{aligned} \iota \left(\bar{z}_{\alpha j} - \frac{1}{\pi} \frac{\partial}{\partial z_{\alpha j}} \right) \iota^{-1} &= -i \frac{1}{\sqrt{2}\pi} z_{\alpha j}'', & \iota \left(\bar{z}_{\alpha j} + \frac{1}{\pi} \frac{\partial}{\partial z_{\alpha j}} \right) \iota^{-1} &= 2\sqrt{2}i \frac{\partial}{\partial z_{\alpha n+j}'}, \\ \iota \left(z_{\alpha j} - \frac{1}{\pi} \frac{\partial}{\partial \bar{z}_{\alpha j}} \right) \iota^{-1} &= -i \frac{1}{\sqrt{2}\pi} z_{\alpha n+j}', & \iota \left(z_{\alpha j} + \frac{1}{\pi} \frac{\partial}{\partial \bar{z}_{\alpha j}} \right) \iota^{-1} &= 2\sqrt{2}i \frac{\partial}{\partial z_{\alpha j}''}, \\ \iota \left(\bar{z}_{\mu j} - \frac{1}{\pi} \frac{\partial}{\partial z_{\mu j}} \right) \iota^{-1} &= i \frac{1}{\sqrt{2}\pi} z_{\mu n+j}'', & \iota \left(\bar{z}_{\mu j} + \frac{1}{\pi} \frac{\partial}{\partial z_{\mu j}} \right) \iota^{-1} &= -2\sqrt{2}i \frac{\partial}{\partial z_{\mu j}'}, \\ \iota \left(z_{\mu j} - \frac{1}{\pi} \frac{\partial}{\partial \bar{z}_{\mu j}} \right) \iota^{-1} &= i \frac{1}{\sqrt{2}\pi} z_{\mu j}', & \iota \left(z_{\mu j} + \frac{1}{\pi} \frac{\partial}{\partial \bar{z}_{\mu j}} \right) \iota^{-1} &= -2\sqrt{2}i \frac{\partial}{\partial z_{\mu n+j}''}. \end{aligned}$$

References

- [1] Jeffrey Adams. The theta correspondence over \mathbb{R} . In *Harmonic analysis, group representations, automorphic forms and invariant theory*, volume 12 of *Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap.*, pages 1–39. World Sci. Publ., Hackensack, NJ, 2007.
- [2] Richard E. Borcherds. Automorphic forms with singularities on Grassmannians. *Invent. Math.*, 132(3):491–562, 1998.

- [3] Richard E. Borcherds. The Gross-Kohnen-Zagier theorem in higher dimensions. *Duke Math. J.*, 97(2):219–233, 1999.
- [4] Jan H. Bruinier. *Borcherds products on $O(2, 1)$ and Chern classes of Heegner divisors*, volume 1780 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2002.
- [5] Jan Hendrik Bruinier. Regularized theta lifts for orthogonal groups over totally real fields. *J. Reine Angew. Math.*, 672:177–222, 2012.
- [6] Jan Hendrik Bruinier and Jens Funke. On two geometric theta lifts. *Duke Math. J.*, 125(1):45–90, 2004.
- [7] Jan-Hendrik Bruinier, Ben Howard, Stephen S. Kudla, Michael Rapoport, and Tonghai Yang. Modularity of generating series of divisors on unitary shimura varieties. preprint, 2017.
- [8] Jan-Hendrik Bruinier, Ben Howard, Stephen S. Kudla, Michael Rapoport, and Tonghai Yang. Modularity of generating series of divisors on unitary shimura varieties ii. preprint, 2017.
- [9] Jan Hendrik Bruinier, Benjamin Howard, and Tonghai Yang. Heights of Kudla-Rapoport divisors and derivatives of L -functions. *Invent. Math.*, 201(1):1–95, 2015.
- [10] Jan Hendrik Bruinier and Ulf Kühn. Integrals of automorphic Green’s functions associated to Heegner divisors. *Int. Math. Res. Not.*, (31):1687–1729, 2003.
- [11] Jan Hendrik Bruinier and Tonghai Yang. Arithmetic degrees of special cycles and derivatives of siegel eisenstein series. preprint, 2018.
- [12] Stephan Ehlen and Siddharth Sankaran. On two arithmetic theta lifts. *Compos. Math.*, 154(10):2090–2149, 2018.
- [13] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi. *Tables of integral transforms. Vol. I*. McGraw-Hill Book Company, Inc., New York-Toronto-London, 1954. Based, in part, on notes left by Harry Bateman.
- [14] Mogens Flensted-Jensen. Discrete series for semisimple symmetric spaces. *Ann. of Math. (2)*, 111(2):253–311, 1980.
- [15] Jens Funke and John Millson. Cycles with local coefficients for orthogonal groups and vector-valued Siegel modular forms. *Amer. J. Math.*, 128(4):899–948, 2006.
- [16] Jens P. Funke and Stephen S. Kudla. Mock Modular Forms and Geometric Theta Functions for Indefinite Quadratic Forms. preprint, 2017.
- [17] Luis Garcia and Siddharth Sankaran. Green forms and the arithmetic siegel-weil formula. *Inv. Math.* to appear.

- [18] Jeffrey A. Harvey and Gregory Moore. Algebras, BPS states, and strings. *Nuclear Phys. B*, 463(2-3):315–368, 1996.
- [19] Eric Hofmann. Borcherds products on unitary groups. *Mathematische Annalen*, 354:799–832, 2014.
- [20] Tobias Hüfler. *Automorphe Formen auf orthogonalen und unitären Gruppen*. PhD thesis, TU Darmstadt, 2017.
- [21] Markus Schwagenscheidt Kathrin Bringmann, Stephan Ehlen. On the modular completion of certain generating functions. *ArXiv e-prints*, 2018.
- [22] Stephen S. Kudla. On certain arithmetic automorphic forms for $SU(1, q)$. *Invent. Math.*, 52(1):1–25, 1979.
- [23] Stephen S. Kudla. Central derivatives of Eisenstein series and height pairings. *Ann. of Math. (2)*, 146(3):545–646, 1997.
- [24] Stephen S. Kudla. Integrals of Borcherds forms. *Compositio Math.*, 137(3):293–349, 2003.
- [25] Stephen S. Kudla. Special cycles and derivatives of Eisenstein series. In *Heegner points and Rankin L-series*, volume 49 of *Math. Sci. Res. Inst. Publ.*, pages 243–270. Cambridge Univ. Press, Cambridge, 2004.
- [26] Stephen S. Kudla and John J. Millson. The theta correspondence and harmonic forms. I. *Math. Ann.*, 274(3):353–378, 1986.
- [27] Stephen S. Kudla and John J. Millson. The theta correspondence and harmonic forms. II. *Math. Ann.*, 277(2):267–314, 1987.
- [28] Stephen S. Kudla and John J. Millson. Intersection numbers of cycles on locally symmetric spaces and Fourier coefficients of holomorphic modular forms in several complex variables. *Inst. Hautes Études Sci. Publ. Math.*, (71):121–172, 1990.
- [29] Yifeng Liu. Arithmetic theta lifting and L -derivatives for unitary groups. I. *Algebra Number Theory*, 5(7):849–921, 2011.
- [30] Takayuki Oda and Masao Tsuzuki. The secondary spherical functions and automorphic Green currents for certain symmetric pairs. *Pure Appl. Math. Q.*, 5(3, Special Issue: In honor of Friedrich Hirzebruch. Part 2):977–1028, 2009.
- [31] Takuro Shintani. On construction of holomorphic cusp forms of half integral weight. *Nagoya Math. J.*, 58:83–126, 1975.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF DURHAM, SOUTH ROAD, DURHAM DH13LE, UK

E-mail address: jens.funke@durham.ac.uk

MATHEMATISCHES INSTITUT, UNIVERSITÄT HEIDELBERG, IM NEUENHEIMER FELD 205,
D-69120 HEIDELBERG, GERMANY
E-mail address: hofmann@mathi.uni-heidelberg.de