

On different notions of tameness in arithmetic geometry

Moritz Kerz and Alexander Schmidt

March 17, 2009

Abstract: The notion of a tamely ramified covering is canonical only for curves. Several notions of tameness for coverings of higher dimensional schemes have been used in the literature. We show that all these definitions are essentially equivalent. Furthermore, we prove finiteness theorems for the tame fundamental groups of arithmetic schemes.

1 Introduction

Let \bar{C} be a proper, connected and regular curve (i.e. $\dim C = 1$) of finite type over $\text{Spec}(\mathbb{Z})$ and let $C \subset \bar{C}$ be a nonempty open subscheme. Every point $x \in \bar{C} \setminus C$ defines a discrete rank one valuation v_x on the function field $k(C)$. One says that an étale covering $C' \rightarrow C$ is *tamely ramified along $\bar{C} \setminus C$* if for every $x \in \bar{C} \setminus C$ the valuation v_x is tamely ramified in $k(C')|k(C)$. Since the proper, regular curve \bar{C} is determined by C , we can say that the étale covering $C' \rightarrow C$ is *tame* if it is tamely ramified along $\bar{C} \setminus C$.

Following [SGA1, GM], one possible extension of this definition to higher dimensions is the following. We denote by $Sch(\mathbb{Z})$ the category of separated schemes of finite type over $\text{Spec}(\mathbb{Z})$. Let $X \in Sch(\mathbb{Z})$ be a regular scheme together with an open embedding into a regular, proper scheme $\bar{X} \in Sch(\mathbb{Z})$ such that $\bar{X} \setminus X$ is a normal crossing divisor (NCD) on \bar{X} . Then an étale covering $Y \rightarrow X$ is called *tamely ramified along $\bar{X} \setminus X$* if the discrete valuations associated with the generic points of $\bar{X} \setminus X$ are tamely ramified in $k(Y)|k(X)$. However, there might exist many or (at our present knowledge about resolution of singularities) even no regular compactifications \bar{X} of X such that $\bar{X} \setminus X$ is a NCD. Furthermore, there is no obvious functoriality for the tame fundamental group.

So the question for a good notion of tameness of an étale covering $Y \rightarrow X$ of regular schemes in $Sch(\mathbb{Z})$ naturally occurs. In this paper we compare several possible definitions:

curve-tameness: for every morphism $C \rightarrow X$ with $C \in Sch(\mathbb{Z})$ a regular curve, the base change $Y \times_X C \rightarrow C$ is tame.

divisor-tameness: for every normal compactification \bar{X} of X and every point $x \in \bar{X} \setminus X$ with $\text{codim}_{\bar{X}} x = 1$, the discrete rank one valuation v_x on $k(X)$ associated with x is tamely ramified in $k(Y)|k(X)$.

chain-tameness: There exists a compactification \bar{X} of X such that every discrete valuation of rank $d = \dim X$ on $k(X)$ which dominates a Parshin chain on \bar{X} is tamely ramified in $k(Y)|k(X)$.

valuation-tameness: every nonarchimedean valuation of $k(X)$ with center outside X is tamely ramified in $k(Y)|k(X)$.

The notion of curve-tameness has been considered in [Wi] and the notion of chain-tameness in [Sc1]. Curve-tameness is the maximal definition of tameness which is stable under base change and extends the given definition for curves. Valuation tameness is obviously stronger than divisor-tameness and chain-tameness. Our first result is the following:

Theorem 1.1 (see Theorem 4.4). *The notions of curve-tameness, divisor-tameness and chain-tameness are equivalent. If every intermediate field between $k(X)$ and the Galois closure of $k(Y)$ over $k(X)$ admits a regular proper model, then they are equivalent to valuation-tameness.*

Moreover, if there exists a regular compactification \bar{X} such that $\bar{X} \setminus X$ is a NCD, then all these notions of tameness are equivalent to the notion of tame ramification along $\bar{X} \setminus X$ of [SGA1, GM].

Suppose that $\pi : Y \rightarrow X$ is Galois with group G and that we are given a fixed normal compactification \bar{X} of X . Denoting the normalization of \bar{X} in $k(Y)$ by \bar{Y} , there are the following additional notions of tameness:

numerical tameness: for every $y \in \bar{Y} \setminus Y$ the inertia group $T_y(\bar{Y}|\bar{X}) \subset G$ is of order prime to the characteristic of the residue field $k(y)$.

cohomological tameness: for every $x \in \bar{X} \setminus X$ the semi-local ring $\mathcal{O}_{\bar{Y}, \pi^{-1}(x)}$ is a cohomologically trivial G -module.

The notion of numerical tameness has been considered in [Ab], [CE], [Sc1] and [Wi]. Cohomological tameness has been considered in [CEPT] (in the more general context of group scheme actions).

Our second result is :

Theorem 1.2 (see Theorems 5.3 and 6.2). *Numerical tameness and cohomological tameness are equivalent and imply valuation tameness. If $\bar{X} \setminus X$ is a NCD or G is nilpotent, then all definitions are equivalent.*

We show Theorems 1.1 and 1.2 in the more general context of regular schemes, separated and of finite type over an integral, pure-dimensional and excellent base scheme S . In the appendix we give illustrating examples, in particular, an example of an étale covering of a regular scheme which is numerically tame with respect to one regular compactification (hence curve-, divisor- and chain-tame) but not numerically tame with respect to another regular compactification.

In the case that the function field of S is absolutely finitely generated, we show the following tame variant of a finiteness theorem of Katz and Lang [KL].

Theorem 1.3 (see Theorem 7.1). *Let S be an integral, pure-dimensional and excellent base scheme whose function field is absolutely finitely generated. Let $f : Y \rightarrow X$ be a smooth, surjective morphism of connected regular schemes which are separated and of finite type over S . Assume that the generic fibre of f is geometrically connected and that one of the following conditions (i) and (ii) is satisfied:*

- (i) *X has a regular compactification \bar{X} over S such that the boundary $\bar{X} \setminus X$ is a normal crossing divisor.*
- (ii) *The generic fibre of f has a rational point.*

Then the group

$$\ker_S^t(Y/X) := \ker(\pi_1^{t,ab}(Y/S) \rightarrow \pi_1^{t,ab}(X/S))$$

is finite.

The authors thank A. Holschbach and F. Pop for helpful discussions on the subject, and T. Chinburg for his proposal to consider cohomological tameness.

2 The Key Lemma

We start by recalling the following desingularization results. See [Lip] for a proof of (i) and [Sha], Lecture 3, Theorem on p. 38 and Remark 2 on p. 43, for (ii). We denote the set of regular points of a scheme X by X^{reg} .

Proposition 2.1. *Let X be a two-dimensional, normal, connected and excellent scheme. Then the following hold.*

- (i) *After a finite number of blow-ups in closed points followed by normalizations, we obtain a proper morphism $p : X' \rightarrow X$ with X' regular and such that $p^{-1}(X^{reg}) \rightarrow X^{reg}$ is an isomorphism of schemes.*
- (ii) *Assume that X is regular and let Z be a proper closed subset on X . After a finite number of blow-ups in closed points, we obtain a proper morphism $p : X' \rightarrow X$ such that $p^{-1}(Z)$ is a strict normal crossing divisor on X' .*

Let X be a normal, noetherian scheme and let $X' \subset X$ be a dense open subscheme. Assume we are given an étale covering $Y' \rightarrow X'$.

Definition. Let $x \in X \setminus X'$ be a point. We say that $Y' \rightarrow X'$ is *unramified along x* if it extends to an étale covering of some open subscheme $U \subset X$ which contains X' and x . Otherwise we say that $Y' \rightarrow X'$ *ramifies along x* .

Remark 2.2. Let Y be the normalization of X in $k(Y')$. Then the branch locus of $Y \rightarrow X$ consists of all points $x \in X \setminus X'$ such that $Y' \rightarrow X'$ ramifies along x .

Remark 2.3. If $\text{codim}_X\{x\} = 1$, then $Y' \rightarrow X'$ ramifies along x if and only if the discrete valuation of $k(X')$ associated with x ramifies in $k(Y')$. In this case we can speak about *tame* and *wild* ramification along x by referring to the associated valuation.

The Key Lemma for our investigations is the following Lemma 2.4. An integral, one-dimensional scheme will be called a curve. We denote the normalization of a curve C in its function field $k(C)$ by \tilde{C} .

Lemma 2.4 (Key Lemma). *Let A be a local, normal and excellent ring and let $X' \subset X = \text{Spec}(A)$ be a nonempty open subscheme. Let $Y' \rightarrow X'$ be an étale Galois covering of prime degree p . Assume that $X \setminus X'$ contains an irreducible component D of codimension one in X such that $Y' \rightarrow X'$ is ramified along the generic point of D . Then there exists a curve C on X with $C' := C \cap X' \neq \emptyset$ such that the base change $Y' \times_{X'} \tilde{C}' \rightarrow \tilde{C}'$ is ramified along a point of $\tilde{C} \setminus \tilde{C}'$.*

In the proof of the Key Lemma we use a reduction argument to dimension two which is due to Wiesend [Wi, Proof of Thm. 2]. Wiesend's arguments for the two-dimensional case are however incomplete. O. Gabber has obtained a similar result using a slightly different technique.

In the proof of our Key Lemma we need the following ramification criterion.

Lemma 2.5. *Let R be a discrete valuation ring with prime element π , K the quotient field of R and p a prime number.*

- (i) *If $K'|K$ is a separable extension of discretely valued fields of degree p with trivial residue field extension and ramification index p , then the minimal polynomial f of any prime element of R' is of the form $f = T^p + a_{p-1}T^{p-1} + \cdots + a_0 \in R[T]$ with $\pi|a_i$ for every i and $\pi^2 \nmid a_0$.*
- (ii) *Let $f \in K[T]$ be a separable polynomial of the form $f = T^p + a_{p-1}T^{p-1} + \cdots + a_0$ with $a_0 \neq 0$, $p \nmid v_R(a_0)$, and such that*

$$\min_{i \geq 0} v_R(a_i) > p \max_{i \geq 0} [v_R(a_0) - v_R(a_i)].$$

Then $K' = K[T]/(f)$ is a discretely valued field with ramification index p over K .

Proof. Assertion (i) is standard, see for example [Se, I.6 Proposition 18]. For assertion (ii), we substitute T by $\pi^l T$, where $l = \max_{i \geq 0} [v_R(a_0) - v_R(a_i)]$, and divide f by π^{p^l} . Let

$$g = T^p + b_{p-1}T^{p-1} + \cdots + b_0$$

denote the resulting monic polynomial. We claim that $0 < v_R(b_0) \leq v_R(b_i)$ for all $i \geq 0$ and still $p \nmid v_R(b_0)$. In fact, we have $b_i = \pi^{l(i-p)} a_i$ so that $v_R(b_i) = v_R(a_i) - l(p-i) > 0$. We have $l \geq v_R(a_0) - v_R(a_i)$ by definition, which implies

$$v_R(b_i) - v_R(b_0) = v_R(a_i) - v_R(a_0) + lp - l(p-i) = v_R(a_i) - v_R(a_0) + li \geq 0,$$

validating our claim.

Let R' be the normalization of R in $K[T]/(g) \cong K[T]/(f) = K'$. Let R'' be the localization of R' at some maximal ideal. We will show that the ramification index e of $R''|R$ is p . This shows that $R' = R''$ and completes the proof of (ii). Let $t \in R''$ be the image of the variable T under the homomorphism $R[T]/(g) \rightarrow R''$. The equation

$$t^p = -b_{p-1}t^{p-1} - \cdots - b_0$$

shows that $t \notin R''^\times$. Furthermore, it gives the equality in the middle of

$$p v_{R''}(t) = v_{R''}(t^p) = v_{R''}(b_0) = e v_R(b_0).$$

As $p \nmid v_R(b_0)$, this implies $p|e$ and, as $e \leq p$, this indeed means $e = p$. \square

Proof of the Key Lemma 2.4. Let Y be the normalization of X in $k(Y')$. Then $Y = \text{Spec}(B)$ for some semi-local ring B .

Reduction to $\dim(A) = 2$. The assertion is trivial if $\dim(A) = 1$. Assume that we have shown Lemma 2.4 for $\dim(A) = 2$. We use induction on $\dim(A)$ to prove the general case. For $\dim(A) > 2$, choose a point $x_1 \in D$ of dimension one. Set $X_1 = \text{Spec}(A_{x_1})$, where A_{x_1} means the localization of A at the prime ideal corresponding to x_1 , and $X'_1 = X_1 \cap X'$. We deduce by our induction assumption that there exists a curve C_1 on X_1 with $C'_1 = C_1 \cap X'_1 \neq \emptyset$ such that $Y' \times_{X'} \tilde{C}'_1 \rightarrow \tilde{C}'_1$ is ramified along $D \times_X \tilde{C}'_1$. Let X_2 be the normalization of the closure of C_1 in X , X'_2 be $X_2 \times_X X'$ and Y'_2 be the normalization of X'_2 in $Y' \otimes k(C)$. Finally, the two-dimensional case produces a curve C_2 on X_2 with $C'_2 = C_2 \cap X'_2 \neq \emptyset$ such that $Y'_2 \times_{X'_2} \tilde{C}'_2 \rightarrow \tilde{C}'_2$ is ramified along $D \times \tilde{C}'_2$. Let C be the image of C_2 under the morphism $X_2 \rightarrow X$, C is a curve on X with $C' = C \cap X' \neq \emptyset$ such that $Y' \times_{X'} \tilde{C}' \rightarrow \tilde{C}'$ is ramified along $D \cap C$. The latter since $Y \otimes_X k(C)$ is ramified over some point of $D \times \tilde{C}$ if $Y \otimes_X k(C_2) = Y_2 \otimes_{X_2} k(C_2)$ is ramified over some point of $D \times \tilde{C}'_2$ by the base change invariance of étale morphisms.

Reduction to $\dim(A) = 2$ and A and D regular. Applying Proposition 2.1 to $\text{Spec}(A)$ and D , and localizing at a closed point of the strict transform of D , we can assume without loss generality that A and D are regular.

Now Lemma 2.4 will be proved by distinguishing three possible cases with different ramification indices. Since A is factorial, there exists an irreducible $\pi \in A$ with $D = V(\pi)$. Now $R_D = A/(\pi)$, $R = A_{(\pi)}$ and $R' = B_{(\pi)}$ are discrete valuation rings. We distinguish several possible cases:

1st case: $v_{R'}(\pi) = p$.

Let π' be a prime element of R' . After multiplication by units of R' we can assume that $\pi' \in A'$. The minimal polynomial $f \in K[T]$ of π' lies in $A[T]$ and, according to Lemma 2.5 (i), is of the form $f = T^p + a_{p-1}T^{p-1} + \cdots + a_0$ with $\pi|a_i$ for every i but $\pi^2 \nmid a_0$. Observe that $A[T]/(f)$ and B are isomorphic over some dense open subscheme of X . We use Proposition 2.1 to resolve singularities of the divisor generated by the coefficients of f and localize at the closed point of the strict transform of D . This new ring will still be denoted by A , and the induced open subscheme of $X = \text{Spec}(A)$ by X' .

Let π be a prime element in the new desingularized ring with $D = V(\pi)$ and denote by $t \in A$ an element such that $V(t)$ is the exceptional divisor. Then (t, π) is the maximal ideal of A and the a_i are supported on $V(t) \cup D$, so that a_i is up to a unit of A equal to $t^{l_i} \pi^{e_i}$ with $e_0 = 1$ and $e_i \geq 1$ for $i > 0$. In order to prove the Key Lemma 2.4, we will construct a regular curve C on X of the form $C = V(t^N - \pi)$ with the required properties.

For this we choose a natural number N such that

- $N > p \max_{i \geq 0} (l_0 - l_i)$.
- $C = V(t^N - \pi)$ meets X' and over the generic point of C the algebras $A[T]/(f)$ and B are isomorphic.
- $N + l_0$ is not divisible by p .

Then $R_C = A/(t^N - \pi)$ is a discrete valuation ring and, since $(t, t^N - \pi) = (t, \pi)$ is the maximal ideal of A , the element t induces a prime element of R_C . Let $K_C = Q(R_C)$. Our assumption on N shows that $K'_C := B \otimes_A K_C \cong K_C[T]/(f_C)$, where $f_C \in R_C[T]$ is the polynomial induced by f . The polynomial f_C is separable as B is étale over X' and of the form

$$f_C = T^p + a_{p-1}^C T^{p-1} + \cdots + a_0^C$$

with $v_{R_C}(a_i) = N e_i + l_i$ for $i \geq 0$. Therefore the assumptions of Lemma 2.5 (ii) are fulfilled, so that K'_C is discretely valued and has ramification index p over K_C .

2nd case: $v_{R'}(\pi) = 1$.

In this case the residue field extension $K'_D|K_D$ of $R'|R$ is an inseparable extension of degree p . In particular, $K_D = Q(R_D)$ has characteristic p . Let R'_D be the normalization of R_D in K'_D and let π_D be a prime element of R_D . Then R'_D is a discrete valuation ring by [Bo, VI, 8.6 Corollary 2]. There are two subcases:

Subcase 2a: $v_{R'_D}(\pi_D) = p$.

In this subcase we have $K'_D = K_D(\sqrt[p]{\pi_D})$ if we choose the prime element π_D appropriately. Let $f \in R[T]$ be the minimal polynomial of a lift to R' of $\sqrt[p]{\pi}$. After a resolution of singularities as in Proposition 2.1 of the divisor generated by D and the coefficients of f and after localizing at the closed point of the strict transform of D , we can assume the following:

- The coefficients of f are supported on $E \cup D$, where E is the exceptional divisor,
- E and D intersect transversely.

The constant coefficient t of f still induces a prime element in R_D and by assumption t is supported on E so that we conclude $E = V(t)$. So f is of the form $f = T^p + a_{p-1}T^{p-1} + \cdots + a_1T + t$ with a_i for $i > 0$ up to a unit of A equal to $\pi^{e_i}t^{l_i}$ with $e_i > 0$ and $l_i \in \mathbb{Z}$. The ideal (t, π) of A is the maximal ideal, since E and D intersect transversely. There exists some dense open subscheme of $\text{Spec}(A_t)$ such that B and $A_t[T]/(f)$ are isomorphic over this subscheme. We will construct a natural number N such that the regular curve $C = V(t^N - \pi)$ has the required properties. In fact, choose $N > 0$ such that the following properties are satisfied:

- $N + l_i > 0$ for all $i > 0$.
- The generic point of $V(t^N - \pi)$ lies in X' and in the open subscheme of $\text{Spec}(A_t)$ where B and $A_t[T]/(f)$ are isomorphic.

Since $(t, t^N - \pi) = (t, \pi)$ is the maximal ideal of A , the element t induces a prime element of the discrete valuation ring $R_C = A/(t^N - \pi)$. With $K_C = Q(R_C)$, we have $B \otimes_A K_C \cong K_C[T]/(f_C)$, where f_C is the separable polynomial in $K_C[T]$ induced by f . Then f_C is of the form

$$f_C = T^p + a_{p-1}^C T^{p-1} + \cdots + a_0^C$$

with $v_{R_C}(a_i^C) > 0$ for every i and $v_{R_C}(a_0^C) = 1$. Finally, Lemma 2.5 (ii) shows that K'_C is discretely valued and ramified over K_C with ramification index p .

Subcase 2b: $v_{R'_D}(\pi_D) = 1$.

In this subcase the residue extension of $R'_D|R_D$ is an inseparable extension of degree p , so that $K'_D = K_D(\sqrt[p]{\xi})$ for some $\xi \in R_D^\times$ which is not a p -th power in the residue field of R_D . Lift ξ to R' and denote its minimal polynomial by $f \in R[T]$. As in the previous subcase, we resolve singularities and localize so that without loss of generality we can assume the coefficients of f are supported on $E \cup D$, where $E = V(t)$ is the exceptional divisor, and so that E and D intersect transversely. The prime element t can be chosen arbitrarily now, in contrast to subcase 2a. Observe that the residue field of A does not change in the desingularization process (we blow up a regular scheme in a regular center). Then f is of the form $f = T^p + a_{p-1}T^{p-1} + \cdots + a_1T + a_0$ with a_i for $i > 0$ up to a unit of A equal to $\pi^{e_i}t^{l_i}$ with $e_i > 0$ and $l_i \in \mathbb{Z}$ and such that $a_0 \in A^\times$ induces an element in the residue field of A which is not a p -th power. There exists an open subscheme of $\text{Spec}(A_t)$ such that B and $A_t[T]/(f)$ are isomorphic over this subscheme. Again the curve C we are searching for will be of the form $C = V(t^N - \pi)$ for some natural number N . We choose $N > 0$ such that the following properties are satisfied:

- $N + l_i > 0$ for all $i > 0$,
- For $C = V(t^N - \pi)$ the intersection $C \cap X'$ is nonempty and the generic point of C lies in the open subscheme of $\text{Spec}(A_t)$ over which B and $A_t[T]/(f)$ are isomorphic.

As in the previous cases it follows that t induces a prime element of the discrete valuation ring $R_C = A/(t^N - \pi)$ and that, with $K_C = Q(R_C)$, we have $K'_C := B \otimes_A K_C \cong K_C[T]/(f_C)$, where f_C is the polynomial in $K_C[T]$ induced by f . Then f_C is of the form

$$f_C = T^p + a_{p-1}^C T^{p-1} + \cdots + a_0^C$$

with $v_{R_C}(a_i^C) > 0$ for every $i > 0$ and such that $a_0 \in R_C^\times$ is not a p -th power in the residue field of R_C . It follows that K'_C is a discretely valued field ramified over K_C with ramification index $e = 1$. \square

3 Some valuation theory

Working over a general base scheme S , we first have to fix some notation.

Definition. We call an integral noetherian scheme X *pure-dimensional* if $\dim X = \dim \mathcal{O}_{X,x}$ for every closed point $x \in X$.

Remark 3.1. Any integral scheme of finite type over a field or over a Dedekind domain with infinitely many prime ideals is pure-dimensional. A proper scheme over a pure-dimensional universally catenary scheme is pure-dimensional, see [EGA4], IV, 5.6.5. The affine line $\mathbb{A}_{\mathbb{Z}_p}^1$ over the ring of p -adic integers gives an example of a regular scheme which is not pure-dimensional.

Let from now on S be an integral, pure-dimensional, separated and excellent base scheme. We work in the category $Sch(S)$ of separated schemes of finite type over S . In order to avoid the effect that open subschemes might have smaller (Krull-)dimension than the ambient scheme (e.g. $Spec(\mathbb{Q}_p) \subset Spec(\mathbb{Z}_p)$), we redefine the notion of dimension for schemes in $Sch(S)$ as follows:

Let $X \in Sch(S)$ be integral and let T be the closure of the image of X in S . Then we put

$$\dim_S X := \deg.\text{tr.}(k(X)|k(T)) + \dim_{\text{Krull}} T.$$

If the image of X in S contains a closed point of T , then $\dim_S X = \dim_{\text{Krull}} X$ by [EGA4], IV, 5.6.5. This equality holds for arbitrary $X \in Sch(S)$ if S is of finite type over a field or over a Dedekind domain with infinitely many prime ideals.

Definition. A *function field over S* is the function field $K = k(X)$ of some integral scheme $X \in Sch(S)$. We call X a *model of K* and put $\dim_S K = \dim_S X$.

Every function field over S admits a proper model, see [Lue] for a scheme theoretic proof. Let $K = k(X)$ be a function field over S and let T be the closure of the image of X in S (with reduced scheme structure). Then T is an integral closed subscheme in S .

Definition. We denote by $\text{Val}_S(K)$ the set of nonarchimedean valuations v on K such that \mathcal{O}_v dominates $\mathcal{O}_{T,t}$ for some point $t \in T$. For $v \in \text{Val}_S(K)$ and for a given proper S -model X of K we denote by Z_v the center of v on X , which is the (uniquely defined) integral closed subscheme of X such that \mathcal{O}_v dominates \mathcal{O}_{X,Z_v} .

Let $n \leq \dim_S K$ be a natural number. We consider \mathbb{Z}^n as an ordered group with the lexicographic ordering.

Definition. We call an S -valuation $v : K^\times \rightarrow \mathbb{Z}^n$, i.e. a valuation with ordered value group \mathbb{Z}^n , a *discrete rank n valuation on K* . The residue field is denoted by Kv .

The following proposition, due to Abhyankar, is well known. The inequality in the proposition is called the Abhyankar inequality.

Proposition 3.2. *Let K be a function field over S and let \bar{X} be a proper model of K . Let v be a discrete rank n valuation on K with center Z_v on \bar{X} . Then we have the inequality*

$$\deg.\text{tr.}(Kv|k(Z_v)) + \dim_S Z_v + n \leq \dim_S K.$$

Suppose that equality holds. Then $Kv|k(Z_v)$ is a finitely generated field extension. Moreover, there exists a proper model \bar{X}' of K such that $\dim_S Z'_v = \dim_S K - n$ and $Kv = k(Z'_v)$.

Proof. Let z_v be the generic point of Z_v . Since S is pure-dimensional, also \bar{X} is pure-dimensional and the Krull dimension of the ring $A := \mathcal{O}_{\bar{X}, z_v}$ is equal to $\dim_S K - \dim_S Z_v$. So [Va, Théorème 9.2] comprises our lemma except for the last statement. In order to prove the latter, let $s_1, \dots, s_m \in \mathcal{O}_v^\times$ be a finite family of elements which generate the residue field of \mathcal{O}_v over the residue field of the local ring A . Let $\text{Spec}(B) \subset \bar{X}$ be an open neighbourhood of z_v and set $X' := \text{Spec}(B[s_1, \dots, s_m])$. Then v has nontrivial center in X' and the function field of this center coincides with Kv . Finally, we let $X' \subset \bar{X}' \rightarrow \bar{X}$ be a compactification, which exists by [Lue]. \square

We call v *geometric* if equality holds in the Abhyankar inequality in Proposition 3.2. This notion does not depend on the proper model we have chosen. If v is geometric, then Kv is a function field over S of dimension $\dim_S K - n$. In case $n = \dim_S K$ the Abhyankar inequality is automatically an equality, so that every discrete rank $\dim_S K$ valuation is geometric and the center on every proper model is a closed point.

If v is a valuation on K and w a valuation on Kv , then the ring

$$\mathcal{O}_{v \circ w} := \{x \in \mathcal{O}_v \mid \bar{x} \in \mathcal{O}_w\}$$

is a valuation ring and the associated valuation on K is called the composite valuation $v \circ w$.

The following lemma is ‘folklore’ but we could not find a reference.

Lemma 3.3. *Let K be a function field over S .*

- (i) *If v is a geometric discrete S -valuation of rank n on K and w is a geometric discrete S -valuation of rank m on Kv , then $v \circ w$ is a geometric discrete rank $(n + m)$ valuation.*
- (ii) *Each discrete geometric S -valuation v on K of rank n can be written in the form $v = v_1 \circ \dots \circ v_n$, where the v_i , $i = 1, \dots, n$, are geometric discrete rank 1 valuations on Kv_{i-1} (set $Kv_0 := K$).*

Proof. (i) We have to show that $v \circ w$ is geometric. We use Proposition 3.2 to find a proper model \bar{X} of K such that $Kv = k(Z_v)$, where Z_v is the center of v on \bar{X} . Then Z_v is a proper model of Kv and we denote the center of w on Z_v by Z_w . Note that $Z_w = Z_{v \circ w}$. The residue fields of w and of $v \circ w$ are the same. We get the equality

$$\begin{aligned} n + (\text{deg.tr.}(Kw|k(Z_w)) + m) &= (\dim_S K - \dim_S Z_v) + (\dim_S Z_v - \dim_S Z_w) \\ &= \dim_S K - \dim_S Z_{v \circ w}, \end{aligned}$$

which shows that $v \circ w$ is geometric.

(ii) Every discrete rank n valuation v can be uniquely decomposed into a chain $v = v_1 \circ \dots \circ v_n$ of discrete rank 1 valuations. By Proposition 3.2, for any proper model \bar{X} of K , the transcendence degree $\text{deg.tr.}(Kv_1|k(Z_{v_1}))$ is finite. In a similar way as in the proof of Proposition 3.2, we find a proper model \bar{X} of K such that $\text{deg.tr.}(Kv_1|k(Z_{v_1})) = 0$, i.e. the extension is algebraic. Let w be the restriction of $v_2 \circ \dots \circ v_n$ to $k(Z_{v_1})$. Then Z_{v_1} is a proper model of $k(Z_{v_1})$ and the center Z_w of w on Z_{v_1} is equal to Z_v . Furthermore, the residue fields of v and of $v_2 \circ \dots \circ v_n$ are the same, hence Kv is an algebraic extension of the residue field Kw of w . Using the Abhyankar inequality for v_1 and for w , we obtain

$$\begin{aligned} \text{deg.tr.}(Kv|k(Z_v)) + n &= 1 + (\text{deg.tr.}(Kw|k(Z_w)) + n - 1) \\ &\leq (\dim_S K - \dim_S Z_{v_1}) + (\dim_S Z_{v_1} - \dim_S Z_w) \\ &= \dim_S K - \dim_S Z_v. \end{aligned}$$

Since v is geometric, equality holds in both Abhyankar inequalities; hence v_1 and w are geometric. By Proposition 3.2, the extension $Kv_1|k(Z_{v_1})$ is finite, and so Kv_1 is a function field over S and $v_2 \circ \dots \circ v_n$ is a geometric discrete S -valuation of rank $n - 1$ on this field. Now the result follows by induction. \square

Let v be a valuation on the field K , $L|K$ a finite Galois extension with Galois group G and w a valuation on L which extends v . Recall that the decomposition and the inertia group $G_w(L|K)$ and $T_w(L|K)$ of w in $L|K$ are defined by

$G_w(L|K) = \{g \in G \mid gw = w\}$ and $T_w(L|K) = \ker(G_w(L|K) \rightarrow \text{Aut}(Lv|Kv))$, respectively. Furthermore, we have the ramification group

$$V_w(L|K) = \{g \in G_w(L|K) \mid w(gx/x - 1) > 0 \text{ for all } x \in L^\times\}.$$

If $\text{char}(Kv) = 0$, then $V_w(L|K) = 1$, and if $p := \text{char}(Kv) > 0$, then $V_w(L|K)$ is the unique p -Sylow subgroup of $T_w(L|K)$. If w and w' are different extensions of v , then the decomposition, inertia and ramification groups of w and w' are conjugate and we sometimes write $G_v(L|K)$, $T_v(L|K)$ and $V_v(L|K)$ for these groups if we don't care for conjugation. There are several definitions of ramification for valuations used in the literature. We use the following:

Definition. Let v be a valuation on a field K and let $L|K$ be a finite Galois extension. We say that v is *unramified* (resp. *tamely ramified*) in $L|K$ if $T_v(L|K)$ is trivial (resp. if $V_v(L|K)$ is trivial). We say that v is unramified (resp. tamely ramified) in a finite separable extension $L|K$ if it is unramified (resp. tamely ramified) in the Galois closure of $L|K$.

Remark 3.4. Let v be a valuation with residue characteristic p on a field K and let $L|K$ be a finite separable extension. Let w_1, \dots, w_n be the different extensions of v to L . We denote by $e(w_i|v)$ the ramification index of w_i over v and we put $f(w_i|v) = [Lw_i : Kv]$. By the results of [End], §22, the valuation v is unramified (resp. tamely ramified) in $L|K$ in the sense of the above definition if and only if the following conditions (1)–(3) hold:

- (1) $e(w_i|v) = 1$ (resp. $e(w_i|v)$ is prime to p) for $i = 1, \dots, n$,
- (2) $Lw_i|Kv$ is separable for $i = 1, \dots, n$,
- (3) $\sum_{i=1}^n e(w_i|v)f(w_i|v) = [L : K]$.

If v is a discrete rank n valuation, then conditions (1) and (2) already imply (3).

Lemma 3.5. *Let $L|K$ be a finite separable extension of function fields over S and let $d = \dim_S K$. Then the following are equivalent.*

- (i) $L|K$ is ramified at a (geometric) discrete rank d valuation.
- (ii) $L|K$ is ramified at a geometric discrete rank 1 valuation.

Proof. (ii) \Rightarrow (i) Let v be a ramified geometric discrete rank 1 valuation. Choose a geometric discrete rank $(d - 1)$ valuation v_1 on Kv and put $w = v \circ v_1$; to find such a valuation v_1 one can use Parshin chains, see Section 4. Then w is a ramified discrete rank d valuation.

(i) \Rightarrow (ii) We use induction on $d = \dim_S K$. For $d = 1$ the lemma is trivial. In case $d = 2$ and v is a ramified discrete rank 2 valuation of K choose a proper normal model X of K . Then consider the two-dimensional local ring $A = \mathcal{O}_{X,x}$ where x is the generic point of the center of v on X . After a modification we may assume that A is regular, see Proposition 2.1. Since the normal closure of A in L is ramified over A , purity of the branch locus [SGA2, X.3.4] gives us a ramified divisor on A which gives us a ramified geometric discrete rank 1 valuation. For $d > 2$ write $v = v_1 \circ w$ where v_1 is a discrete rank 1 valuation, which is geometric

according to Lemma 3.3 (ii). If v_1 ramifies in L we are done. Otherwise consider the finite product L_v of the residue fields of all extensions of v_1 to L . Then, as v ramifies in L and as v_1 does not ramify in L , one of the extensions of w to L_v ramifies (see [ZS], Ch. VI §11 Cor. 2 to Lem. 4). Now we use the induction assumption to find a geometric discrete valuation v_2 of Kv_1 of rank 1 which ramifies in L_v . So $v_1 \circ v_2$ is a geometric discrete rank 2 valuation on K which ramifies in the extension $L|K$. Using the induction assumption again, we find a geometric discrete rank 1 valuation on K which ramifies in $L|K$. \square

Lemma 3.6. *Let L, K and d be as in Lemma 3.5. Assume there exists a wildly ramified discrete rank d valuation v on K . Then there exists a wildly ramified geometric discrete rank 1 valuation w on K .*

Proof. Let p be the characteristic of Kv . We may replace $L|K$ by its Galois closure. Since the ramification group of v is p -group, there exists an intermediate extension $K \subset K' \subset L' \subset L$ such that $L'|K'$ is Galois of prime order p and such that an extension v' of v to K' is wildly ramified in $L'|K'$. Choose a proper normal model X of K' over S . By the Lemma 3.5, we find a discrete rank 1 valuation w' of K' which ramifies in $L'|K'$. Choose a point $x \in Z_{w'}$ of codimension 2 in X and of residue characteristic p . Set $A = \mathcal{O}_{X,x}$. By Proposition 2.1, we may assume that A is regular and that the ramification locus of $L'|K'$ in $\text{Spec}(A)$ has normal crossings. Then Abhyankar's Lemma [SGA1, XIII Proposition 5.2] applied to the strict henselization of A shows that there is a ramified prime divisor on $\text{Spec}(A)$ of function field characteristic p . Let w'' be the discrete rank 1 valuation corresponding to this ramified prime divisor. Then the restriction w of w'' to K is a wildly ramified geometric discrete rank 1 valuation. \square

4 Curve-, Divisor- and Chain-Tameness

We start with the following applications of our Key Lemma 2.4.

Proposition 4.1. *Let X be a regular, pure-dimensional, excellent scheme, $X' \subset X$ a dense open subscheme, $Y' \rightarrow X'$ an étale covering and Y the normalization of X in $k(Y')$. Suppose that for every curve C on X with $C' = C \cap X' \neq \emptyset$, the étale covering $Y' \times_X C' \rightarrow X' \times_X C'$ extends to an étale covering of \tilde{C} . Then $Y \rightarrow X$ is étale.*

Proof. Without loss of generality we can assume that $Y' \rightarrow X'$ is a Galois covering. Assume $Y \rightarrow X$ were not étale. We have to find a curve C on X with $C' = C \cap X' \neq \emptyset$ such that $Y' \times_{X'} \tilde{C}' \rightarrow \tilde{C}'$ is ramified along $\tilde{C} \setminus \tilde{C}'$. By the purity of the branch locus [SGA2, X.3.4], there exists a component D of $X \setminus X'$ of codimension one in X such that $Y \rightarrow X$ is ramified over the generic point of D . Let G be a cyclic subgroup of prime order of the inertia group of some point of Y which lies over the generic point of D . Let Y'_G be the quotient of Y' by the action of G . Consider the Galois covering $Y' \rightarrow Y'_G$ of prime degree and let Y_G be the normalization of X in $k(Y'_G)$. By considering the localization at any closed point of Y_G lying over D , Lemma 2.4 produces a curve C_G on Y_G with $C'_G = C_G \cap Y'_G \neq \emptyset$ such that $Y' \times_{Y'_G} C'_G \rightarrow C'_G$ is ramified along $\tilde{C}_G \setminus \tilde{C}'_G$. Let C be the image of C_G under the morphism $Y_G \rightarrow X$. Then C is the curve we are looking for. \square

Proposition 4.2. *Let X be a normal, pure-dimensional, excellent scheme and let D be a NCD on X (in particular, X is regular in a neighbourhood of D). Then an étale covering $Y' \rightarrow X' := X \setminus D$ is tamely ramified along D if and only if for each closed curve $C \subset X$ not contained in D the base change $Y' \times_X C' \rightarrow \tilde{C}'$ is tamely ramified along $D_{\tilde{C}}$.*

Proof. If $Y' \rightarrow X' := X \setminus D$ is tamely ramified along D and C is a curve on X we use Abhyankar's Lemma to show that $Y' \times_X \tilde{C}' \rightarrow \tilde{C}'$ is tamely ramified along $D_{\tilde{C}}$. Let x be a point of the intersection $D \cap C$ and A the strict henselization of $\mathcal{O}_{X,x}$. We can replace X by $\text{Spec}(A)$. Then $|D| = V(\pi_1) \cup \cdots \cup V(\pi_r)$ for some irreducible elements $\pi_1, \dots, \pi_r \in A$ which are part of a parameter system of A and $X' = \text{Spec}(A')$ with $A' = A_{\pi_1 \dots \pi_r}$. By Abhyankar's Lemma [SGA1, XIII Proposition 5.2], we can assume without loss of generality that $Y' \rightarrow X'$ is a Galois covering of the form

$$\text{Spec}(A'[T_1, \dots, T_r] / (T_1^{n_1} - \pi_1, \dots, T_r^{n_r} - \pi_r)) \rightarrow \text{Spec}(A')$$

for natural numbers n_1, \dots, n_r which are prime to the residue characteristic p of A . So, in particular, p does not divide $\deg(Y'|X') = n_1 \cdots n_r$. In this situation the degree of the Galois covering $Y' \times_X \tilde{C}' \rightarrow \tilde{C}'$ divides $\deg(Y'|X')$, so it must be tamely ramified along $D_{\tilde{C}}$.

For the reverse direction assume that $Y' \rightarrow X' := X \setminus D$ is not tamely ramified along some irreducible component D_0 of D . We will construct a curve C on X such that the base change $Y' \times_X \tilde{C}' \rightarrow \tilde{C}'$ is not tamely ramified along $D_0 \times_X \tilde{C}$. We can replace Y' by its Galois closure over X' . Let G be a subgroup of prime order of the wild ramification group of $Y' \rightarrow X'$ at some point over the generic point of D_0 . Let Y'_G be the quotient of Y' under the action of G and let Y_G be the normalization of X in $k(Y'_G)$. Then a localization of $Y' \rightarrow Y'_G \subset Y_G$ at a closed point of Y_G lying over D_0 satisfies the assumptions of Lemma 2.4, so that we find a curve $C_G \subset Y_G$ with $C'_G = C_G \cap Y'_G \neq \emptyset$ such that $Y' \times_X \tilde{C}'_G \rightarrow \tilde{C}'_G$ is not tamely ramified along $D_0 \times_X \tilde{C}'_G$. Finally, let the curve C we are searching for be the image of C_G under the morphism $Y_G \rightarrow X$. \square

Let S be an integral, pure-dimensional and excellent base scheme and $\text{Sch}(S)$ the category of separated schemes of finite type over S .

We call $C \in \text{Sch}(S)$ a *curve* if C is integral and $\dim_S C = 1$. For a regular curve $C \in \text{Sch}(S)$ there exists a unique regular curve $P(C) \in \text{Sch}(S)$ which is proper over S and contains C as a dense open subscheme. Note that $P(C)$ has Krull-dimension 1. So there is a unique notion of tameness for étale coverings of regular curves in $\text{Sch}(S)$. The next definition is motivated by Proposition 4.2. It is the 'maximal' definition of tameness which is stable under base change and extends the given one for curves.

Definition. Let $Y \rightarrow X$ be an étale covering in $\text{Sch}(S)$. We say that $Y \rightarrow X$ is *curve-tame* if for any morphism $C \rightarrow X$ in $\text{Sch}(S)$ with C a regular curve, the base change $Y \times_X C \rightarrow C$ is tamely ramified along $P(C) \setminus C$.

Below we will introduce versions of tameness which use Parshin chains on schemes. For the convenience of the reader we recall the definition of Parshin chains and their connection with valuations.

Definition. Let $X \in \text{Sch}(S)$ be a scheme. A finite family of points $P = (P_0, \dots, P_r)$ on the scheme X is called a *chain* if

$$\overline{\{P_0\}} \subset \overline{\{P_1\}} \subset \cdots \subset \overline{\{P_r\}}.$$

The chain P is called a *Parshin chain* if $\dim_S \overline{\{P_i\}} = i$ for $0 \leq i \leq r$.

Assume we are given a scheme $X \in \text{Sch}(S)$ and a Parshin chain $P = (P_0, \dots, P_d)$ of length $d = \dim_S(X)$ on X . We say that a discrete valuation $v \in \text{Val}_S(k(X))$ of rank d *dominates* the Parshin chain P if, for the unique decomposition $v = v_1 \circ \cdots \circ v_d$ into discrete valuations of rank 1 and $i = 1, \dots, d$, the valuation ring

corresponding to $v_1 \circ \dots \circ v_i$ dominates $\mathcal{O}_{X, P_{d-i}}$. A Parshin chain P is dominated by at least one and by at most finitely many discrete valuations. If, for $i = 1, \dots, d$, P_{i-1} is a regular point on $\overline{\{P_i\}}$, then there is exactly one such discrete valuation.

We introduce further definitions of tameness. Let $Y \rightarrow X$ be an étale covering of connected normal schemes in $Sch(S)$. Then every $v \in \text{Val}_S(k(X))$ with center in X is unramified in $k(Y)|k(X)$.

Definition. Assume that Y and X are connected and normal. We say that $Y \rightarrow X$ is

- *divisor-tame* if for every normal compactification \bar{X} of X and every point $x \in \bar{X} \setminus X$ with $\text{codim}_{\bar{X}} x = 1$ the discrete rank one valuation v_x on $k(X)$ associated with x is tamely ramified in $k(Y)|k(X)$.
- *valuation-tame* if every $v \in \text{Val}_S(k(X))$ is tamely ramified in $k(Y)|k(X)$.
- *discrete-valuation-tame*, if every discrete valuation $v \in \text{Val}_S(k(X))$ of rank $d = \dim_S X$ is tamely ramified in $k(Y)|k(X)$.
- *chain-tame* if there exists a normal compactification \bar{X} of X such that each discrete valuation $v \in \text{Val}_S(k(X))$ of rank $d = \dim_S X$ which dominates a Parshin-chain on \bar{X} is tamely ramified in $k(Y)|k(X)$.
- *weakly chain-tame* if there exists a normal compactification \bar{X} of X such that each discrete valuation $v \in \text{Val}_S(k(X))$ of rank $d = \dim_S X$ which dominates a Parshin-chain $P = (P_0, \dots, P_d)$ on \bar{X} with $P_1, \dots, P_d \in X$ is tamely ramified in $k(Y)|k(X)$.

These definitions extend to the non-connected case by requiring the corresponding property for every connected component.

Valuation-tameness obviously implies discrete-valuation-tameness and divisor-tameness. Discrete-valuation-tameness implies chain-tameness, which implies weak chain-tameness.

Remark 4.3. The question whether an étale scheme morphism $Y \rightarrow X$ is tame or not (in any of the above versions), depends on the category $Sch(S)$ in which it is considered. For example, the étale morphism $\text{Spec}(\mathbb{Z}[\frac{1}{2}, \sqrt{-1}]) \rightarrow \text{Spec}(\mathbb{Z}[\frac{1}{2}])$ is not tame in $Sch(\mathbb{Z})$, but is tame as a morphism in $Sch(\mathbb{Z}[\frac{1}{2}])$. Another example is the following: any étale covering $Y \rightarrow X$ of varieties over \mathbb{Q}_p is tame when considered in $Sch(\mathbb{Q}_p)$. This is in general not the case if we consider $Y \rightarrow X$ as a covering in $Sch(\mathbb{Z}_p)$.

Now we formulate our first main result.

Theorem 4.4. *Let S be an integral, excellent and pure-dimensional base scheme and let $Y \rightarrow X$ be an étale covering of regular schemes in $Sch(S)$. Then the following are equivalent:*

- (i) $Y \rightarrow X$ is curve-tame.
- (ii) $Y \rightarrow X$ is divisor-tame.
- (iii) $Y \rightarrow X$ is discrete-valuation-tame.
- (iv) $Y \rightarrow X$ is chain-tame.
- (v) $Y \rightarrow X$ is weakly chain-tame.

If every intermediate field between $k(X)$ and the Galois closure of $k(Y)$ over $k(X)$ admits a regular proper model, then then (i)–(v) are equivalent to

- (vi) $Y \rightarrow X$ is valuation-tame.

If there exists a regular compactification \bar{X} of X such that $\bar{X} \setminus X$ is a NCD, then (i)–(vi) are equivalent and there is a further equivalent condition:

(vii) $Y \rightarrow X$ is tamely ramified along $\bar{X} \setminus X$.

Proof. We start making the round. (i) \Rightarrow (ii) follows from the Key Lemma 2.4. Let us show (ii) \Rightarrow (iii). Assume there exists a wildly ramified discrete valuation of rank $d = \dim_S X$. Using Lemma 3.5, we find a wildly ramified geometric discrete rank 1 valuation v on $k(X)$. Choose any normal compactification \bar{X} of X . As $Y \rightarrow X$ is étale, the center of v lies in $\bar{X} \setminus X$. By [Liu] §8, ex. 3.14, after blowing up \bar{X} in centers contained in $\bar{X} \setminus X$ and finally normalizing, we find a normal compactification \bar{X} of X such that v is the valuation associated to a point $x \in \bar{X} \setminus X$ of codimension 1 in \bar{X} . This shows (ii) \Rightarrow (iii).

Furthermore, (iii) \Rightarrow (iv) and (iv) \Rightarrow (v) are obvious. So assume (iv). Let \bar{X} be a compactification of X and let $C \rightarrow X$ be a morphism in $Sch(S)$, where C is a regular curve. If the image of C in X is a closed point, then $Y \times_X C \rightarrow C$ extends to an étale morphism of $P(C)$. So assume that the image x of the generic point of C in X is one-dimensional and that the base change $Y \times_X C \rightarrow C$ is wildly ramified along $P(C) \setminus C$. By the valuative criterion of properness, $C \rightarrow X$ extends to a morphism $P(C) \rightarrow \bar{X}$. Let v be a discrete rank 1 valuation on $k(C)$ with center in $P(C) \setminus C$ which is wildly ramified in $Y \times_X C \rightarrow C$. Let w be the restriction of v to $k(x)$ with respect to the inclusion $k(x) \subset k(C)$. Then w is wildly ramified in $Y \otimes k(x) \rightarrow k(x)$. Let $P_0 \in \bar{X}$ be the center of w . Since x is a regular point on X , we find a Parshin chain P_0, P_1, \dots, P_d on \bar{X} such that $P_1 = x$ and P_i is a regular point on \bar{P}_{i+1} for $i = 1, \dots, d-1$. Then let W be the geometric rank $d-1$ -valuation associated to P_1, \dots, P_d . Then $W \circ w$ is a wildly ramified discrete rank d -valuation. This shows (v) \Rightarrow (i).

Obviously, (vi) implies (iii). Now assume that (v) holds and that every intermediate field between $k(X)$ and the Galois closure of $k(Y)$ over $k(X)$ admits a regular proper model. In order to show (vi), we may replace Y by its Galois closure over X . Assume that there exists a valuation $v \in \text{Val}_S(k(X))$ which is wildly ramified in $k(Y)|k(X)$. Let p be the residue characteristic of v and let $H \subset G(k(Y)|k(X))$ be a cyclic subgroup of order p contained in the ramification group of v . We replace X by the quotient scheme Y/H , i.e. we may assume that $Y|X$ is cyclic of order p . Let X' be a regular proper model of $k(X)$ and let x' be the center of v in X' . Then x' is ramified in $Y' \rightarrow X'$. By purity of the branch locus, we find a prime divisor $D \subset X'$ which contains x' and is ramified in $Y' \rightarrow X'$. Now choose a Parshin-chain from x to D and a discrete rank d valuation which dominates this chain. This discrete rank d valuation is wildly ramified in $k(Y)|k(X)$. Hence (iii) is violated.

Finally, assume that there exists a regular compactification \bar{X} such that $\bar{X} \setminus X$ is a normal crossing divisor. Then (vii) is equivalent to (i) by Proposition 4.2. In the next section we will see (Theorem 5.4) that in this situation $Y \rightarrow X$ is numerically tame along $\bar{X} \setminus X$, and therefore valuation-tame by Theorem 5.3. So (vii) implies (vi). \square

Remark 4.5. If the scheme X in Theorem 4.4 is only assumed to be normal instead of regular, one still gets the implications

$$(i) \Rightarrow (ii) \Leftrightarrow (iii) \Rightarrow (iv) \Rightarrow (v).$$

5 Numerical tameness

Let X be a connected normal scheme with function field $K = k(X)$ and let $L|K$ be a finite Galois extension with Galois group G . Let Y be the normalization of X in L and let $y \in Y$ be a (not necessarily closed) point with image $x \in X$. Recall that the decomposition and the inertia group $G_y(Y|X)$ and $T_y(Y|X)$ of y in $Y|X$ are defined by

$G_y(Y|X) = \{g \in G \mid gy = y\}$ and $T_y(Y|X) = \ker(G_y(Y|X) \rightarrow \text{Aut}(k(y)|k(x)))$, respectively. If y and y' have the same image $x \in X$, then the decomposition and inertia groups of y and y' are conjugate and we sometimes write $G_x(Y|X)$ and $T_x(Y|X)$ for these groups if we don't care for conjugation.

Next we introduce the notion of numerical tameness.

Definition. Let $\bar{X} \in \text{Sch}(S)$ be normal connected and proper, and let $X \subset \bar{X}$ be a dense open subscheme. Let $Y \rightarrow X$ be an étale Galois covering and let \bar{Y} be the normalization of \bar{X} in the function field $k(Y)$ of Y . We say that $Y \rightarrow X$ is *numerically tamely ramified along $\bar{X} \setminus X$* if the order of the inertia group $T_x(\bar{Y}|\bar{X}) \subset G(\bar{Y}|\bar{X}) = G(Y|X)$ of each point $x \in \bar{X} \setminus X$ is prime to the residue characteristic of x . An étale covering $Y \rightarrow X$ is called *numerically tamely ramified along $\bar{X} \setminus X$* if it can be dominated by a Galois covering which is numerically tamely ramified along $\bar{X} \setminus X$. This definition extends to the non-connected case by requiring numerical tame ramification for all connected components.

Remark 5.1. If a point x_0 lies in the closure of another point x_1 , then $T_{x_1}(Y|X) \subset T_{x_0}(Y|X)$. It therefore suffices to check numerical tameness on closed points.

Numerical tameness is stable under base change in the following sense.

Lemma 5.2. *Let $\bar{X} \in \text{Sch}(S)$ be normal and proper, $X \subset \bar{X}$ a dense open subscheme and $Y \rightarrow X$ an étale covering. Let $f : \bar{X}' \rightarrow \bar{X}$ be a proper morphism in $\text{Sch}(S)$ with \bar{X}' normal such that $X' := f^{-1}(X)$ is dense in \bar{X}' . If $Y \rightarrow X$ is numerically tamely ramified along $\bar{X} \setminus X$, then the base change $Y \times_X \bar{X}' \rightarrow \bar{X}'$ is numerically tamely ramified along $\bar{X}' \setminus X'$.*

Proof. This follows since the inertia groups of the base change are subgroups of the inertia groups of $Y \rightarrow X$. \square

Theorem 5.3. *Let $X \in \text{Sch}(S)$ be a regular scheme, \bar{X} a normal compactification and $Y \rightarrow X$ an étale covering. If $Y \rightarrow X$ is numerically tamely ramified along $\bar{X} \setminus X$, then it is valuation-tame. In particular, $Y \rightarrow X$ is curve-, divisor- and chain-tame.*

Proof. We may suppose that $Y \rightarrow X$ is Galois. Let v be an S -valuation on $k(X)$ with residue characteristic $p > 0$. Let $x \in \bar{X}$ be a closed point the center of v . Then x has residue characteristic p and we have an inclusion $T_v(Y|X) \subset T_x(Y|X)$. Hence $T_v(Y|X)$ is of order prime to p and v is tamely ramified. \square

A partial converse in the case that \bar{X} is regular is given by the following

Theorem 5.4. *Let $\bar{X} \in \text{Sch}(S)$ be a regular, proper scheme, $X \subset \bar{X}$ a dense open subscheme and $Y \rightarrow X$ an étale covering. Assume that one of the following conditions is satisfied:*

- (a) $\bar{X} \setminus X$ is a NCD and $Y \rightarrow X$ is tamely ramified along $\bar{X} \setminus X$, or
- (b) $Y \rightarrow X$ is curve-tame and can be dominated by a Galois covering with nilpotent Galois group.

Then $Y \rightarrow X$ is numerically tamely ramified along $\bar{X} \setminus X$.

Remark 5.5. The equivalence of numerical tameness and chain-tameness for nilpotent coverings has been shown in [Sc1]. Wiesend has given an incomplete proof of the equivalence of numerical tameness and curve-tameness for nilpotent coverings in [Wi]. In the appendix we will give examples for curve-tame Galois coverings $Y \rightarrow X$ with non-nilpotent Galois group which are not numerically tamely ramified along $\bar{X} \setminus X$ for some regular compactification \bar{X} . The first such example has been given by the referee of [Wi].

Proof. Assume that $\bar{X} \setminus X$ is a NCD and $Y \rightarrow X$ is tamely ramified along $\bar{X} \setminus X$. By Abhyankar's Lemma [SGA1, XIII Proposition 5.2], the inertia group of every closed point $x \in \bar{X}$ has order prime to the residue characteristic of x . Hence $Y|X$ is numerically tamely ramified along $\bar{X} \setminus X$.

Now assume that $\bar{X} \setminus X$ is not necessarily a NCD but that $Y \rightarrow X$ can be dominated by a Galois covering with nilpotent group. Since a finite nilpotent group is the product of its Sylow subgroups, we may assume that $Y \rightarrow X$ is Galois and that $G = \text{Gal}(Y|X)$ is a finite p -group, where p is some prime number. Assume that $Y|X$ were not numerically tamely ramified along $\bar{X} \setminus X$. Then we find a closed point in $x_0 \in \bar{X} \setminus X$ with residue characteristic p which ramifies in $\bar{Y}|\bar{X}$. Now we factor $Y \rightarrow X$ in the form

$$Y = X_n \rightarrow X_{n_1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = X,$$

such that $X_{i+1} \rightarrow X_i$ is Galois of degree p for $i = 0, \dots, n-1$. We denote by \bar{X}_i the normalization of \bar{X} in $k(X_i)$. Let $a, 0 \leq a \leq n-1$, be the unique index such that $\bar{X}_a|\bar{X}$ is étale over x_0 but $\bar{X}_{a+1}|\bar{X}_a$ is not étale over the preimage x_0 . Note that \bar{X}_a is regular in a neighbourhood of the preimage of x_0 , since there is a neighbourhood which is étale over \bar{X} . By purity, there exists a prime divisor on \bar{X}_a which meets the preimage of x_0 in a closed point x_a and which ramifies in $\bar{X}_{a+1}|\bar{X}_a$. Applying the Key Lemma 2.4 to the localization of $\bar{X}_{a+1}|\bar{X}_a$ at x_a , we find a curve $\bar{C} \subset \bar{X}_a$ containing x_a and with $C := \bar{C} \cap X_a \neq \emptyset$, such that the base change $X_{a+1} \times_{X_a} \bar{C} \rightarrow \bar{C}$ ramifies along some point over x_a . We conclude that $X_{a+1}|X_a$ is not curve-tame. Hence $Y|X$ is not curve-tame. A contradiction. \square

6 Cohomological tameness

Let $\bar{X} \in \text{Sch}(S)$ be normal connected and proper, and let $X \subset \bar{X}$ be a dense open subscheme. Let $Y \rightarrow X$ be an étale Galois covering with group $G = \text{Gal}(Y|X)$ and let \bar{Y} be the normalization of \bar{X} in the function field $k(Y)$ of Y . We denote the projection by $\pi : \bar{Y} \rightarrow \bar{X}$.

Lemma 6.1. *Using the notation as above let $(U_i = \text{Spec}(A_i))_{i \in I}$ be an affine Zariski-open covering of \bar{X} and put $\pi^{-1}(U_i) =: V_i = \text{Spec}(B_i)$. Then the following are equivalent.*

- (i) For every closed point $x \in \bar{X} \setminus X$ the semi-local ring $\mathcal{O}_{\bar{Y}, \pi^{-1}(x)}$ is a cohomologically trivial G -module.
- (ii) For every point $x \in \bar{X}$ the semi-local ring $\mathcal{O}_{\bar{Y}, \pi^{-1}(x)}$ is a cohomologically trivial G -module.
- (iii) B_i is a cohomologically trivial G -module for all i .

Proof. For any prime ideal $\mathfrak{p} \in A_i$, the localization $A_i \rightarrow (A_i)_{\mathfrak{p}}$ is flat. Hence for every subgroup $H \subset G$ and all $i \in \mathbb{Z}$, we have

$$\hat{H}^i(H, B_i \otimes_{A_i} (A_i)_{\mathfrak{p}}) \cong \hat{H}^i(H, B_i) \otimes_{A_i} (A_i)_{\mathfrak{p}},$$

where $\hat{H}^i(H, -)$ denotes the Tate cohomology of the finite group H . We conclude that B_i is a cohomologically trivial G -module if and only if $B_i \otimes_{A_i} (A_i)_{\mathfrak{p}}$ is cohomologically trivial for every prime ideal, or even for every maximal ideal. In order to conclude the proof, it remains to show, that $B_i \otimes_{A_i} (A_i)_{\mathfrak{p}}$ is a cohomologically trivial G -module for such \mathfrak{p} which are unramified B_i . To this end note that the natural homomorphism $(A_i)_{\mathfrak{p}} \rightarrow (A_i)_{\mathfrak{p}}^{sh}$ to the strict henselization is faithfully flat and that $B_i \otimes_{A_i} (A_i)_{\mathfrak{p}}^{sh}$ is an induced G -module if \mathfrak{p} is unramified. \square

Definition. We say that $Y \rightarrow X$ is *cohomologically tamely ramified along $\bar{X} \setminus X$* if the equivalent conditions of Lemma 6.1 are satisfied. An étale covering $Y \rightarrow X$ is called *cohomologically tamely ramified along $\bar{X} \setminus X$* if it can be dominated by a Galois covering which is cohomologically tamely ramified along $\bar{X} \setminus X$. This definition extends to the non-connected case by requiring cohomological tame ramification for all connected components.

Theorem 6.2. *Let $\bar{X} \in \text{Sch}(S)$ be normal, connected and proper, and let $X \subset \bar{X}$ be a dense open subscheme. Then an étale covering $Y \rightarrow X$ is cohomologically tamely ramified along $\bar{X} \setminus X$ if and only if it is numerically tamely ramified along $\bar{X} \setminus X$.*

Proof. Let $(U_i = \text{Spec}(A_i))_{i \in I}$ be an affine Zariski-open covering of \bar{X} and put $\pi^{-1}(U_i) =: V_i = \text{Spec}(B_i)$. Let $\text{tr}_{B_i|A_i} : B_i \rightarrow A_i$ be the trace map $b \mapsto \sum_{g \in G} gb$. We omit the indices i from the notation and call $B|A$ tamely ramified if for every maximal ideal $\mathfrak{m} \subset A$ the inertia group $T_{\mathfrak{m}}(B|A)$ is of order prime to the characteristic of A/\mathfrak{m} . The following claims are standard, cf. [NSW], Theorem 6.1.10.

Claim 1: $\text{tr}_{B|A} : B \rightarrow A$ is surjective if and only if $B|A$ is cohomologically tamely ramified.

Proof of claim 1. For a maximal ideal $\mathfrak{m} \subset A$, we denote the henselization of $A_{\mathfrak{m}}$ by $A_{\mathfrak{m}}^h$. Recall that $A_{\mathfrak{m}} \rightarrow A_{\mathfrak{m}}^h$ is faithfully flat. Hence $\text{tr}_{B|A} : B \rightarrow A$ is surjective if and only if $\text{tr}_{B|A} \otimes_A A_{\mathfrak{m}}^h : B \otimes_A A_{\mathfrak{m}}^h \rightarrow A_{\mathfrak{m}}^h$ is surjective for all $\mathfrak{m} \subset A$. Therefore we may assume that A is local henselian with maximal ideal \mathfrak{m} . Then B is a finite product of local henselian rings and we may reduce to the case that B is local henselian with maximal ideal \mathfrak{m}_B . Let κ and κ' be the residue fields of A and B , respectively. By Nakayama's Lemma, $\text{tr}_{B|A} : B \rightarrow A$ is surjective if and only if the induced map $B/\mathfrak{m}_B \rightarrow A/\mathfrak{m}$ is surjective. Since the trace sends \mathfrak{m}_B to \mathfrak{m} , this map factors through a map $\text{tr} : \kappa' \rightarrow \kappa$. Let p be the residue characteristic. By definition, the inertia group acts trivially on κ' . Hence tr is the zero map if $p \mid \#T_{\mathfrak{m}}(B|A)$. If $e = \#T_{\mathfrak{m}}(B|A)$ is prime to p , then $\kappa'|\kappa$ is separable and $\text{tr} = e \cdot \text{tr}_{\kappa'|\kappa}$ is surjective. This shows the claim.

Claim 2: B is a cohomologically trivial G -module if and only if $B|A$ is cohomologically tamely ramified.

Proof of claim 2. If B is a cohomologically trivial G -module, then $tr_{B|A}$ is surjective since $0 = \hat{H}^0(G, B) = A/tr_{B|A}(B)$. Hence $B|A$ is cohomologically tamely ramified by Claim 1. If $B|A$ is cohomologically tamely ramified and $H \subset G$ is a subgroup, then also $B|B^H$ is cohomologically tamely ramified. Hence $tr_{B|B^H}$ is surjective by Claim 1, which implies $\hat{H}^0(H, B) = 0$.

Next we prove that $H^1(H, B) = 0$. Let $a(\sigma) \in B$ be a 1-cocycle and let $x \in B$ be such that $tr_{B|A}(x) = 1$. Setting

$$b := \sum_{\sigma \in H} a(\sigma)\sigma x,$$

we obtain for $\tau \in H$,

$$\tau b = \sum_{\sigma \in H} \tau a(\sigma)(\tau\sigma x) = \sum_{\sigma \in H} (a(\tau\sigma) - a(\tau))(\tau\sigma x) = b - a(\tau)tr_{B|A}(x).$$

Therefore $a(\tau) = (1 - \tau)b$, hence $a(\tau)$ is a 1-coboundary. By [NSW], Proposition 1.8.4, we conclude that B is cohomologically trivial. This shows Claim 2.

Now we show the theorem. If $Y \rightarrow X$ is numerically tamely ramified, then B_i is a cohomologically trivial G -module for all i by Claim 2. Hence $Y \rightarrow X$ is cohomologically tamely ramified by Lemma 6.1. The same arguments also show the reverse direction. \square

7 Finiteness theorems

In this section we will prove a tame version of a finiteness result due to Katz and Lang [KL] and will deduce finiteness theorems for the abelianized tame fundamental groups of arithmetic schemes, which had been previously shown in [Sc2].

Let, as before, S be an integral, pure-dimensional and excellent base scheme. Let $X \in Sch(S)$ be regular and connected. We use the word tame for the equivalent notions of curve-, divisor-, discrete-valuation- and chain-tameness and *tame covering* means a finite, étale morphism which is tame. The tame coverings of X satisfy the axioms of a Galois category ([SGA1], V, 4). After choosing a geometric point \bar{x} of X , we have the fibre functor $(Y \rightarrow X) \mapsto Mor_X(\bar{x}, Y)$ from the category of tame coverings of X to the category of sets, whose automorphism group is called the **tame fundamental group** $\pi_1^t(X/S, \bar{x})$. It classifies étale coverings of X which are tame when considered in $Sch(S)$. Denoting the usual étale fundamental group by $\pi_1(X, \bar{x})$, we have an obvious surjection

$$\pi_1(X, \bar{x}) \twoheadrightarrow \pi_1^t(X/S, \bar{x}),$$

which is an isomorphism if X is proper over S . The fundamental groups to different base points are isomorphic, the isomorphism being canonical up to inner automorphisms. Therefore the abelianized fundamental groups

$$\pi_1^{ab}(X, \bar{x}) \text{ and } \pi_1^{t,ab}(X/S, \bar{x})$$

are canonically independent of the base point \bar{x} , which we will exclude from notation. Note that the notion of (curve-) tameness is stable under base change. Therefore, given a morphism $f : Y \rightarrow X$ between regular, connected schemes in $Sch(S)$, a geometric point \bar{y} of Y and its image $\bar{x} = f(\bar{y})$ in X , we obtain an induced homomorphism

$$\pi_1^t(Y/S, \bar{y}) \longrightarrow \pi_1^t(X/S, \bar{x}).$$

Now assume that the function field k of S is absolutely finitely generated and let $p = \text{char}(k)$. Consider the kernel

$$\ker(Y/X) := \ker(\pi_1^{ab}(Y) \rightarrow \pi_1^{ab}(X))$$

If $f : Y \rightarrow X$ is smooth and surjective, then a theorem of Katz and Lang ([KL], Theorem 1) asserts that the prime-to- p part of $\ker(Y/X)$ is finite. We will prove an analogous result for the tame kernel

$$\ker_S^t(Y/X) := \ker(\pi_1^{t,ab}(Y/S) \rightarrow \pi_1^{t,ab}(X/S)),$$

which does not exclude the p -part.

Theorem 7.1. *Let S be an integral, pure-dimensional and excellent base scheme whose function field is absolutely finitely generated. Let $f : Y \rightarrow X$ be a smooth, surjective morphism of connected regular schemes in $\text{Sch}(S)$. Assume that the generic fibre of f is geometrically connected and that one of the following conditions (i) and (ii) is satisfied:*

- (i) *X has a regular compactification \bar{X} over S such that the boundary $\bar{X} \setminus X$ is a normal crossing divisor.*
- (ii) *The generic fibre of f has a rational point.*

Then the group

$$\ker_S^t(Y/X) := \ker(\pi_1^{t,ab}(Y/S) \rightarrow \pi_1^{t,ab}(X/S))$$

is finite.

Remark 7.2. Condition (i) is satisfied if X has Krull-dimension less or equal to 2; conjecturally, it is satisfied for an arbitrary X .

In the proof of Theorem 7.1 we will need the following

Lemma 7.3. *Let A be a discrete valuation ring and let v be the associated discrete rank one valuation on the quotient field k of A . Let $K|k$ be a finitely generated, regular field extension and let $w \in \text{Val}_{\text{Spec}(A)}(K)$ be a geometric discrete rank one valuation on K extending v . Let $\pi \in k$ be a uniformizer for v and let $k'|k$ be a finite Galois extension such that w is unramified in $Kk'|K$. Then*

$$\#T_v(k'|k) \mid w(\pi) \cdot [(kv)_{Kw} : kv]_i,$$

where $[(kv)_{Kw} : kv]_i$ is the index of inseparability of the relative algebraic closure $(kv)_{Kw}$ of kv in Kw (which is finite over kv by Proposition 3.2).

Proof. Without loss of generality we may assume that A is henselian. Let v' be the unique extension of v to k' . Then $\#T_v(k'|k) = e_v(k'|k)[k'v' : kv]_i$. Let $B \subset K$ be the valuation ring of w and let B' be the integral closure of B in Kk' . Then B' is a semi-local principal ideal domain. Let \mathfrak{m} be the maximal ideal of B and let $\mathfrak{m}'_1, \dots, \mathfrak{m}'_g$ be the maximal ideals of B' . By assumption, $B \hookrightarrow B'$ is étale, hence $\mathfrak{m}B' = \mathfrak{m}'_1 \cdots \mathfrak{m}'_g$. This implies

$$\pi B' = \mathfrak{m}^{w(\pi)} B' = (\mathfrak{m}'_1 \cdots \mathfrak{m}'_g)^{w(\pi)},$$

But $\pi B' = (\pi')^{e_v(k'|k)} B'$, where $\pi' \in k'$ is a uniformizer of v' , and so we obtain $e_v(k'|k) \mid w(\pi)$.

Furthermore, for all i the extension $(B'/\mathfrak{m}'_i)|(B/\mathfrak{m}) = Kw$ is separable by assumption. As B'/\mathfrak{m}'_i contains $k'v'$, we conclude that

$$[k'v' : kv]_i \mid [(kv)_{Kw} : kv].$$

Summing up, the result follows. \square

Proof of Theorem 7.1. In the proof we will make frequent use of the following well-known fact: Let X be a normal, connected locally noetherian scheme with generic point η and function field K . Let K^{sep} be a separable closure of K and let $\bar{\eta}$ be the corresponding geometric point of X . Then $\pi_1(X, \bar{\eta})$ is a quotient of the Galois group $G(K^{\text{sep}}|K)$ and the functor “fibre over η ”

$$\{\text{connected étale coverings of } X\} \longrightarrow \{\text{finite separable extensions of } K\}$$

is fully faithful, with image those finite separable extensions $K'|K$ for which the normalization of X in K' is étale over X . With the obvious modifications, the same also holds for the tame fundamental group.

First step: Reduction to $S = X = \text{Spec}(K)$.

Let $\eta \in X$ be the generic point and let $K := k(\eta)$ be its function field. To accomplish this step it clearly suffices to prove the following claim:

Claim: *The natural map $\ker_{\eta}^t(Y_{\eta}/\eta) \rightarrow \ker_S^t(Y/X)$ has finite cokernel.*

Proof of the Claim: Let L be the function field of Y . We choose a separable closure L^{sep} of L and denote the separable closure of K in L by K^{sep} . By assumption, L and K^{sep} are linearly disjoint extensions of K . Therefore the natural map $\pi_1^{t,ab}(Y_{\eta}/\eta) \rightarrow G(K^{\text{sep}}|K)^{ab}$ is surjective. The exact sequence

$$1 \rightarrow \pi_1^t(Y_{\bar{\eta}}/\bar{\eta}) \rightarrow \pi_1^t(Y_{\eta}/\eta) \rightarrow G(K^{\text{sep}}|K) \rightarrow 1$$

implies the exact sequence

$$(*) \quad \pi_1^{t,ab}(Y_{\bar{\eta}}/\bar{\eta})_{G(K^{\text{sep}}|K)} \rightarrow \pi_1^{t,ab}(Y_{\eta}/\eta) \rightarrow G(K^{\text{sep}}|K)^{ab} \rightarrow 0$$

Therefore $\ker_{\eta}^t(Y_{\eta}/\eta)$ is a quotient of $\pi_1^{t,ab}(Y_{\bar{\eta}}/\bar{\eta})_{G(K^{\text{sep}}|K)}$. Consider the commutative diagram

$$\begin{array}{ccccc} \pi_1^{t,ab}(Y_{\bar{\eta}}/\bar{\eta}) & \longrightarrow & \pi_1^{t,ab}(Y_{\eta}/\eta) & \twoheadrightarrow & G(K^{\text{sep}}|K)^{ab} \\ \parallel & & \downarrow & & \downarrow \\ \pi_1^{t,ab}(Y_{\bar{\eta}}/\bar{\eta}) & \longrightarrow & \pi_1^{t,ab}(Y/S) & \twoheadrightarrow & \pi_1^{t,ab}(X/S). \end{array}$$

The upper row is exact and the lower row is a complex. Denoting the cohomology of the lower row by H , we obtain an exact sequence

$$\ker_{\eta}^t(Y_{\eta}/\eta) \rightarrow \ker_S^t(Y/X) \rightarrow H.$$

It therefore remains to show that H is finite. In Galois terms, $H = G(K_2|K_1)$, where

- $K_1|K$ is the maximal abelian extension of K such that the normalization of X in K_1 is tame over X .
- $K_2|K$ is the maximal abelian extension of K such that the normalization of Y in LK_2 is tame over Y .

We consider the cases (i) and (ii) separately.

(i) Let \bar{X} be a regular compactification of X over S such that $\bar{X} \setminus X$ is a normal crossing divisor. Let v_1, \dots, v_n be the discrete rank one valuations of K associated to the irreducible components of $\bar{X} \setminus X$. Then, by Theorem 4.4, $H = G(K_2|K_1)$ is generated by the ramification groups of v_1, \dots, v_n in $G(K_2|K)$, and so it suffices to show that these groups are finite. Let v be one of the v_i and let p be the residue characteristic. If $p = 0$, there is nothing to prove, so assume $p > 0$. Let $K_2(p)$ be the maximal p -subextension of K in K_2 . We have to show that the inertia group of v in $K_2(p)|K$ is finite. Factor $Y \rightarrow \bar{X}$ into an open immersion $Y \subset \bar{Y}$ with dense image and normal \bar{Y} and a proper morphism $\bar{Y} \rightarrow \bar{X}$, which is possible for example by [Lue]. We find a codimension one point on \bar{Y} whose associated discrete valuation w of L extends v . Since the étale covering of Y associated to $LK_2(p)|L$ is divisor-tame, we see that w is unramified in $LK_2(p)|L$. Therefore $T_v(K_2(p)|K)$ is finite by Lemma 7.3.

(ii) As the generic fibre of f has a rational point, there exists a section $s : X_0 \rightarrow Y$ to $f : Y \rightarrow X$ over some dense open subscheme $X_0 \subset X$. If $X' \rightarrow X$ is an étale

covering such that $X' \times_X Y \rightarrow Y$ is tame, then, using the section s and the base change property of (curve-)tame coverings, we see that $X' \times_X X_0 \rightarrow X_0$ is tame. Taking the point of view of divisor-tameness, we conclude that $X' \rightarrow X$ is tame. Hence $H = 0$ in this case.

This completes the proof of the claim and of the first step. If $\text{char}(K) = 0$, we are ready, because in this case

$$\ker_{\eta}^t(Y/\eta) = \ker(Y/\eta)$$

is finite by [KL], Theorem 1. So we may assume that K has characteristic $p > 0$.

Second step: Reduction to the case that $f : Y \rightarrow X$ is an elementary fibration which admits a section $s : X \rightarrow Y$.

Suppose we are in the situation achieved by the first step, i.e. $X = S = \text{Spec}(K)$. We claim that it suffices to prove the finiteness of $\ker_K^t(Y/K)$ after a finite base change $K \subset K'$. In fact, using the exact sequence (*) of the first step, we see that the map $\ker_{K'}^t(Y \times_K K'/K') \rightarrow \ker_K^t(Y/K)$ is surjective, and this reduces us to show the finiteness of $\ker_{K'}^t(Y \times_K K'/K')$.

Let $Y' \rightarrow Y$ be étale and let K' be the algebraic closure of K in $k(Y')$. We claim that we may replace Y by Y' and K by K' . First note that $Y' \rightarrow Y$ factors through $Y' \rightarrow Y \times_K K'$, and so, by the last reduction, we may suppose that $K = K'$. Since the homomorphism $\pi_1^{ab}(Y') \rightarrow \pi_1^{ab}(Y)$ has finite cokernel, the same is true for $\pi_1^{t,ab}(Y'/K) \rightarrow \pi_1^{t,ab}(Y/K)$ and also for $\ker_K^t(Y'/K) \rightarrow \ker_K^t(Y/K)$. This shows the claim.

Recall that an elementary fibration is a complement of a finite étale divisor in a proper and smooth relative curve. By [SGA4, Exp. XI, 3.1], after replacing Y by an étale open, we can factor f into a sequence of elementary fibrations

$$Y = V_n \xrightarrow{f_n} V_{n-1} \xrightarrow{f_{n-1}} \cdots \xrightarrow{f_1} V_0 = X.$$

Moreover, since a surjective, smooth morphism admits a section over an étale open ([EGA4], IV, 17.16.3), we can achieve the existence of sections $s_i : V_{i-1} \rightarrow V_i$ to $f_i : V_i \rightarrow V_{i-1}$ for all i . Because the sequences

$$0 \longrightarrow \ker_K^t(V_i/V_{i-1}) \longrightarrow \ker_K^t(V_i/K) \longrightarrow \ker_K^t(V_{i-1}/K) \longrightarrow 0$$

are exact, it suffices to show the theorem for each elementary fibration in our sequence separately.

Last step: Completion of the proof.

After the second step we are reduced to the case in which $f : Y \rightarrow X$ is an elementary fibration which has a section. A further application of the procedure in the first step allows us to assume that again $S = X = \text{Spec}(K)$, where K is an absolutely finitely generated field K of characteristic $p > 0$. Let \tilde{Y} be the (uniquely defined) smooth compactification of the smooth curve Y over K . We have to show the finiteness of prime-to- p and of the p -primary part of $\ker_K^t(Y/K)$. The first is isomorphic to the prime-to- p part of $\ker(Y/K)$. The latter is isomorphic to the p -part of $\ker(\tilde{Y}/K)$. Therefore the proof of our theorem is finally reduced to [KL, Theorems 1 and 2]. \square

Using Theorem 7.1, one immediately deduces the following finiteness theorems for the tame fundamental group of arithmetic schemes, which had been shown previously in [Sc2].

Theorem 7.4. *If $X \in \text{Sch}(\mathbb{Z})$ is regular, connected and flat over $\text{Spec}(\mathbb{Z})$, then the abelianized tame fundamental group $\pi_1^{t,ab}(X/\text{Spec}(\mathbb{Z}))$ is finite.*

Proof. For a dense open subscheme $X_0 \subset X$, the homomorphism

$$\pi_1^{t,ab}(X_0/\text{Spec}(\mathbb{Z})) \longrightarrow \pi_1^{t,ab}(X/\text{Spec}(\mathbb{Z}))$$

is surjective. Therefore, making X smaller, we may assume that the structural morphism $X \rightarrow \text{Spec}(\mathbb{Z})$ is smooth. Let k be the algebraic closure of \mathbb{Q} in $k(X)$. Then $X \rightarrow \text{Spec}(\mathbb{Z})$ factors through $\text{Spec}(\mathcal{O}_k)$. Denoting the (open) image of X in $\text{Spec}(\mathcal{O}_k)$ by U , the morphism $X \rightarrow U$ is smooth and surjective with geometrically connected generic fibre. Now consider the exact sequence

$$0 \rightarrow \ker_{\text{Spec}(\mathbb{Z})}^t(X/U) \rightarrow \pi_1^{t,ab}(X/\text{Spec}(\mathbb{Z})) \rightarrow \pi_1^{t,ab}(U/\text{Spec}(\mathbb{Z})) \rightarrow 0.$$

The left hand group is finite by Theorem 7.1. The group $\pi_1^{t,ab}(U/\text{Spec}(\mathbb{Z}))$ is finite by class field theory: it is the Galois group of the ray class field with modulus $\prod_{\mathfrak{p} \notin U} \mathfrak{p}$ of the number field k . This finishes the proof. \square

If X is a smooth, connected variety over a finite field \mathbb{F} , then $\pi_1^{t,ab}(X/\text{Spec}(\mathbb{Z})) = \pi_1^{t,ab}(X/\text{Spec}(\mathbb{F}))$, and we have the degree map

$$\text{deg} : \pi_1^{t,ab}(X/\text{Spec}(\mathbb{F})) \longrightarrow \pi_1(\mathbb{F}) \cong \hat{\mathbb{Z}}.$$

The degree map has an open image, which corresponds to the field of constants of X , i.e. the algebraic closure of \mathbb{F} in $k(X)$. In this situation, Theorem 7.1 reads as

Theorem 7.5. *Let X be a smooth, connected variety over a finite field \mathbb{F} . Then $\ker(\text{deg})$ is finite. In particular,*

$$\pi_1^{t,ab}(X) \cong \hat{\mathbb{Z}} \oplus (\text{finite group}).$$

A Examples

Example 1: *We give an example which shows that the notion of tame ramification along a divisor on a regular scheme is not stable under base change if the divisor does not have normal crossings. This example is taken from [Sc1], Example 1.3.*

Let $X = \text{Spec}(\mathbb{Z}[T])$ be the affine line over $\text{Spec}(\mathbb{Z})$ and consider the divisor

$$D = \text{div}(T+4) + \text{div}(T-4),$$

which is not a normal crossing divisor. Let $K = \mathbb{Q}(T)$ be the function field of X and $U = X - D$. Put $f = (T+4)(T-4) = T^2 - 16$, $L = K(\sqrt{f})$ and consider the normalization X_L of X in L . The ramification locus of $X_L \rightarrow X$ is either D or $D \cup X_2$, where X_2 is the unique vertical divisor on X over characteristic 2. Let us show that $X_L \rightarrow X$ is unramified at the generic point of X_2 . This is equivalent to the statement that $L|K$ is unramified at the unique discrete valuation v_2 of K which corresponds to the prime ideal $2\mathbb{Z}[T] \subset \mathbb{Z}[T]$. Therefore it suffices to show that f is a square in the completion K_2 of K with respect to v_2 . Consider the polynomial $F(X) = X^2 - f = X^2 - T^2 + 16$. We have $F(T) \equiv 0 \pmod{16}$ and the derivative $F'(T) = 2T$ has the exact 2-valuation 1. By the usual approximation process (cf. [Se] 2.2. Theorem 1), we see that f has a square root in K_2 . Hence the ramification locus of $X_L \rightarrow X$ is exactly D , and since D is the sum of horizontal prime divisors, the morphism $U_L \rightarrow U$ is tamely ramified along D .

Now consider the closed subscheme $Y \subset X$ given by the equation $T = 0$, so $Y \cong \text{Spec}(\mathbb{Z})$. Then $D_Y = D \cap Y$ is the point on Y which corresponds to the prime number 2. Let $V = U \cap Y = Y - D_Y$. The base change $V' = U_L \times_U V \rightarrow V$ is the normalization of $V \cong \text{Spec}(\mathbb{Z}[\frac{1}{2}])$ in $\mathbb{Q}(\sqrt{-1})$. But 2 is wildly ramified in $\mathbb{Q}(\sqrt{-1})$, and so $V' \rightarrow V$ is not tamely ramified along D_Y .

Example 2: We construct an example of a cyclic étale covering of smooth varieties which is curve-tame but not numerically tame with respect to some normal but not regular compactification. Furthermore we give an example of a curve-tame Galois covering (with non-nilpotent Galois group) which is numerically tame with respect to some regular compactification but not numerically tame with respect to another regular compactification.

Let $k = \bar{k}$ be an algebraically closed field of characteristic $p > 2$. Let E be an ordinary elliptic curve over k and let $y^2 = x(x-1)(x-\lambda)$ be a Weierstraß equation for E . We obtain a cyclic degree 2 covering $\pi : E \rightarrow \mathbb{P}_k^1$ and we have

$$E \setminus \{\infty\} = \text{Spec}(k[x, y]/y^2 - x(x-1)(x-\lambda)) = \pi^{-1}(\mathbb{A}_k^1 = \text{Spec}(k[x])).$$

Let $E' \rightarrow E$ be the unique cyclic étale covering of degree p . Because of uniqueness, $E' \rightarrow \mathbb{P}_k^1$ is a (tamely ramified) Galois covering of degree $2p$. If $\text{Gal}(E'|\mathbb{P}_k^1)$ would be abelian, we would obtain a cyclic covering of degree p of \mathbb{P}_k^1 which is tame, hence étale because $p = \text{char}(k)$. As \mathbb{P}_k^1 does not have nontrivial étale coverings, the Galois group $\text{Gal}(E'|\mathbb{P}_k^1)$ is nonabelian, hence isomorphic to the dihedral group D_{2p} (which is not nilpotent).

Let $X = \pi^{-1}(\mathbb{A}_k^1 \setminus \{0, 1, \lambda\})$ and let X' be the preimage of X in E' . Taking the product with \mathbb{A}_k^1 , we obtain an étale Galois covering with Galois group D_{2p}

$$\phi : X' \times \mathbb{A}_k^1 \xrightarrow{\text{étale degree } p} X' \times \mathbb{A}_k^1 \xrightarrow{\text{étale degree } 2} (\mathbb{A}_k^1 \setminus \{0, 1, \lambda\}) \times \mathbb{A}_k^1.$$

The covering ϕ is numerically tame with respect to the compactification

$$\bar{\phi} : E' \times \mathbb{P}_k^1 \xrightarrow{\text{étale degree } p} E \times \mathbb{P}_k^1 \xrightarrow{\text{degree } 2} \mathbb{P}_k^1 \times \mathbb{P}_k^1,$$

in particular, ϕ is curve-tame. We will show that ϕ is not numerically tame with respect to another regular compactification of $(\mathbb{A}_k^1 \setminus \{0, 1, \lambda\}) \times \mathbb{A}_k^1$:

Consider the embedding

$$(\mathbb{A}_k^1 \setminus \{0, 1, \lambda\}) \times \mathbb{A}_k^1 \hookrightarrow \mathbb{P}_k^2, (x, t) \mapsto (x : t : 1),$$

and denote the normalizations of \mathbb{P}_k^2 in the function fields of $X \times \mathbb{A}_k^1$ and $X' \times \mathbb{A}_k^1$ by Y and Y' , respectively.

We claim that the curve-tame cyclic covering $X' \times \mathbb{A}_k^1 \rightarrow X \times \mathbb{A}_k^1$ is not numerically tame with respect to the normal compactification $X \times \mathbb{A}_k^1 \hookrightarrow Y$. From this it follows that $\phi : X' \times \mathbb{A}_k^1 \rightarrow (\mathbb{A}_k^1 \setminus \{0, 1, \lambda\}) \times \mathbb{A}_k^1$ is not numerically tame with respect to the regular compactification $(\mathbb{A}_k^1 \setminus \{0, 1, \lambda\}) \times \mathbb{A}_k^1 \hookrightarrow \mathbb{P}_k^2$.

In order to show the claim, it suffices to prove that there is exactly one point in Y' over $P = (0 : 1 : 0) \in \mathbb{P}_k^2$. This can be seen as follows: P corresponds to the maximal ideal $\mathfrak{m} = (t^{-1}x, t^{-1}) \subset k[t^{-1}x, t^{-1}]$. This ring is contained in $k[x, t^{-1}]$ and \mathfrak{m} is the preimage of the principal prime ideal $\mathfrak{p} = (t^{-1}) \subset k[x, t^{-1}]$. It therefore suffices to show that there is exactly one prime ideal above \mathfrak{p} in the integral closure of $k[x, t^{-1}]$ in K' . But this is easily seen: setting $E' \setminus \pi'^{-1}(\infty) = \text{Spec}(A')$, where $\pi' : E' \rightarrow \mathbb{P}_k^1$ is the projection, the integral closure of $k[x, t^{-1}]$ in K' is just $A'[t^{-1}]$.

Example 3: We construct a similar example as in Example 2, but with arithmetic surfaces instead of varieties. This example assumes some familiarity with S. Saito's class field theory for curves over local fields, see [Sa].

Let $p \neq 2$ be a prime number, $k|\mathbb{Q}_p$ a p -adic field, $E \rightarrow k$ an elliptic curve and $E \rightarrow \mathbb{P}_k^1$ the degree 2 covering defined by a Weierstraß model. Then the normalization \mathcal{W} of $\mathbb{P}_{\mathcal{O}_k}^1$ in $k(E)$ is a normal model of E over $\text{Spec}(\mathcal{O}_k)$. Let $\mathcal{E} \rightarrow \mathcal{W}$ be a minimal resolution such that \mathcal{E} is a regular model of E and such that the reduced special fibre $(\mathcal{E}_s)_{\text{red}}$ of \mathcal{E} is a normal crossing divisor. Recall that

the dual graph Γ of $(\mathcal{E}_s)_{red}$ is defined as follows: the vertices correspond to the irreducible components and an edge connecting two vertices corresponds to an intersection point of the two components.

Let $E' \rightarrow E$ be the maximal étale elementary-abelian p -covering of E in which all closed points of E split completely. By [Sa], Proposition 2.2, it extends to an étale covering $\mathcal{E}' \rightarrow \mathcal{E}$, hence $E' \rightarrow E$ is numerically tame with respect to the regular compactification $E \subset \mathcal{E}$. Furthermore, by loc.cit. Proposition 2.3 and Theorem 2.4, the covering $E' \rightarrow E$ is finite and there is a natural isomorphism $G(E'|E) \cong \pi_1^{ab}(\Gamma)/p$.

Now assume that there exists a (singular) rational point P in the special fibre \mathcal{W}_s of \mathcal{W} such that the subgraph Γ_P of Γ given by the components over P is not contractible. Then not every elementary-abelian p -covering of Γ splits over Γ_P , hence $\mathcal{E}' \rightarrow \mathcal{E}$ cannot come by base change from an étale covering of $\mathcal{W}' \rightarrow \mathcal{W}$. We conclude that $E' \rightarrow E$ is not numerically tame with respect to the normal compactification $E \subset \mathcal{W}$. Furthermore, considering the composite map $E' \rightarrow E \rightarrow \mathbb{P}_k^1$, we obtain an example of a curve-tame but not numerically tame covering of some open subscheme $U \subset \mathbb{P}_k^1$. This example can be “lifted” to obtain a similar example over the ring of integers of a number field.

References

- [Ab] Abbes, A. *The Grothendieck-Ogg-Shafarevich formula for arithmetic surfaces*. J. Algebraic Geom. **9** (2000), no. 3, 529–576.
- [Bo] Bourbaki, N. *Commutative algebra*. Elements of Mathematics. Springer-Verlag, Berlin, 1998.
- [CE] Chinburg, T.; Erez, B. *Equivariant Euler-Poincaré characteristics and tameness*. Journées arithmétiques 1991. Société Math. de France, Astérisque. **209** (1992), 179–194.
- [CEPT] Chinburg, T.; Erez, B.; Pappas, G.; Taylor, M. J. *Tame actions of group schemes: integrals and slices*. Duke Math. J. **82** (1996), no. 2, 269–308.
- [EGA4] Grothendieck, A. *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas (EGA 4)*. Inst. Hautes Études Sci. Publ. Math. I. No. 20 1964, II. No. 24 1965, No. 28 1966, IV. No. 32 1967.
- [End] Endler, O. *Valuation theory*. Universitext. Springer-Verlag, New York-Heidelberg, 1972.
- [GM] Grothendieck, A.; Murre, J. P. *The tame fundamental group of a formal neighbourhood of a divisor with normal crossings on a scheme*. Lecture Notes in Math., Vol. 208, Springer-Verlag, Berlin-New York, 1971.
- [KL] Katz, N.; Lang, S. *Finiteness theorems in geometric classfield theory. With an appendix by Kenneth A. Ribet*. Enseign. Math. (2) **27** (1981), no. 3–4, 285–319.
- [Lip] Lipman, J. *Desingularization of two-dimensional schemes*. Ann. Math. **107** (1978), 151–207.
- [Liu] Liu, Q. *Algebraic geometry and arithmetic curves*. Oxford Graduate Texts in Mathematics, 6. Oxford Science Publications. Oxford University Press, Oxford, 2002.

- [Lue] Lütkebohmert, W. *On compactification of schemes*. *Manuscripta Math.* **80** (1993), 95–111.
- [NSW] Neukirch, J.; Schmidt, A.; Wingberg, K. *Cohomology of number fields*. 2nd ed. *Grundlehren der Math. Wiss.*, Bd. 323. Springer-Verlag, Berlin, 2008.
- [Sa] Saito, S. *Class field theory for curves over local fields*. *J. Number Theory* **21** (1985), no. 1, 44–80.
- [Sc1] Schmidt, A. *Tame coverings of arithmetic schemes*. *Math. Ann.* **322** (2002), no. 1, 1–18.
- [Sc2] Schmidt, A. *Singular homology of arithmetic schemes*. *Algebra & Number Theory* **1** (2007), no. 2, 183–222.
- [Se] Serre, J.-P. *Local fields*. *Graduate Texts in Math.*, Vol. 67, Springer-Verlag, New York-Berlin, 1979.
- [SGA1] Grothendieck, A. *Revêtements étales et groupe fondamental (SGA 1)*. *Lecture Notes in Math.*, 224, Springer-Verlag, Berlin-New York, 1971.
- [SGA2] Grothendieck, A. *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*. *Advanced Studies in Pure Math.*, Vol. 2. North-Holland Publishing Co., Amsterdam; Masson & Cie, Éditeur, Paris, 1968.
- [SGA4] Artin, M.; Grothendieck, A. and et Verdier, J. L. *Théorie des topos et cohomologie étale des schémas (SGA 4)*. *Lecture Notes in Math.*, 269, 270 and 305, 1972/3.
- [Sha] Shafarevich, I. G. *Lectures on minimal models and birational transformations of two dimensional schemes*. *Tata Inst. Fundam. Res.* 37, Bombay 1966.
- [Va] Vaquié, M. *Valuations. Resolution of singularities (Obergrugl, 1997)*, 539–590, *Progr. Math.*, 181, Birkhäuser, Basel, 2000.
- [Wi] Wiesend, G. *Tamely ramified covers of varieties and arithmetic schemes*. *Forum Math.* **20** (2008), no. 3, 515–522.
- [ZS] Zariski, O.; Samuel, P. *Commutative algebra. Vol. II*. *Graduate Texts in Mathematics*, Vol. 29. Springer-Verlag, New York-Heidelberg, 1975.

NWF I-Mathematik, Universität Regensburg, D-93040 Regensburg, Deutschland. *E-mail address*: moritz.kerz@mathematik.uni-regensburg.de

NWF I-Mathematik, Universität Regensburg, D-93040 Regensburg, Deutschland. *E-mail address*: alexander.schmidt@mathematik.uni-regensburg.de