

WACH MODULES, REGULATOR MAPS AND ε -ISOMORPHISMS IN FAMILIES

REBECCA BELLOVIN AND OTMAR VENJAKOB

ABSTRACT. In this paper, we prove the “local ε -isomorphism” conjecture of Fukaya and Kato [FK06] for (crystalline) families of $G_{\mathbf{Q}_p}$ -representations. This can be regarded as a local analogue of the Iwasawa main conjecture for families, extending earlier work of Kato for rank one modules (see [Ven13]), of Benois and Berger for crystalline $G_{\mathbf{Q}_p}$ -representations with respect to the cyclotomic extension (see [BB08]) as well as of Loeffler, Venjakob and Zerbes (see [LVZ15]) for crystalline $G_{\mathbf{Q}_p}$ -representations with respect to abelian p -adic Lie extensions of \mathbf{Q}_p . Nakamura [Nak13, Nak14] has shown Kato’s ε -conjecture for rank one (φ, Γ) -modules over the Robba ring, which means in particular only after inverting p ; moreover he attaches an ε -isomorphism to trianguline (φ, Γ) -modules over the Robba ring depending on a fixed triangulation. As a consequence of our result one obtains the independence of the chosen triangulation for crystalline families a posteriori.

The main ingredient of (the integrality part of) our proof consists of the construction of families of Wach modules generalizing work of Wach and Berger [Ber04] and following Kisin’s approach via a corresponding moduli space [Kis09].

CONTENTS

1. Introduction	2
Acknowledgements	3
Notation and Conventions	3
2. Rings of p -adic Hodge theory	4
3. Families of Wach modules	6
4. Generic fibre	16
4.1. Families of Wach modules and (φ, Γ) -modules	16
4.2. Comparisons between $\mathbf{N}(M)$ and $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$	17
5. Galois cohomology	21
6. Regulator maps for families	24
6.1. Γ -regulator	24
6.2. G -regulator	25
7. Construction of the isomorphism	26
7.1. Construction of Θ over $\mathcal{K}_{\tilde{R}}(G)$	28
7.2. Definition of Θ over $\Lambda_R(G)[1/p]$	29
7.3. Definition of the epsilon-isomorphism	31
7.4. Properties of the epsilon-isomorphism	32
7.5. Epsilon-isomorphisms for more general modules	34
8. Application to deformation rings	37
9. Relation to Nakamura’s work	38
Appendix A. Generic fibers	38
A.1. Quasi-Stein spaces	38
A.2. Partial generic fibers	40
Appendix B. Determinant functors	41
References	43

2010 *Mathematics Subject Classification*. Primary: 11R23. Secondary: 11S40.

The first author was partially supported by an NSF Mathematical Sciences Postdoctoral Research Fellowship, and the second author was supported by the DFG-Forschergruppe ”Symmetrie, Geometrie und Arithmetik” .

1. INTRODUCTION

The significance of (local) ε -factors à la Deligne and Tate, or more generally of the (conjectural) ε -isomorphism suggested by Fukaya and Kato in [FK06, §3], is at least twofold: First, they are important ingredients to obtain a precise functional equation for L -functions or more generally for (conjectural) ζ -isomorphism (loc. cit., §2) of motives in the context of Tamagawa number conjectures; second, they are essential in interpolation formulae of (actual) p -adic L -functions and for the relation between ζ -isomorphisms and (conjectural) p -adic L -functions as discussed in (loc. cit., §4). Of course the two occurrences are closely related; for a survey on these ideas see also [Ven07].

The ε -isomorphism conjecture asserts roughly speaking the existence of a canonical trivialization of the determinant of a cohomology complex associated to any representation M of the Galois group $G_{\mathbf{Q}_p} = \text{Gal}(\overline{\mathbf{Q}_p}/\mathbf{Q}_p)$ with coefficients in a p -adically complete local ring R . This trivialization is required to be compatible with a specific “standard” trivialization of the corresponding complex for de Rham representations with coefficients in finite extensions of \mathbf{Q}_p obtained by specializing M at ideals, or more generally at representations, of R . This standard trivialization contains information about the ε -factor of the corresponding Weil–Deligne representation, hence the terminology “ ε -isomorphism”, and the Bloch–Kato-exponential map.

The aim of this article is to prove, for any crystalline family M of Galois representations of $G_{\mathbf{Q}_p} = \text{Gal}(\overline{\mathbf{Q}_p}/\mathbf{Q}_p)$ over a complete local noetherian \mathbf{Z}_p -algebra R with finite residue field, the existence of the ε -isomorphism (see Theorem 7.13)

$$\varepsilon_{\Lambda_R(G),\xi}(M) : \text{Det}_{\Lambda_{\tilde{R}}(G)}(0) \xrightarrow{\sim} \left[\text{Det}_{\Lambda_{\tilde{R}}(G)}(\Lambda_{\tilde{R}}(G) \otimes_{\Lambda_R(G)} R\Gamma_{\text{Iw}}(K_\infty, M)) \right] \left[\text{Det}_{\Lambda_{\tilde{R}}(G)}(\Lambda_{\tilde{R}}(G) \otimes_R M) \right]$$

such that the specializations $\varepsilon_{\Lambda_R(G),\xi}(M) \otimes_R \mathcal{O}_L$ with respect to any \mathbf{Z}_p -algebra homomorphism $R \rightarrow \mathcal{O}_L$ into the ring of integers \mathcal{O}_L of a finite extension L of \mathbf{Q}_p agree with the ε -isomorphisms $\varepsilon_{\Lambda_{\mathcal{O}_L}(G),\xi}(M \otimes_R \mathcal{O}_L)$ established in [LVZ15]. In particular $\varepsilon_{\Lambda_R(G),\xi}(M)$ satisfies the above mentioned compatibility with the standard trivializations for corresponding specializations (see Proposition 7.21). For further properties we refer the reader to section 7.4.

Here $\xi = (\xi_n)_{n \geq 1}$ denotes a compatible system of p -power roots of unity generating $\mathbf{Z}_p(1)$ as a \mathbf{Z}_p -module, $\Lambda_R(G)$ is the Iwasawa algebra of $G = G(K_\infty/\mathbf{Q}_p)$ for $K_\infty = K(\mu(p))$ with K any (possibly infinite) unramified extension K of \mathbf{Q}_p and $R\Gamma_{\text{Iw}}(K_\infty, M)$ denotes the complex calculating local Iwasawa cohomology of M over K_∞ . Furthermore, for an associative ring A with one, Det_A denotes the determinant functor à la Knudsen–Mumford (if A is commutative) or à la Deligne/Fukaya–Kato (even for non-commutative A) (see Appendix B); from a technical point of view we want to stress that we use the realisation via line bundles whenever it is possible in order to take advantage of (stronger) arguments and higher flexibility from commutative algebra. The Iwasawa algebra $\Lambda_{\tilde{R}}(G)$ is defined in (11). We are mainly interested in the case, where $G \cong \mathbf{Z}_p^2 \times \Delta$ for a finite group Δ . This specific instance of the local ε -isomorphism conjecture can be thought of as a “local Iwasawa main conjecture” for M over the extension $\mathbf{Q}_p^{\text{ab}}/\mathbf{Q}_p$; compare the discussion at the end of §2 in [Ven13].

The key ingredient in the construction of the local ε -isomorphism and the technical heart of this paper is the construction of Wach modules $\mathbf{N}(M)$ for formal families of Galois representations M . The theory of (φ, Γ) -modules and several other aspects of p -adic Hodge theory had been generalized to formal and rigid-analytic families by work of Dee [Dee01], Berger and Colmez [BC08], Kedlaya and Liu [KL10], and Bellovin [Bel15]. In §3, we give a theory of Wach modules for formal families of crystalline representations of G_F , following Kisin’s strategy for the construction of potentially semi-stable deformation rings [Kis08].

We then compare Iwasawa cohomology to our Wach modules, extending the classical case. With this in hand, we can define ε -isomorphisms for families of crystalline representations of $G_{\mathbf{Q}_p}$ with coefficients in rings R which are complete, local, noetherian \mathbf{Z}_p -algebras which are Cohen–Macaulay, normal, and flat over \mathbf{Z}_p , with reduced generic fiber, and we show that the ε -isomorphisms we construct are compatible with base change and with the ε -isomorphisms constructed in [LVZ15].

With this machinery at hand we can sharpen some work of Nakamura [Nak13], who had given a different construction of ε -isomorphism even for not necessarily étale, but trianguline (φ, Γ) -modules (in families) over the Robba ring, but in which the final integrality statement is missing and which a priori depends on the choice of a triangulation. For crystalline families our results implies the independence of such a choice (Corollary 9.1). See section 9 for a comparison with his result.

The advantage of providing ε -isomorphisms for families is as follows: First of all, Iwasawa main conjectures [SU14] or (Equivariant) Tamagawa Number Conjectures are nowadays often considered over (e.g. Hida-) families. Second, we hope that our work can be used to obtain results towards the construction of ε -isomorphisms attached to torsion representations: In previous work [LVZ15] by the second author together with David Loeffler and Sarah Zerbes we constructed an “ ε -isomorphism” for crystalline p -adic Galois representations. Unfortunately, we were not able to address the following general question: let T, T' be two torsion-free \mathbf{Z}_p -representations of $G_{\mathbf{Q}_p}$ such that $T \cong T'$ modulo p^k for some k . Does it follow that the ε -isomorphisms for T and T' agree modulo p^k ? In this paper, we give a partial answer and show that if T and T' both are crystalline with Hodge-Tate weights in $[a, b]$, then the desired congruence holds, if the framed deformation ring $R_{\text{cris}}^{\square, [a, b]}$ is normal and Cohen-Macaulay. We do this by showing that the ε -isomorphisms for T and T' are both specializations of a “universal” isomorphism over $R_{\text{cris}}^{\square, [a, b]}$; see section 8.

We expect that it is possible to extend our constructions to the setting of affinoid families of Galois representations, using techniques similar to those of [Wan15]. However, we have not done so here.

We quickly outline the content of the various sections: After recalling the basic rings of p -adic Hodge theory in section 2, we construct Wach modules $\mathbf{N}(M)$ for formal families of Galois representations M in section 3; the main result is summarized in Theorem 3.31. In section 4 we relate (the base change $\mathbf{N}(M)^{\text{rig}}$ of) the Wach module $\mathbf{N}(M)$ to the (φ, Γ) -module $\mathbf{D}_{\text{rig}}^{\dagger}(M^{\text{rig}})$ attached to the associated family M^{rig} of Galois representations over the rigid analytic generic fiber R^{rig} of R . In particular, we discuss the relation with $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$, generalizing parts of [Ber04] to families. This permits us to define $\mathbf{D}_{\text{cris}}(M)$ *without* inverting p . In section 5 we explain how to express Iwasawa cohomology of M in terms of $\mathbf{N}(M)$, generalizing to families a variant of Fontaine’s isomorphism due to Loeffler and Zerbes in [LZ14]; see Propositions 5.1 and 5.6. Again following [LZ14], we construct the regulator map attached to M in section 6 and we study its specializations in Theorem 6.1. Finally, in section 7 we put everything together in order to construct an isomorphism Θ — roughly speaking the determinant of the regulator map — first over the total ring of fractions and then over the Iwasawa algebra with p inverted. The isomorphism Θ relates the Iwasawa cohomology of M to $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$. Using Theorem 7.10 which relates the determinants of $\mathbf{D}_{\text{cris}}(M)[1/p]$ and $M[1/p]$, we are able to define the ε -isomorphism (see Definition 7.12) over the Iwasawa algebra with p inverted. By a specialization argument we show its integrality in Theorem 7.13. Finally, we briefly discuss the application to deformation rings in section 8 before relating our ε -isomorphism to that of Nakamura in section 9.

ACKNOWLEDGEMENTS

This project grew out of natural questions arising from the previous work [LVZ15], and we are very grateful to David Loeffler and Sarah Zerbes for many suggestions. We would also like to thank Toby Gee, Brandon Levin, James Newton, and Carl Wang Erickson for helpful discussions.

NOTATION AND CONVENTIONS

Let p be an odd prime. For H a p -adic analytic group, and L a complete discretely valued extension of \mathbf{Q}_p with ring of integers \mathcal{O} and uniformiser ϖ , we write $\Lambda_{\mathcal{O}}(H)$ and $\Lambda_L(H) = L \otimes_{\mathcal{O}} \Lambda_{\mathcal{O}}(H)$ for the Iwasawa algebras of H with \mathcal{O} and L coefficients, and $\mathcal{H}_L(H)$ for the algebra of L -valued locally analytic distributions on H , which is the completion of $\Lambda_L(H)$ in a certain Fréchet topology. We shall only use these constructions in cases where H is abelian and p -torsion-free, in which case all of these algebras are reduced commutative semi-local rings, and can be interpreted as algebras of functions on the p -adic analytic space parameterising characters of H .

We shall also need the notation $\mathcal{K}_L(H)$, signifying the total ring of quotients of $\mathcal{H}_L(H)$, which is a finite direct product of fields.

Let $\mathbf{Q}_{p, \infty} = \mathbf{Q}_p(\mu_{p^\infty})$ and $\Gamma = \text{Gal}(\mathbf{Q}_{p, \infty}/\mathbf{Q}_p)$, and let $\chi_{\text{cyc}} : G_{\mathbf{Q}_p} \twoheadrightarrow \Gamma \rightarrow \mathbf{Z}_p^\times$ be the p -cyclotomic character. We write γ_{-1} for the unique element of Γ such that $\chi(\gamma_{-1}) = -1$.

If M denotes any $G_{\mathbf{Q}_p}$ -module over \mathbf{Z}_p and i is any integer, we write $M(i)$ for the i th Tate twist of M .

By R we shall denote a complete local noetherian \mathbf{Z}_p -algebra with maximal ideal \mathfrak{m}_R . We write $H_{\mathbf{Q}_p}$ for the kernel of χ_{cyc} . If A is any associative ring, M a (left) A -module and Y a (right) A -module, we denote the base change $Y \otimes_A M$ often just by M_Y . If φ denotes an injective ring homomorphism of A ,

which is now assumed to be commutative, we shall as usual set $\varphi^*M := A \otimes_{A,\varphi} M$ where the left copy of A is viewed as A -module via φ . Assume, moreover, that M itself bears a φ -linear action φ_M . Then we obtain a canonical map $\varphi^*M \rightarrow M$ sending $a \otimes m$ to $a\varphi_M(m)$. In case the latter map is injective we shall sometimes identify φ^*M with its image in M without further indication.

We write $\widehat{\mathbf{Q}}_p^{\text{nr}}$ for the completion of the maximal unramified extension \mathbf{Q}_p^{nr} of \mathbf{Q}_p while the ring of integers are denoted by $\widehat{\mathbf{Z}}_p^{\text{nr}}$ and \mathbf{Z}_p^{nr} , respectively. F shall usually denote a finite *unramified* extension of \mathbf{Q}_p .

2. RINGS OF p -ADIC HODGE THEORY

The rings (and their properties) we will use are primarily defined in [Ber02].

Let F be a finite unramified extension of \mathbf{Q}_p with ring of integers \mathcal{O}_F and residue field k_F . Let $\widetilde{\mathbf{E}} := \varprojlim_{x \rightarrow x^p} \mathbf{C}_p$, and let $\widetilde{\mathbf{E}}^+$ be the subset of $x \in \widetilde{\mathbf{E}}$ such that $x^{(0)} \in \mathcal{O}_{\mathbf{C}_p}$. Then $\widetilde{\mathbf{E}}$ is an algebraically closed field of characteristic p . There is a valuation $\text{ord}_{\mathbf{E}}$ defined by $v_{\mathbf{E}}(x^{(i)}) = v_p(x^{(0)})$, and a Frobenius (given by raising to the p th power). The valuation ring of $\widetilde{\mathbf{E}}$ with respect to this valuation is $\widetilde{\mathbf{E}}^+$, and $\widetilde{\mathbf{E}}$ is complete with respect to $v_{\mathbf{E}}$.

Let $\varepsilon := (\varepsilon^{(0)}, \varepsilon^{(1)}, \varepsilon^{(2)} \dots) \in \widetilde{\mathbf{E}}^+$ be a choice of compatible p th power roots of unity with $\varepsilon^{(0)} = 1$ and $\varepsilon^{(1)} \neq 1$. There is a natural map $k_F((\bar{\pi})) \rightarrow \widetilde{\mathbf{E}}$ given by sending $\bar{\pi}$ to $\varepsilon - 1$; we denote its image by \mathbf{E}_F , we denote by \mathbf{E} the separable closure of \mathbf{E}_F inside $\widetilde{\mathbf{E}}$, and we denote by \mathbf{E}^+ the valuation ring of \mathbf{E} .

Let $\widetilde{\mathbf{A}}^+ := W(\widetilde{\mathbf{E}}^+)$ and $\widetilde{\mathbf{A}} := W(\widetilde{\mathbf{E}})$. There are two possible topologies on $\widetilde{\mathbf{A}}^+$ and $\widetilde{\mathbf{A}}$, the p -adic topology or the weak topology; they are complete for both.

The *p -adic topology* is defined by putting the discrete topology on $W(\widetilde{\mathbf{E}})/p^n W(\widetilde{\mathbf{E}})$ for all n , and taking the projective limit topology on $\widetilde{\mathbf{A}}$; $\widetilde{\mathbf{A}}^+$ is given the subspace topology. The *weak topology* is defined by putting the valuation topology on $\widetilde{\mathbf{E}}$ and giving $\widetilde{\mathbf{A}}$ the product topology; $\widetilde{\mathbf{A}}^+$ is again given the subspace topology.

Alternatively, the weak topology on $\widetilde{\mathbf{A}}$ is given by taking the sets

$$U_{k,n} := p^k \widetilde{\mathbf{A}} + [\tilde{p}]^n \widetilde{\mathbf{A}}^+ \text{ for } k, n \geq 0$$

to be a basis of neighborhoods around 0, where $\tilde{p} \in \widetilde{\mathbf{E}}^+$ is any fixed element with $\tilde{p}^{(0)} = p$ (i.e., \tilde{p} is a system of compatible p -power roots of p). The weak topology on $\widetilde{\mathbf{A}}^+$ is similarly generated by the sets $U_{k,n} \cap \widetilde{\mathbf{A}}^+ = p^k \widetilde{\mathbf{A}}^+ + [\tilde{p}]^n \widetilde{\mathbf{A}}^+$. The weak topology can also be defined by a family of semi-norms, but we will not use this.

Both rings carry continuous actions of Frobenius (for either topology). However, the Galois action is continuous for the weak topology, but it is not continuous for the p -adic topology because the Galois actions on $\widetilde{\mathbf{E}}^+$ and $\widetilde{\mathbf{E}}$ are not discrete.

There are versions of all of these rings with no tilde; they are imperfect versions of the rings with tildes.

Let $\pi \in \widetilde{\mathbf{A}}$ denote $[\varepsilon] - 1$, where $[\varepsilon]$ denotes the Teichmüller lift of ε . Then there is a well-defined injective map $\mathcal{O}_F[[X]][X^{-1}] \rightarrow \widetilde{\mathbf{A}}$ given by sending X to π ; we let \mathbf{A}_F denote the p -adic completion of the image. This is a Cohen ring for \mathbf{E}_F .

Let $\mathbf{B}_F := \mathbf{A}_F[1/p]$. Then \mathbf{B}_F is a field, and we let \mathbf{B} be the completion of the maximal unramified extension of \mathbf{B}_F inside $\widetilde{\mathbf{B}}$. Further, we put $\mathbf{A} := \mathbf{B} \cap \widetilde{\mathbf{A}}$ and $\mathbf{A}^+ := \mathbf{A} \cap \widetilde{\mathbf{A}}^+$. Alternatively, \mathbf{A} is the completion of the integral closure of the image of $\mathbf{Z}_p[[X]][X^{-1}]$ in $\widetilde{\mathbf{A}}$.

Because extensions of \mathbf{E}_F correspond to unramified extensions of \mathbf{B}_F , we get a natural Galois action on \mathbf{A} and \mathbf{A}^+ . For any extension K/\mathbf{Q}_p , we may therefore define $\mathbf{A}_K := \mathbf{A}^{H_K}$ and $\mathbf{B}_K := \mathbf{A}_K[1/p]$. When K is unramified over \mathbf{Q}_p , this agrees with our original definition of these rings.

In fact, we can describe \mathbf{A}_K more explicitly: If $F' \subset K$ is the maximal unramified subfield of K , then every element of \mathbf{A}_K can be written uniquely in the form $\sum_{k \in \mathbf{Z}} a_k \pi_K^k$, where $\{a_k\}_{k \in \mathbf{Z}} \in \mathcal{O}_{F'}$ is a sequence tending to 0 as k tends to $-\infty$ and π_K is a (non-canonical) lift of a uniformizer of \mathbf{E}_K [Col08, Proposition 7.1]. The subring $\{\sum_{k \geq 0} a_k \pi_K^k\}$ is contained in \mathbf{A}_K^+ ; on the other hand, if $f = \sum_{k \in \mathbf{Z}} a_k \pi_K^k \in \mathbf{A}_K^+$, then $f \in \widetilde{\mathbf{A}}^+$ and therefore $a_k \equiv 0 \pmod{p}$ for $k < 0$ (since $\pi_K^{-1} \in \mathbf{E}_K \setminus \mathbf{E}_K^+$).

We also get an action of Frobenius induced from that on $\widetilde{\mathbf{A}}$; however, because \mathbf{E} is imperfect, the action of Frobenius is no longer surjective.

If R is a complete local noetherian \mathbf{Z}_p -algebra with maximal ideal \mathfrak{m}_R and finite residue field, we define $R \widehat{\otimes} \mathbf{A}$, $R \widehat{\otimes} \mathbf{A}^+$, $R \widehat{\otimes} \mathbf{A}_F$, and $R \widehat{\otimes} \mathbf{A}_F^+$ to be the tensor products over \mathbf{Z}_p , completed with respect to \mathfrak{m}_R . If R is discrete, these completed tensor products become ordinary tensor products over \mathbf{Z}_p .

This permits us to define $\mathbf{D}(M)$ and $\mathbf{D}^+(M)$ for R -linear representations of $G_{\mathbf{Q}_p}$ as follows:

$$\mathbf{D}(M) = ((R \widehat{\otimes} \mathbf{A}) \otimes_R M)^{H_{\mathbf{Q}_p}} \quad \text{and} \quad \mathbf{D}^+(M) = ((R \widehat{\otimes} \mathbf{A}^+) \otimes_R M)^{H_{\mathbf{Q}_p}}.$$

Note that there are canonical isomorphisms

$$(1) \quad \mathbf{D}(M) \cong \varprojlim_n \mathbf{D}(M/\mathfrak{m}_R^n M) \quad \text{and} \quad \mathbf{D}^+(M) \cong \varprojlim_n \mathbf{D}^+(M/\mathfrak{m}_R^n M).$$

A consideration of Witt coordinates shows that the cokernel C of the injection $\widetilde{\mathbf{A}}^+ \hookrightarrow \widetilde{\mathbf{A}}$ has no p -torsion, and is therefore \mathbf{Z}_p -flat. It follows that the cokernel of $\mathbf{A}^+ \hookrightarrow \mathbf{A}$ is also \mathbf{Z}_p -flat. As a consequence, the natural maps $R \widehat{\otimes} \widetilde{\mathbf{A}}^+ \rightarrow R \widehat{\otimes} \widetilde{\mathbf{A}}$ and $R \widehat{\otimes} \mathbf{A}^+ \rightarrow R \widehat{\otimes} \mathbf{A}$ are injective.

We record the following useful lemma:

Lemma 2.1. *Let R be a discrete \mathbf{Z}_p -algebra, and let $I \subset R$ be an ideal. Then $I(R \otimes_{\mathbf{Z}_p} \mathbf{A}) \cap (R \otimes_{\mathbf{Z}_p} \mathbf{A}^+) = I(R \otimes_{\mathbf{Z}_p} \mathbf{A}^+)$.*

Proof. Suppose that $f := r_1 f_1 + \dots + r_k f_k \in R \otimes_{\mathbf{Z}_p} \mathbf{A}^+$, where $r_i \in I$ and $f_i \in \mathbf{A}$, and let \bar{f}_i denote the image of f_i in the cokernel $R \otimes_{\mathbf{Z}_p} C$ of the map $R \otimes_{\mathbf{Z}_p} \mathbf{A}^+ \rightarrow R \otimes_{\mathbf{Z}_p} \mathbf{A}$. By assumption, $\sum_i r_i \bar{f}_i = 0$, so there exist $\bar{f}'_j \in R \otimes_{\mathbf{Z}_p} C$ and $r'_{ij} \in R$ such that $\bar{f}_i = \sum_j r'_{ij} \bar{f}'_j$ for all i and $\sum_i r_i r'_{ij} = 0$ by [Mat89, Theorem 7.6] (since $R \otimes_{\mathbf{Z}_p} C$ is R -flat). We may choose $\tilde{f}'_j \in \mathbf{A}$ lifting \bar{f}'_j ; setting $f'_i := \sum_j r'_{ij} \tilde{f}'_j$, we have $f_i - f'_i \in \mathbf{A}^+$ for all i and $\sum_i r_i (f_i - f'_i) = \sum_i r_i f_i = f$ so we are done. \square

Remark 2.2. This does not imply that if $f \in \mathbf{A}$ satisfies $rf \in \mathbf{A}^+$ then $f \in \mathbf{A}^+$, merely that there is some $f' \in \mathbf{A}^+$ such that $rf = rf'$.

Furthermore, \mathbf{A}/p is the separable closure of $\mathbf{A}_F/p = k((\pi))$, where k is the residue field of F , and \mathbf{A}^+/p is its ring of integers [Fon90, §1.8(a)].

Lemma 2.3. *Let R be a discrete \mathbf{Z}/p^n -algebra.*

- (1) *For any finite extension K/F , the natural map $R \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+ \rightarrow (R \otimes_{\mathbf{Z}_p} \mathbf{A}^+)^{H_K}$ is an isomorphism.*
- (2) *If $I \subset R$ is an ideal, then $(IR \otimes_{\mathbf{Z}_p} \mathbf{A}^+)^{H_K} = IR \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+.$*
- (3) *Suppose in addition that R is a finite-type \mathbf{Z}/p^n -algebra. Then every element of $R \otimes_{\mathbf{Z}_p} \mathbf{A}^+$ is contained in $R \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+$ for some finite extension K/F .*

Proof. (1) We proceed by induction on n . If $n = 1$, the result follows by choosing a basis for R over \mathbf{F}_p and using the fact that \mathbf{A}^+/p is the ring of integers of a separable closure of \mathbf{A}_F/p . So we may assume the result for $n - 1$ and consider the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & p^{n-1}R \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+ & \longrightarrow & R \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+ & \longrightarrow & (R \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+)/p^{n-1} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & (p^{n-1}R \otimes_{\mathbf{Z}_p} \mathbf{A}^+)^{H_K} & \longrightarrow & (R \otimes_{\mathbf{Z}_p} \mathbf{A}^+)^{H_K} & \longrightarrow & ((R \otimes_{\mathbf{Z}_p} \mathbf{A}^+)/p^{n-1})^{H_K} \end{array}$$

Let $f \in (R \otimes_{\mathbf{Z}_p} \mathbf{A}^+)^{H_K}$ and let \bar{f} be its image in $((R \otimes_{\mathbf{Z}_p} \mathbf{A}^+)/p^{n-1})^{H_K}$. The right-most arrow is an isomorphism, by the inductive hypothesis, so \bar{f} lifts to $\tilde{f} \in R \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+$. Then $f - \tilde{f}$ is in the image of $(p^{n-1}R \otimes_{\mathbf{Z}_p} \mathbf{A}^+)^{H_K}$. But $p^{n-1}R$ is a finite R/p -module, so it is an \mathbf{F}_p -vector space. Choosing a basis for $p^{n-1}R$ and considering the structure of \mathbf{A}^+/p again implies that $f - \tilde{f}$ lifts to $p^{n-1}R \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+$, and therefore to $R \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+$.

Injectivity follows by another diagram chase.

- (2) There is a natural commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & IR \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+ & \longrightarrow & R \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+ & \longrightarrow & (R/I) \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+ \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & (IR \otimes_{\mathbf{Z}_p} \mathbf{A}^+)^{H_K} & \longrightarrow & (R \otimes_{\mathbf{Z}_p} \mathbf{A}^+)^{H_K} & \longrightarrow & ((R/I) \otimes_{\mathbf{Z}_p} \mathbf{A}^+)^{H_K} \end{array}$$

Then the first part of this lemma and the snake lemma imply the desired result.

- (3) We proceed by induction on n . If $n = 1$, then $R \otimes_{\mathbf{Z}_p} \mathbf{A}^+ = \varinjlim_K R \otimes_{\mathbf{F}_p} (\mathbf{A}^+/p)^{H_K}$. Since R is assumed finite-type over \mathbf{F}_p , the result follows. We now assume the result holds for $n - 1$. Consider $f \in R \otimes_{\mathbf{Z}_p} \mathbf{A}^+$, and let \bar{f} denote its image in $(R/p^{n-1}) \otimes_{\mathbf{Z}_p} \mathbf{A}^+$. By the inductive hypothesis, $\bar{f} \in (R/p^{n-1}) \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+$ for some finite extension K/F , and \bar{f} lifts to $\tilde{f} \in R \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+$. Then $f - \tilde{f}$ lies in the finite $(R/p) \otimes_{\mathbf{Z}_p} \mathbf{A}^+$ -module $p^{n-1} R \otimes_{\mathbf{Z}_p} \mathbf{A}^+$, so we see that $f - \tilde{f} \in R \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+$ for some finite extension K'/F and we are done. \square

3. FAMILIES OF WACH MODULES

families

Let R be a complete local noetherian \mathbf{Z}_p -algebra with finite residue field. We give R its natural profinite topology.

Definition 3.1. Let K/\mathbf{Q}_p be a finite extension. A *family of G_K -representations* over R is a finite free R -module M endowed with a continuous R -linear action of G_K . If X is a rigid analytic space over \mathbf{Q}_p , a *family of G_K -representations* over X is a finite projective \mathcal{O}_X -module endowed with a continuous \mathcal{O}_X -linear action of G_K .

Note that to a family M of G_K -representations over R we can naturally assign a family M^{rig} of G_K -representations over $X = \text{Spf}(R)^{\text{rig}}$; see Appendix A.1 for details.

For any rigid analytic space X over \mathbf{Q}_p , we further define the sheaves of rings $\mathcal{B}_{X,\max}^+$ and $\mathcal{B}_{X,\max}$ by setting $\mathcal{B}_{X,\max}^+(U) := \mathcal{O}_X(U) \widehat{\otimes}_{\mathbf{Q}_p} \mathbf{B}_{\max}^+$ and $\mathcal{B}_{X,\max}(U) := \cup_{i \in \mathbf{N}} t^{-i} \mathcal{B}_{X,\max}^+(U)$. If M is a family of $G_{\mathbf{Q}_p}$ -representations over X , we define a sheaf of abelian groups $\mathcal{D}_{\text{cris}}(M)$ by setting $\mathcal{D}_{\text{cris}}(M)(U) := (M(U) \widehat{\otimes}_{\mathcal{O}_X(U)} \mathcal{B}_{X,\max}(U))^{G_K}$; we let $\mathbf{D}_{\text{cris}}(M)$ denote $\mathcal{D}_{\text{cris}}(M)(X)$. We say that M is *crystalline* if $\mathcal{D}_{\text{cris}}(M)$ is a vector bundle over \mathcal{O}_X of the same rank as M and the natural map $\mathcal{B}_{X,\max} \otimes_{\mathcal{O}_X} \mathcal{D}_{\text{cris}}(M) \rightarrow \mathcal{B}_{X,\max} \otimes_{\mathcal{O}_X} M$ is an isomorphism.

We now fix an interval $[a, b]$ of integers.

prop:crisfam

Proposition 3.2. *Suppose that $R[1/p]$ is reduced, and let M be a family of G_K -representations over R . Then the following are equivalent:*

- (1) *For every homomorphism $R \rightarrow E'$, where E'/E is a finite extension, the representation $E' \otimes_R M$ is crystalline with Hodge–Tate weights in $[a, b]$.*
- (2) *For every homomorphism $R \rightarrow S$, where S is a finite flat \mathbf{Z}_p -algebra, the representation $S \otimes_R M[1/p]$ is crystalline with Hodge–Tate weights in $[a, b]$.*

(The same holds if “crystalline” is replaced by “semistable”, although we shall not need this.)

Proof. The implication (2) \Rightarrow (1) is clear. To show that (1) \Rightarrow (2) we will use the results of [Bel15]. Suppose we are given a homomorphism $R \rightarrow S$ as in (2). Let R' be an affinoid algebra such that there is a natural flat homomorphism $R \rightarrow R'$ with dense image. By Theorem 1.1.3 of *op.cit* we know that there is a unique closed rigid-analytic subvariety of $X' = \text{Sp}(R')$ such that a homomorphism from R' to an artinian E -algebra factors through this quotient if and only if the corresponding specialization of M is crystalline with Hodge–Tate weights in $[a, b]$. By assumption, this subvariety contains every point of X' . But X' is reduced, so this subvariety is the whole of X' . Since the generic fiber $\text{Spf}(R)^{\text{rig}}$ can be exhausted by affinoid subspaces as above, we see that (1) \Rightarrow (2) as required. \square

We say M is *crystalline with Hodge–Tate weights in $[a, b]$* if R is flat over \mathbf{Z}_p and the above equivalent conditions hold. This is equivalent to the family of representations M^{rig} over the quasi-Stein space $\text{Spf}(R)^{\text{rig}}$ being crystalline with Hodge–Tate weights in $[a, b]$. If R is \mathbf{Z}_p -finite and flat and $K = F$ is unramified over \mathbf{Q}_p , there is an associated *Wach module* $\mathbf{N}(M)$. We wish to extend this to the general case.

The problem is that we would like to have a good notion of Wach modules for families of crystalline representations with coefficients in power series rings. But because the (integral) Wach module functor can fail to be exact, naive constructions do not seem to have good properties. For example, if R is a finite flat \mathbf{Z}_p -algebra and M is an R -linear lattice in a crystalline representation of $G_{\mathbf{Q}_p}$, there is an associated Wach module $\mathbf{N}(M)$; it is a finite $R \otimes_{\mathbf{Z}_p} \mathbf{A}_{\mathbf{Q}_p}^+$ -module, but it is in general not projective over

$R \otimes_{\mathbf{Z}_p} \mathbf{A}_{\mathbf{Q}_p}^+$. This makes it difficult to verify that $\mathbf{N}(M)$ has good base change properties, or to show that $\mathbf{N}(M)$ has good finiteness properties when R is a power series ring.

We give a definition of Wach modules. As we are primarily interested in positive representations (that is, Galois representations with non-positive Hodge–Tate weights, where the cyclotomic character has weight 1), we restrict to that case.

def:Wach

Definition 3.3. A *Wach module* with weights in $[-h, 0]$ and coefficients in a complete local noetherian \mathbf{Z}_p -algebra R with finite residue field is a finite $R \widehat{\otimes}_{\mathbf{Z}_p} \mathbf{A}_F^+$ -module N equipped with an R -linear action of Γ such that Γ acts trivially on $N/\pi N$, and equipped with an R -linear Frobenius $\varphi : N \rightarrow N$ commuting with Γ such that the cokernel $N/\varphi^*(N)$ is killed by q^h , where $q = \varphi(\pi)/\pi$.

Remark 3.4. If $R = \mathbf{Z}_p$, this is weaker than the definition given in [Ber04], which requires N to additionally be a free \mathbf{A}_F^+ -module.

We are interested in Wach modules because of their connection to crystalline representations. More precisely, if M is a \mathbf{Z}_p -lattice in a crystalline representation of G_F , there is an associated Wach module $\mathbf{D}(M)$, which is an \mathbf{A}_F^+ -lattice in the (φ, Γ) -module $\mathbf{D}(M)$ (which has \mathbf{A}_F -coefficients).

While $\mathbf{D}(M)$ behaves well with respect to coefficients, by the work of [Dee01], $\mathbf{D}^+(M)$ does not, nor does $\mathbf{N}(M)$. To get around this problem, we follow Kisin and construct a moduli space of \mathbf{A}_F^+ -lattices in \mathbf{A}_F -modules.

Recall that for a \mathbf{Z}_p -finite G_K -representation M , the (φ, Γ) -module $\mathbf{D}(M)$ is defined by $\mathbf{D}(M) = (\mathbf{A} \otimes_{\mathbf{Z}_p} M)^{H_K}$. By the work of Dee, if M is a finite free rank- d R -module for a finite local \mathbf{Z}_p -algebra R and the Galois action is R -linear, then $\mathbf{D}(M)$ is finite projective of rank d over $R \otimes_{\mathbf{Z}_p} \mathbf{A}_K$. Moreover, if $R \rightarrow R'$ is a homomorphism of such rings, then $\mathbf{D}(R' \otimes_R M) = R' \otimes_R \mathbf{D}(M)$.

coeffs-a-a+-i

Lemma 3.5. *Let A be an artin local \mathbf{Z}_p -algebra, and let M be a finite A -module equipped with a continuous A -linear action of G_K . Then*

- (1) *If M' is a finite A -module (with trivial G_K -action), then $\mathbf{D}(M \otimes_A M') \cong \mathbf{D}(M) \otimes_A M'$.*
- (2) *If B is an A -algebra, then $(\mathbf{A} \otimes_{\mathbf{Z}_p} M_B)^{H_K} \cong \mathbf{D}(M) \otimes_A B$.*

Proof. (1) If M' is free over A , this is clear. Otherwise, we may choose a finite presentation of M' and use the exactness of the functor $\mathbf{D}(-)$.

- (2) We may write $B = \varinjlim_i B_i$, where the B_i are A -finite submodules of B and the transition maps are injections. Then the functor of H_K -invariants commutes with the direct limit, and so does the tensor product with $\mathbf{D}(M)$, so this follows from the first part. \square

Thus, under these hypotheses we may refer to $\mathbf{D}(M_B)$ without any ambiguity.

We give a generalization of [Ber04, Lemme III.3.2]:

Lemma 3.6. *Let A be an artin local \mathbf{Z}_p -algebra with finite residue field, let B be an A -algebra, and let M be an A -linear representation of G_F of rank d . If $N \subset \mathbf{D}(M_B)$ is a free $B \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -module of rank d , stable by the actions of φ and Γ , such that $(B \otimes_{\mathbf{Z}_p} \mathbf{A}_F) \otimes_{B \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+} N = \mathbf{D}(M_B)$, then for any ideal $I \subset B$,*

$$\left((B \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_{B \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+} N \right) \cap I \left((B \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_B M_B \right) = I \left((B \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_{B \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+} N \right)$$

Proof. The proof is essentially identical to the proof of [Ber04, Lemme III.3.2]. A basis $\{n_i\}$ of N is also a basis of $\mathbf{D}(M_B)$, and therefore of $(B \otimes_{\mathbf{Z}_p} \mathbf{A}) \otimes_{B \otimes_{\mathbf{Z}_p} \mathbf{A}_F} \mathbf{D}(M_B) = (B \otimes_{\mathbf{Z}_p} \mathbf{A}) \otimes_B M_B$. If we write $x \in (\mathbf{A}^+ \otimes_{\mathbf{Z}_F^+} N) \cap I(\mathbf{A}^+ \otimes_{\mathbf{Z}_p} M_B)$ with respect to this basis as $\sum_i x_i n_i$ we see that $x_i \in IA$. But $x_i \in \mathbf{A}^+$, as well, so $x_i \in IA^+$, by Lemma 2.1. \square

Definition 3.7. Let A be an artin local \mathbf{Z}_p -algebra with finite residue field, and let M be a finite free A -module of rank d equipped with a continuous A -linear action of G_F . An $A \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -lattice in $\mathbf{D}(M)$ of height $\leq h$ is a finite projective $A \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -submodule \mathbf{N} of rank d which generates $\mathbf{D}(M)$ as an $A \otimes_{\mathbf{Z}_p} \mathbf{A}_F$ -module, such that \mathbf{N} is φ - and Γ -stable, the cokernel of $\varphi^* \mathbf{N} \rightarrow \mathbf{N}$ is killed by q^h , and Γ acts trivially on $\mathbf{N}/\pi \mathbf{N}$.

We define a functor on A -algebras by letting $L_M^{\leq h}(B)$ denote the set of $B \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -lattices in $\mathbf{D}(M \otimes_A B)$ of height $\leq h$.

Proposition 3.8. *The functor $L_M^{\leq h}(B)$ is representable by a projective A -scheme $\mathcal{L}_M^{\leq h}$. If $A \rightarrow A'$ is a map of local artin rings with finite residue field, then there is a canonical isomorphism $\mathcal{L}_M^{\leq h} \times_A A' \rightarrow \mathcal{L}_{M_{A'}}^{\leq h}$. Moreover, $\mathcal{L}_M^{\leq h}$ is equipped with a canonical very ample line bundle.*

Proof. The proof is essentially identical to the proof of [Kis08, Proposition 1.3] and [Kis09, Proposition 2.1.7]. We begin by noting that since p is torsion in A , $A \otimes_{\mathbf{Z}_p} \mathbf{A}_F = (A \otimes_{\mathbf{Z}_p} \mathcal{O}_F)[\pi][\pi^{-1}]$. Choosing a basis for $\mathbf{D}(M)$ yields a distinguished projective rank- d $A \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -module $\mathbf{N}_0 \subset \mathbf{D}(M)$ which spans $\mathbf{D}(M)$, and every $A \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -lattice \mathbf{N} satisfies $\pi^{-i}\mathbf{N}_0 \subset \mathbf{N} \subset \pi^i\mathbf{N}_0$ for some i . Thus, all $B \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -lattices are points of the affine Grassmannian for GL_d over A , which is an ind-projective scheme.

Furthermore, φ - and Γ -stability define closed conditions on the affine Grassmannian, as do the requirements that $\mathbf{N}/\varphi^*\mathbf{N}$ be killed by q^h and that Γ act trivially on $\mathbf{N}/\pi\mathbf{N}$. Finally, the φ -stability of $A \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -lattices implies that if r is the least integer such that $\pi^r\mathbf{N}_0 \subset (1 \otimes \varphi)\varphi^*(\mathbf{N}_0) \subset \pi^{-r}\mathbf{N}_0$, s is the least integer such that $p^s = 0$ in A , and $i = \max\{\frac{(p-1)sh+r}{p-1}, \frac{r}{p-1}\}$, then $\pi^{-i}\mathbf{N}_0 \subset \mathbf{N} \subset \pi^i\mathbf{N}_0$ for every $A \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -lattice \mathbf{N} .

Indeed, let B be an A -algebra, let \mathbf{N} be a $B \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -lattice, and let i be the smallest integer such that $B \widehat{\otimes}_A \mathbf{N}_0 \subset \pi^{-i}\mathbf{N}$. Then the smallest integer j such that $(1 \otimes \varphi)\varphi^*(B \widehat{\otimes}_A \mathbf{N}_0) \subset \pi^{-j}(1 \otimes \varphi)\varphi^*\mathbf{N}$ is ip . Moreover,

$$(1 \otimes \varphi)\varphi^*(B \widehat{\otimes}_A \mathbf{N}_0) \subset \pi^{-r}(B \widehat{\otimes}_A \mathbf{N}_0) \subset \pi^{-i-r}\mathbf{N} = q^{-h}\pi^{-i-r}(q^h\mathbf{N}) \subset \pi^{-i-r-h(p-1)s}(1 \otimes \varphi)\varphi^*\mathbf{N}$$

since $\pi^{(p-1)s} \in q^h(A \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+)$. Thus, $ip \leq i + r + h(p-1)s$, and so $i \leq \frac{r+h(p-1)s}{p-1}$.

Similarly, if i is the smallest integer such that $\mathbf{N} \subset \pi^{-i}(B \widehat{\otimes}_A \mathbf{N}_0)$, then

$$(1 \otimes \varphi)\varphi^*(\mathbf{N}) \subset \mathbf{N} \subset \pi^{-i}(B \widehat{\otimes}_A \mathbf{N}_0) \subset \pi^{-i-r}(1 \otimes \varphi)\varphi^*(B \widehat{\otimes}_A \mathbf{N}_0)$$

so $ip \leq i + r$ and $i \leq \frac{r}{p-1}$.

Therefore, $L_M^{\leq h}$ is representable by a closed subscheme of a Grassmannian, which is projective with a canonical very ample line bundle. \square

Let $\mathcal{N}^{\leq h}(M)$ denote the universal $\mathcal{O}_{\mathcal{L}_M^{\leq h}} \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -lattice on $\mathcal{L}_M^{\leq h}$. It is a sheaf of coherent $\mathcal{O}_{\mathcal{L}_M^{\leq h}} \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -modules, and on any open affine $\mathrm{Spec} A' \subset \mathcal{L}_M^{\leq h}$, it is an $A' \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -lattice; in particular, it is equipped with actions of φ and Γ . We write the structure morphism $\Theta_A : \mathcal{L}_M^{\leq h} \rightarrow \mathrm{Spec} A$, and we write $\tilde{A} := \Theta_{A*} \mathcal{O}_{\mathcal{L}_M^{\leq h}}$.

Lemma 3.9. *There is a canonical isomorphism $M_{\tilde{A}} \cong \left(\mathbf{A} \otimes_{\mathbf{A}_F^+} \Theta_{A*} \mathcal{N}^{\leq h}(M) \right)^{\varphi=1}$.*

Proof. By construction,

$$\Theta_A^* \mathbf{A}_F \otimes_{\Theta_A^* \mathbf{A}_F^+} \mathcal{N}^{\leq h}(M) \cong \mathbf{D}(\mathcal{O}_{\mathcal{L}_M^{\leq h}} \otimes_A M)$$

so

$$\Theta_A^* \mathbf{A} \otimes_{\Theta_A^* \mathbf{A}_F^+} \mathcal{N}^{\leq h}(M) \xrightarrow{\sim} \Theta_A^* \mathbf{A} \otimes_{\mathcal{O}_{\mathcal{L}_M^{\leq h}}} \Theta_A^* M$$

Pushing forward yields

$$\mathbf{A} \otimes_{\mathbf{A}_F^+} \Theta_{A*}(\mathcal{N}^{\leq h}(M)) \xrightarrow{\sim} \Theta_{A*} \left(\Theta_A^* \mathbf{A} \otimes_{\Theta_A^* \mathbf{A}_F^+} \mathcal{N}^{\leq h}(M) \right) \xrightarrow{\sim} \mathbf{A} \otimes_{\mathbf{Z}_p} M_{\tilde{A}}$$

by the projection formula, and taking the $\varphi = 1$ part on each side yields the desired result. \square

Lemma 3.10. *There is a finite covering $\{U_i = \mathrm{Spec} A_i\}$ of $\mathcal{L}_M^{\leq h}$ by open affine subsets such that $\mathcal{N}^{\leq h}(M)(U_i)$ is a free $A_i \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -module of rank d .*

Proof. Choose any affine covering $\{U'_j = \mathrm{Spec} A'_j\}$ of $\mathcal{L}_M^{\leq h}$; by construction, $\mathcal{N}^{\leq h}(M)(U'_j)$ is a finite projective $A'_j \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -module. Every maximal ideal of $A'_j \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ is of the form (\mathfrak{m}', π) , where \mathfrak{m}' is a maximal ideal of $A'_j \otimes_{\mathbf{Z}_p} \mathcal{O}_F$, and $\mathfrak{m}' \cap A'_j$ is a maximal ideal \mathfrak{m} such that $\mathfrak{m}(A'_j \otimes_{\mathbf{Z}_p} \mathcal{O}_F) = \prod_k \mathfrak{m}_k$ is a finite product of maximal ideals of $A'_j \otimes_{\mathbf{Z}_p} \mathcal{O}_F$. Therefore, the localization $(A'_j \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+) \otimes_{A'_j} (A'_j)_{\mathfrak{m}}$ has Jacobson radical generated by (\mathfrak{m}, π) . Then Nakayama's lemma implies that $\mathcal{N}^{\leq h}(M)(U'_j) \otimes_{A'_j} (A'_j)_{\mathfrak{m}}$ is free of rank d , so there is an open affine neighborhood $U_{j,\mathfrak{m}} = \mathrm{Spec} A_{j,\mathfrak{m}} \subset U'_j$ of the point of $\mathcal{L}_M^{\leq h}$ corresponding to \mathfrak{m} such that $\mathcal{N}^{\leq h}(M)(U_{j,\mathfrak{m}})$ is free of rank d over $A_{j,\mathfrak{m}} \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$. \square

Proposition 3.11. *There are natural inclusions of sheaves $\pi^h \left((\mathcal{O}_{\mathcal{L}_M^{\leq h}} \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_A M \right) \subset \mathbf{A}^+ \otimes_{\mathbf{A}_F^+} \mathcal{N}^{\leq h}(M) \subset (\mathcal{O}_{\mathcal{L}_M^{\leq h}} \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_A M$, and therefore natural inclusions of sheaves $\pi^h \mathbf{D}^+(\mathcal{O}_{\mathcal{L}_M^{\leq h}} \otimes_A M) \subset \mathcal{N}^{\leq h}(M) \subset \mathbf{D}^+(\mathcal{O}_{\mathcal{L}_M^{\leq h}} \otimes_A M)$.*

Proof. We first need to check that the presheaf $U \mapsto \mathbf{A}^+ \otimes_{\mathbf{A}_F^+} \mathcal{N}^{\leq h}(M)(U)$ is actually a sheaf. But this follows from Lemma 2.3 and the description of \mathbf{A}_K^+ in § 2 (which implies that $A \otimes \mathbf{A}_K^+$ is finite flat over $A \otimes \mathbf{A}_F^+$).

By Lemma 3.10, $\mathcal{N}^{\leq h}(M)$ is free locally on $\mathcal{L}_M^{\leq h}$; let $\{U_i\}$ be an affine cover of $\mathcal{L}_M^{\leq h}$ such that $\mathcal{N}^{\leq h}(M)(U_i)$ is free for all i .

Let $X_{U_i} \in \text{Mat}_{d \times d}(\mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_{\mathbf{Z}_p} \mathbf{A})$ be a matrix of a basis of $\mathcal{N}^{\leq h}(M)(U_i)$ with respect to a basis of $\mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_A M$. Since p is torsion in A , $\mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_{\mathbf{Z}_p} \mathbf{A} = \mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_{\mathbf{Z}_p} \mathbf{A}^+[1/\pi]$. Moreover, there is some integer $s > 0$ such that $\varphi^s(\pi) \in \pi^p(\mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_{\mathbf{Z}_p} \mathbf{A}^+)$, because $\varphi^s(\pi) = (1 + \pi)^{p^s} - 1$ and $\text{ord}_p \binom{p^s}{n} = s$ for $1 \leq n \leq p - 1$; let P_{U_i} be the matrix of φ^s with respect to the chosen basis of $\mathcal{N}^{\leq h}(M)(U_i)$. Since the action of Frobenius is trivial on M , we see that $X_{U_i} P_{U_i} = \varphi^s(X_{U_i})$, so $\varphi^s(\pi^h X_{U_i}^{-1}) = ((\varphi^{s-1}(q) \cdots \varphi(q)q)^h P_{U_i}^{-1})(\pi^h X_{U_i}^{-1})$.

If X_{U_i} has a coefficient in $\pi^{-r} \mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_{\mathbf{Z}_p} \mathbf{A}^+ \setminus \pi^{-r+1} \mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_{\mathbf{Z}_p} \mathbf{A}^+$ for $r \geq 0$ (and no coefficients with poles of higher order), then $\varphi^s(X_{U_i})$ has a coefficient in $\pi^{-r+p} \mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_{\mathbf{Z}_p} \mathbf{A}^+ \setminus \pi^{-r+p+1} \mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_{\mathbf{Z}_p} \mathbf{A}^+$, whereas $X_{U_i} P_{U_i}$ has coefficients in $\pi^{-r} \mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_{\mathbf{Z}_p} \mathbf{A}^+$. It follows that $r = 0$ and $\mathcal{N}^{\leq h}(M)(U_i) \subset \mathbf{A}^+ \widehat{\otimes}_{\mathbf{Z}_p} (\mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_A M)$.

The other inclusion follows similarly. Since $\mathcal{N}^{\leq h}(M)/\varphi^* \mathcal{N}^{\leq h}(M)$ is killed by q^h by assumption, $((\varphi^{s-1}(q) \cdots \varphi(q)q)^h P_{U_i}^{-1})$ has coefficients in $\mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$. Now if $\pi^h X_{U_i}^{-1}$ has a coefficient with a pole of order r (and no coefficients with poles of higher order), then $\varphi^s(\pi^h X_{U_i}^{-1})$ has a coefficient with a pole of order at least pr , whereas $((\varphi^{s-1}(q) \cdots \varphi(q)q)^h P_{U_i}^{-1})(\pi^h X_{U_i}^{-1})$ has coefficients in $\pi^{-r} \mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_{\mathbf{Z}_p} \mathbf{A}^+$. It follows that $r = 0$ and therefore $\pi^h \left((\mathcal{O}_{\mathcal{L}_M^{\leq h}}(U_i) \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_A M \right) \subset \mathbf{A}^+ \widehat{\otimes}_{\mathbf{A}_F^+} \mathcal{N}^{\leq h}(M)(U_i)$.

Now we take H_F -invariants to obtain the second pair of inclusions. To see that $\left(\mathbf{A}^+ \otimes_{\mathbf{A}_F^+} \mathcal{N}^{\leq h}(M) \right)^{H_F} = \mathcal{N}^{\leq h}(M)$, we observe that the presheaf $U \mapsto \left(\mathbf{A}^+ \otimes_{\mathbf{A}_F^+} \mathcal{N}^{\leq h}(M)(U) \right)^{H_F}$ is a sheaf on $\mathcal{L}_M^{\leq h}$ and the set of open affine subsets $U \subset \mathcal{L}_M^{\leq h}$ such that $\mathcal{N}^{\leq h}(M)(U)$ is free is a basis for the Zariski topology. But if $\mathcal{N}^{\leq h}(M)(U)$ is free, it is clear that $\left(\mathbf{A}^+ \otimes_{\mathbf{A}_F^+} \mathcal{N}^{\leq h}(M)(U) \right)^{H_F} = \mathcal{N}^{\leq h}(M)(U)$, so we are done. \square

Lemma 3.12. *Let $A \rightarrow A'$ be a local homomorphism of local artin rings with finite residue fields. Then there is a homomorphism $A' \otimes_A \Gamma(\mathcal{L}_M^{\leq h}, \mathcal{N}^{\leq h}(M)) \rightarrow \Gamma(\mathcal{L}_{M_{A'}}^{\leq h}, \mathcal{N}^{\leq h}(M_{A'}))$, and the image contains the image of $\pi^h \mathbf{D}^+(M_{A'})$.*

Proof. By the base change properties of Θ_A and $\mathcal{L}_M^{\leq h}$ we have a commutative diagram

$$\begin{array}{ccccc} A' \otimes_A \pi^h((A \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_A M) & \longrightarrow & A' \otimes_A \pi^h((\tilde{A} \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_A M) & \longrightarrow & A' \otimes_A \left(\mathbf{A}^+ \otimes_{\mathbf{A}_F^+} \mathcal{N}^{\leq h}(M)(\mathcal{L}_M^{\leq h}) \right) \\ \parallel & & \downarrow & & \downarrow \\ \pi^h((A' \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_{A'} M_{A'}) & \longrightarrow & \pi^h((\tilde{A}' \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_{A'} M_{A'}) & \xrightarrow{\subset} & \mathbf{A}^+ \otimes_{\mathbf{A}_F^+} \mathcal{N}^{\leq h}(M_{A'})(\mathcal{L}_{M_{A'}}^{\leq h}) \end{array}$$

We first observe that if A' is a finite free A -module, then $\mathbf{D}^+(M_{A'}) = A' \otimes_A \mathbf{D}^+(M)$ and the result follows by taking H_F -invariants of the diagram. Thus, we may assume that $A \rightarrow A'$ induces an isomorphism on residue fields. Indeed, if k' denotes the residue field of A' , then $W(k')$ is a finite free \mathbf{Z}_p -module and A' is a $W(k')$ -algebra by Hensel's lemma. Then $W(k') \otimes_{\mathbf{Z}_p} A$ is a finite free A -module and there is a natural homomorphism $W(k') \otimes_{\mathbf{Z}_p} A \rightarrow A'$; the source is semi-local, but its local summands are summands of a finite free module, hence projective, hence free, and they have residue field k' .

Moreover, for some $N \gg 0$ and some $k \geq 0$, there is a surjection $A[X_1, \dots, X_k]/(X_1, \dots, X_k)^N \twoheadrightarrow A'$ sending the X_i to elements of $\mathfrak{m}_{A'}$. Since $A[X_1, \dots, X_k]/(X_1, \dots, X_k)^N$ is finite free over A , we may assume $A \rightarrow A'$ is surjective.

So suppose that $A \rightarrow A'$ is surjective with kernel I . Given an element $\beta \in \mathbf{A}^+ \otimes_{\mathbf{A}_F^+} \Gamma(\mathcal{L}_{M_A'}^{\leq h}, \mathcal{N}^{\leq h}(M_{A'}))$ in the image of $\pi^h \mathbf{D}^+(M_{A'})$, by the commutativity of the above diagram it lifts to an element $\tilde{\beta} \in \mathbf{A}^+ \otimes_{\mathbf{A}_F^+} \Gamma(\mathcal{L}_M^{\leq h}, \mathcal{N}^{\leq h}(M))$. Since $\tilde{\beta}$ is defined up to an element of $\pi^h I(\mathbf{A}^+ \otimes_A M)$, we need to show that

$$\tilde{\beta} \in \mathcal{N}^{\leq h}(M)(\mathcal{L}_M^{\leq h}) + I(A \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_{A \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+} \mathcal{N}^{\leq h}(M)(\mathcal{L}_M^{\leq h})$$

In order to show this, we first note that for any $h \in H_F$,

$$h(\tilde{\beta}) - \tilde{\beta} \in I((A \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_A M) \cap \mathbf{A}^+ \otimes_{\mathbf{A}_F^+} \mathcal{N}^{\leq h}(M)(\mathcal{L}_M^{\leq h})$$

since β is fixed by H_F . If $U = \text{Spec } A_U \subset \mathcal{L}_M^{\leq h}$ is an open affine subspace such that $\mathcal{N}^{\leq h}(M)(U)$ is free (compare Lemma 3.10), then $h(\tilde{\beta}) - \tilde{\beta}|_U \in I(\mathbf{A}^+ \otimes_{\mathbf{A}_F^+} \mathcal{N}^{\leq h}(M)(U))$ by Lemma 3.5. Indeed, if n_1, \dots, n_d is a basis for $\mathcal{N}^{\leq h}(M)(U)$ and $\tilde{\beta} = \sum_i b_i \otimes n_i$ for some $b_i \in A_U \otimes_{\mathbf{Z}_p} \mathbf{A}^+$, then $h(b_i) - b_i \in I(A_U \otimes_{\mathbf{Z}_p} \mathbf{A}^+)$. Since $(A_U/I) \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+ \rightarrow ((A_U/I) \otimes_{\mathbf{Z}_p} \mathbf{A}^+)^{H_F}$ is an isomorphism, $b_i \in A_U \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+ + IA_U \otimes_{\mathbf{Z}_p} \mathbf{A}^+$. Therefore,

$$\tilde{\beta}|_U \in \mathcal{N}^{\leq h}(M)(U) + I(A_U \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_{A_U \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+} \mathcal{N}^{\leq h}(M)(U).$$

Now consider $\tilde{\beta}|_{U \cap U'}$. If $\tilde{\beta}|_U = n_U + b_U$ and $\tilde{\beta}|_{U'} = n_{U'} + b_{U'}$, with $n_U, n_{U'} \in \mathcal{N}^{\leq h}(M)(U \cap U')$ and $b_U, b_{U'} \in I(A_{U \cap U'} \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_{A_{U \cap U'} \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+} \mathcal{N}^{\leq h}(M)(U \cap U')$, then $n_U - n_{U'} = b_{U'} - b_U$ as elements of the free $(A_{U \cap U'} \otimes_{\mathbf{Z}_p} \mathbf{A}^+)$ -module $(A_{U \cap U'} \otimes_{\mathbf{Z}_p} \mathbf{A}^+) \otimes_{A_{U \cap U'} \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+} \mathcal{N}^{\leq h}(M)(U \cap U')$. But $n_U - n_{U'}$ is fixed by H_F , by construction, so $b_{U'} - b_U$ must have coefficients in $(I(A_{U \cap U'} \otimes_{\mathbf{Z}_p} \mathbf{A}^+))^{H_F} = I(A_{U \cap U'} \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+)$. Thus, $U \cap U' \mapsto b_{U'} - b_U$ is a 1-cocycle valued in $I \mathcal{N}^{\leq h}(M)$, and when viewed as a 1-cocycle valued in $I(\mathbf{A}^+) \otimes_{\mathbf{A}_F^+} \mathcal{N}^{\leq h}(M)$, it is trivial (since it is the 1-coboundary of $U \mapsto b_U$). But all of the b_U have coefficients in $IA \otimes_{\mathbf{Z}_p} \mathbf{A}_K^+$ for some finite extension K/F by Lemma 2.3, and \mathbf{A}_K^+ is a finite free \mathbf{A}_F^+ -module by the description of \mathbf{A}_K^+ in 2. Therefore, there exist $\tilde{b}_U \in IA \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ such that the 1-coboundary $U \cap U' \mapsto b_{U'} - b_U$ is equal to the 1-coboundary $U \cap U' \mapsto \tilde{b}_{U'} - \tilde{b}_U$ (we may choose \tilde{b}_U by projecting the coefficients of b_U down to $IA \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$). Now we rewrite $\tilde{\beta}|_U = (n_U + \tilde{b}_U) + (b_U - \tilde{b}_U)$ and we see that $\{n_U + \tilde{b}_U\}$ glues to a global section of $\mathcal{N}^{\leq h}(M)$, so we are done. \square

Now we wish to consider the situation when A is a complete local noetherian \mathbf{Z}_p -algebra with finite residue field and maximal ideal \mathfrak{m}_A . We may extend the definition of the functor $L_M^{\leq h}(B)$ to A -algebras B such that $\mathfrak{m}_A^i B = 0$ for some i in a natural way. It follows from formal GAGA that $L_M^{\leq h}$ is also representable in this setting by a projective A -scheme. Namely, we consider the formal scheme $\widehat{\mathcal{L}}_M^{\leq h} := \{\widehat{\mathcal{L}}_{M_{A/\mathfrak{m}_A^i}}^{\leq h}\}$, which is equipped with a morphism $\widehat{\Theta}_{A^*} : \widehat{\mathcal{L}}_M^{\leq h} \rightarrow \text{Spf}(A)$; it is a formal scheme equipped with a very ample line bundle, so it is the \mathfrak{m}_A -adic completion of a projective A -scheme $\mathcal{L}_M^{\leq h}$. As in the artinian case, we write the projective structure morphism $\Theta_A : \mathcal{L}_M^{\leq h} \rightarrow \text{Spec } A$, and we write $\tilde{A} := \Theta_{A^*} \mathcal{O}_{\mathcal{L}_M^{\leq h}}$.

Then $\widehat{\mathcal{L}}_M^{\leq h}$ carries a universal $\mathcal{O}_{\widehat{\mathcal{L}}_M^{\leq h}} \widehat{\mathbf{A}}_F^+$ -lattice $\widehat{\mathcal{N}}^{\leq h}(M)$. We may view $\widehat{\mathcal{N}}^{\leq h}(M)$ as a formal coherent sheaf on $\widehat{\mathcal{L}}_M^{\leq h} \times \text{Spf } \mathbf{A}_F^+ \cong \widehat{\mathcal{L}}_M^{\leq h} \times_{\text{Spf } A} \text{Spf}(A \widehat{\otimes} \mathbf{A}_F^+)$, and by formal GAGA, $\widehat{\mathcal{N}}^{\leq h}(M)$ is the completion of a coherent sheaf (in fact, a vector bundle) on $\mathcal{L}_M^{\leq h} \times_{\text{Spec } A} \text{Spec}(A \widehat{\otimes} \mathbf{A}_F^+)$.

If $A \rightarrow A'$ is a local homomorphism of local rings and A' is \mathfrak{m}_A -adically complete, then an A' -point of $\mathcal{L}_M^{\leq h}$ induces a $\{A'/\mathfrak{m}_A^i\}$ -point of $\widehat{\mathcal{L}}_M^{\leq h}$, and therefore a system of $(A'/\mathfrak{m}_A^i) \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -lattices in $\{\mathbf{D}(M_{A'/\mathfrak{m}_A^i})\}$. But this is the same as an $A' \widehat{\otimes} \mathbf{A}_F^+$ -lattice in $\mathbf{D}(M_{A'})$, where A' is given the \mathfrak{m}_A -adic topology. Thus, we may view $\mathcal{L}_M^{\leq h}$ as a moduli space of $A \widehat{\otimes} \mathbf{A}_F^+$ -lattices in $\mathbf{D}(M)$. In particular, \mathbf{Z}_p -points of $\mathcal{L}_M^{\leq h}$ (if they exist) correspond to Wach modules of $M_{\mathbf{Z}_p}$ in the sense defined in [Ber04], i.e., satisfying the properties given in Def. 3.3.

Lemma 3.13. *There is a canonical isomorphism $M_{\tilde{A}} \cong \left(\mathbf{A} \otimes_{\mathbf{A}_F^+} \Theta_{A^*} \mathcal{N}^{\leq h}(M) \right)^{\varphi=1}$.*

Proof. This follows from 3.9 by taking limits. \square

Lemma 3.14. *Let A be a finite flat \mathbf{Z}_p -algebra, let $B := A[1/p]$, and let M be a finite free A -module of rank d equipped with a continuous A -linear action of G_F such that the underlying \mathbf{Z}_p -linear Galois*

representation is a lattice in a crystalline representation with Hodge–Tate weights in $[-h, 0]$. Then $\Theta_{A^*} \mathcal{N}^{\leq h}(M)[1/p]$ is free over $B \otimes_{\mathbf{Q}_p} \mathbf{B}_F^+$ of rank d .

Proof. We first forget the A -linear structures on M and $\Theta_{A^*} \mathcal{N}^{\leq h}(M)$ and consider them as \mathbf{Z}_p - and \mathbf{A}_F^+ -modules, respectively. Then $\mathcal{L}_M^{\leq h} \rightarrow \text{Spec } \mathbf{Z}_p$ has a unique point in the fiber over $\text{Spec } \mathbf{Q}_p$. Indeed, any \mathbf{Q}_p -point extends to a \mathbf{Z}_p -point of $\mathcal{L}_M^{\leq h}$, which induces an \mathbf{A}_F^+ -submodule \mathbf{N} of $\mathbf{D}(M)$ with actions of φ and Γ , such that \mathbf{N}/φ^* is annihilated by q^h and Γ acts trivially on \mathbf{N}/π . Therefore, $\Theta_{\mathbf{Z}_p^*} \mathcal{N}^{\leq h}(M)_{\mathbf{Q}_p}$ is the unique rational Wach module $\mathbf{N}(M[1/p])$ constructed in [Ber04].

Next, recall that $\mathbf{D}_{\text{cris}}(M_B) \cong \mathbf{N}(M)[1/p]/\pi \mathbf{N}(M)[1/p]$. But by [Bel15, Proposition 4.1.3], $\mathbf{D}_{\text{cris}}(M)$ is free over $B \otimes_{\mathbf{Q}_p} F$ of rank d , so we are done by Nakayama’s lemma. \square

Lemma 3.15. *Let N be a finite torsion-free \mathbf{A}_F^+ -module of generic rank d and define $N^{\text{sat}} := N[1/p] \cap (N \otimes_{\mathbf{A}_F^+} \mathbf{A}_F)$, where the intersection is taken inside the finite \mathbf{B}_F -vector space $(N \otimes_{\mathbf{A}_F^+} \mathbf{A}_F)[1/p]$. Then N^{sat} is a finite free \mathbf{A}_F^+ -module of rank d .*

Proof. We first reduce to the case where N is free of rank d over \mathbf{A}_F^+ . Indeed, consider the kernel of the natural map $N \rightarrow N[1/p]/\pi$. This is a finite \mathbf{A}_F^+ -module generated by elements $n_1, \dots, n_{d'} \in N$ such that $p^{e_i} n_i = \pi^{f_i} n'_i$ for all i , for some integers e_i, f_i and $n'_i \in N$. Clearly the elements $n_i/\pi^{f_i} = n'_i/p^{e_i} \in (N \otimes_{\mathbf{A}_F^+} \mathbf{A}_F)[1/p]$ live in N^{sat} , so we may replace N by $N + \mathbf{A}_F^+(\frac{n_1}{\pi^{f_1}}) + \dots + \mathbf{A}_F^+(\frac{n_{d'}}{\pi^{f_{d'}}})$. Now the natural map $N/\pi \rightarrow N[1/p]/\pi$ is injective. But the image of N/π in $N[1/p]/\pi$ is a lattice in a d -dimensional F -vector space, so it is free of rank d over \mathcal{O}_F . Then by Nakayama’s lemma, there is a surjection $(\mathbf{A}_F^+)^{\oplus d} \twoheadrightarrow N$; the kernel must be p -torsion, hence trivial, so this surjection must be an isomorphism.

We may thus assume that N is free of rank d over \mathbf{A}_F^+ . But it is clear that

$$((\mathbf{A}_F^+)^{\oplus d})^{\text{sat}} = ((\mathbf{A}_F^+)^{\text{sat}})^{\oplus d} = (\mathbf{A}_F^+)^{\oplus d}$$

so we are done. \square

The key result is the following, which is the same argument as in the proof of [Kis08, Proposition 1.6.4]:

Proposition 3.16. *Let A be a complete local noetherian \mathbf{Z}_p -algebra together with a finite free A -module M of rank d , and a continuous A -linear action of G_F on M .*

- (1) *The map $\Theta_A : \mathcal{L}_M^{\leq h} \rightarrow \text{Spec } A$ becomes a closed immersion after inverting p .*
- (2) *Let $A^{\leq h}$ denote the quotient of A corresponding to the scheme-theoretic image of Θ_A . Then for any \mathbf{Q}_p -finite artin ring B , a map $A \rightarrow B$ factors through $A^{\leq h}$ if and only if M_B is crystalline with Hodge–Tate weights in $[-h, 0]$.*

Proof. The first part essentially follows from the uniqueness of Wach modules for crystalline representations of G_F cf. [Ber04, Proposition III.4.2]. Indeed, if B is a \mathbf{Q}_p -finite local artin ring, a B -point $x : \text{Spec } B \rightarrow \text{Spec } A[1/p]$ is induced by a homomorphism $A \rightarrow B^\circ$, where $B^\circ \subset B$ is \mathbf{Z}_p -finite and flat. Any B -point of $\mathcal{L}_M^{\leq h}$ over x extends to a B° -point, by the valuative criterion for properness. This corresponds to a system of $(B/\mathfrak{m}_A^n) \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -lattices in $\mathbf{D}(M \otimes_A B/\mathfrak{m}_A^n)$, and therefore induces a $B^\circ \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -submodule of $\mathbf{D}(M \otimes_A B^\circ)$. Inverting p and forgetting about the B -linear structure, we obtain a \mathbf{B}_F^+ -lattice in $\mathbf{D}(M \otimes_R B)$. But there is at most one such lattice, by [Ber04, Proposition III.4.2].

Thus, $\Theta_A[1/p] : \mathcal{L}_M^{\leq h}[1/p] \rightarrow \text{Spec } A[1/p]$ is quasi-finite; since it is projective, it is therefore finite (and is an injection on closed points). Setting $B = E[\varepsilon]/\varepsilon^2$, it also induces injections on tangent spaces, and is therefore a closed immersion.

For the second part, we first consider a \mathbf{Q}_p -finite local artin ring B with residue field E together with a map $\text{Spec } B \rightarrow \text{Spec } A$ which factors through the image of $\mathcal{L}_M^{\leq h}$. We obtain a Wach module with coefficients in $B \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$. Forgetting the B -linear structure, we get a Wach module, so again by [Ber04, Proposition III.4.2], M_B is crystalline with Hodge–Tate weights in $[-h, 0]$.

It remains to show that if M_B is crystalline with Hodge–Tate weights in $[-h, 0]$, it is induced by a point of $\mathcal{L}_M^{\leq h}$. For this, we introduce some new notation: Let $B^\circ \subset B$ denote the subring of elements mapping to $\mathcal{O}_E \subset E$, and we let Int_B denote the set of \mathcal{O}_E -finite subrings of B° (this makes sense because B is canonically an E -algebra). Each such C is a finite flat \mathcal{O}_E -algebra, since B contains no p -torsion.

Observe that the map $A \rightarrow B$ factors through some $C \in \text{Int}_B$, and we may assume that the map $C \rightarrow \mathcal{O}_E$ is a surjection. Then M_C is a lattice in the crystalline representation M_B , so there is some Wach module $\mathbf{N}(M_C) \subset \mathbf{D}(M_C)$. It is an \mathbf{A}_F^+ -lattice in $\mathbf{D}(M_C)$ (forgetting the C -linear structure), and $\mathbf{N}(M_C)[1/p]$ is projective of rank d over $(C \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+)[1/p]$ by Lemma 3.14. Let $\mathbf{N}'_{\mathcal{O}_E}$ be the image of $\mathbf{N}(M_C)$ under the map $C \rightarrow \mathcal{O}_E$ induced by the projection $B \rightarrow E$, and let $\mathbf{N}_{\mathcal{O}_E}$ be the saturation of $\mathbf{N}'_{\mathcal{O}_E}$:

$$\mathbf{N}_{\mathcal{O}_E} := \mathbf{A}_F \otimes_{\mathbf{A}_F^+} \mathbf{N}'_{\mathcal{O}_E} \cap \mathbf{N}'_{\mathcal{O}_E}[1/p]$$

where the intersection is taken inside $\mathbf{D}(M_{\mathcal{O}_E})[1/p]$.

It turns out that $\mathbf{N}_{\mathcal{O}_E}$ is a finite free $\mathcal{O}_E \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -module of rank d . Indeed, it is free over \mathbf{A}_F^+ of rank $d \text{rk}_{\mathbf{Z}_p} \mathcal{O}_E$ by Lemma 3.15, so $\mathbf{N}_{\mathcal{O}_E}/\pi$ is p -torsion-free, hence a lattice in the free $E \otimes_{\mathbf{Q}_p} F$ -module $(\mathbf{N}_{\mathcal{O}_E}/\pi)[1/p]$, hence free over $\mathcal{O}_E \otimes_{\mathbf{Z}_p} \mathcal{O}$ of rank d .

If we choose an $\mathcal{O}_E \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -basis of $\mathbf{N}_{\mathcal{O}_E}$, we can lift it to a $(C \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+)[1/p]$ -basis of $\mathbf{N}(M_C)[1/p]$. After enlarging C to $C' \in \text{Int}_B$, we may assume that the matrices of φ and Γ with respect to this basis have coefficients in $C' \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$. Let $\mathbf{N}_{C'}$ denote the $C' \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$ -module spanned by this basis; $\mathbf{N}_{C'}$ is free over $C' \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$, so in particular is flat over \mathbf{A}_F^+ . Then $\mathbf{N}_{C'}$ is φ - and Γ -stable.

It is clear that Γ acts trivially on $\mathbf{N}_{C'}/\pi$, since this is true for $\mathbf{N}(M_C)$ and therefore for $\mathbf{N}(M_C)[1/p]$. It remains to see that q^h annihilates $\mathbf{N}_{C'}/\varphi^* \mathbf{N}_{C'}$. Certainly this is true after inverting p , since $\mathbf{N}_{C'}[1/p] = \mathbf{N}(M_C)[1/p]$ is a Wach module. So we need to see that $\mathbf{N}_{C'}/\varphi^* \mathbf{N}_{C'}$ has no p -torsion.

To see this, we observe that there is a surjection $C' \rightarrow \mathcal{O}_E$, with kernel \mathfrak{n} , and there is a descending filtration $\{\mathfrak{n}_i\}$ of \mathfrak{n} by subideals such that $\mathfrak{n}_i/\mathfrak{n}_{i+1} \cong \mathcal{O}_E$. Since the action of φ on $\mathbf{N}_{C'}$ is C' -linear, $\mathfrak{n}_i \mathbf{N}_{C'}/\mathfrak{n}_{i+1}$ is isomorphic to $\mathbf{N}_{\mathcal{O}_E}$ as a φ -module. It follows that $\mathbf{N}_{C'}/\varphi^* \mathbf{N}_{C'}$ is an extension of copies of $\mathbf{N}_{\mathcal{O}_E}/\varphi^* \mathbf{N}_{\mathcal{O}_E}$, and is therefore p -torsion free. \square

The upshot of all of this is that if M is a rank- d family of A -linear representations of G_F , $\mathcal{L}_M^{\leq h}$ carries a family $\mathcal{N}^{\leq h}(M)$ of $\mathcal{O}_{\mathcal{L}_M^{\leq h}} \widehat{\otimes}_{\mathbf{Z}_p} \mathbf{A}_F^+$ -modules whose \mathfrak{m}_A -adic completion is the universal family of $\mathcal{O}_{\widehat{\mathcal{L}}_M^{\leq h}} \widehat{\otimes}_{\mathbf{Z}_p} \mathbf{A}_F^+$ -lattices of height $\leq h$. The image of $\Theta : \mathcal{L}_M^{\leq h}[1/p] \rightarrow \text{Spec } A[1/p]$ is exactly the crystalline points with Hodge–Tate weights in the interval $[-h, 0]$.

Definition 3.17. Let M be a family of crystalline representations over A with Hodge–Tate weights in $[-h, 0]$. We define the *total Wach module* to be $\mathbf{N}(M)^{\text{total}} := \Theta_{A^*} \mathcal{N}^{\leq h}(M)$. If M is a family of crystalline representations with Hodge–Tate weights in $[a, b]$, we define $\mathbf{N}(M)^{\text{total}} := \pi^{-b} \mathbf{N}(M(-b))^{\text{total}}$.

Let $\mathcal{I} \subset \mathcal{O}_{\mathcal{L}_M^{\leq h}}$ denotes the ideal sheaf generated by all p -power torsion sections, and let $\mathcal{L}_M^{\leq h, \text{t.f.}}$ denote the corresponding closed subscheme. Let $\mathcal{N}^{\leq h, \text{t.f.}}(M)$ denote the restriction of $\mathcal{N}^{\leq h}(M)$ to $\mathcal{L}_M^{\leq h, \text{t.f.}}$, and write $\mathbf{N}(M)^{\text{t.f.}} := \Theta_{A^*} \mathcal{N}^{\leq h, \text{t.f.}}(M) = \mathbf{N}(M)^{\text{total}}/\mathbf{N}(M)^{p\text{-tors}}$.

We observe that by the theorem on formal functions, the \mathfrak{m}_A -adic completion of $\mathbf{N}(M)^{\text{total}}$ is $\widehat{\Theta}_{A^*} \widehat{\mathcal{N}}^{\leq h}(M)$.

Since Θ_A is a projective map, $\Theta_{A^*} \mathcal{O}_{\mathcal{L}_M^{\leq h}}$ is A -finite, and $\mathbf{N}(M)^{\text{total}}$ is a finite $A \widehat{\otimes}_{\mathbf{Z}_p} \mathbf{A}_F^+$ -module. Since $\Theta_A[1/p] : \mathcal{L}_M^{\leq h}[1/p] \rightarrow \text{Spec } A[1/p]$ is an isomorphism, $\mathbf{N}(M)^{\text{total}}[1/p]$ is projective of rank d and its formation commutes with base change on $A[1/p]$. However, $\mathbf{N}(M)^{\text{total}}$ may not be projective itself.

We record a useful lemma about $\mathbf{N}(M)^{\text{total}}[1/p]$:

Lemma 3.18. *Let \mathbf{N} be a finite $A \widehat{\otimes}_{\mathbf{Z}_p} \mathbf{A}_F^+$ -module such that $\mathbf{N}[1/p]$ is projective. Then there is an affine cover $\{U_k = \text{Spec } A[1/p, 1/f_k]\}$ of $\text{Spec } A[1/p]$ such that $\mathbf{N} \otimes_A A[1/p, 1/f_k]$ is free.*

Proof. It suffices to show that if $\mathfrak{m} \subset A[1/p]$ is a maximal ideal, then $\mathbf{N}[1/p] \otimes_{A[1/p]} A[1/p]_{\mathfrak{m}}$ is free of rank d . If $K := A[1/p]_{\mathfrak{m}}$, then K is a finite extension of \mathbf{Q}_p with ring of integers \mathcal{O} and $(A \widehat{\otimes}_{\mathbf{Z}_p} \mathbf{A}_F^+)[1/p]_{\mathfrak{m}} \cong (\mathcal{O}_K \otimes_{\mathbf{Z}_p} \mathcal{O}_F)[\pi][1/p]$, which is a product of finitely many principal ideal domains. Therefore, $\mathbf{N}[1/p] \otimes_{A[1/p]} A[1/p]_{\mathfrak{m}}$ is free of rank d . But $(A \widehat{\otimes}_{\mathbf{Z}_p} \mathbf{A}_F^+)[1/p] \otimes_{A[1/p]} A[1/p]_{\mathfrak{m}}$ has Jacobson radical generated by \mathfrak{m} , so by Nakayama's lemma, we can lift d generators of $\mathbf{N}[1/p] \otimes_{A[1/p]} A[1/p]_{\mathfrak{m}}$ to generators of $\mathbf{N}[1/p] \otimes_{A[1/p]} A[1/p]_{\mathfrak{m}}$. Since a surjection of projective modules of the same rank is an isomorphism, we are done. \square

Lemma 3.19. *There are natural inclusions of sheaves*

$$\pi^h \left((\mathcal{O}_{\widehat{\mathcal{L}}_M^{\leq h}} \widehat{\otimes} \mathbf{A}^+) \otimes_A M \right) \subset (\mathcal{O}_{\widehat{\mathcal{L}}_M^{\leq h}} \widehat{\otimes} \mathbf{A}^+) \otimes_{\mathcal{O}_{\widehat{\mathcal{L}}_M^{\leq h}}} \widehat{\otimes}_{\mathbf{Z}_p} \mathbf{A}_F^+ \widehat{\mathcal{N}}^{\leq h}(M) \subset (\mathcal{O}_{\widehat{\mathcal{L}}_M^{\leq h}} \widehat{\otimes} \mathbf{A}^+) \otimes_A M$$

and therefore natural inclusions of sheaves

$$\pi^h \mathbf{D}^+(\mathcal{O}_{\widehat{\mathcal{Z}}_M^{\leq h}} \otimes_A M) \subset \widehat{\mathcal{N}}^{\leq h}(M) \subset \mathbf{D}^+(\mathcal{O}_{\widehat{\mathcal{Z}}_M^{\leq h}} \otimes_A M)$$

Proof. This follows from Proposition 3.11 by taking limits. \square

Remark 3.20. Combined with Lemma 3.13, this implies that the representation $M_{\widehat{A}}$ is finite height, i.e., $\mathbf{D}(M_{\widehat{A}})$ is generated by elements of $\mathbf{D}^+(M_{\widehat{A}})$.

Proposition 3.21. *Let $A \rightarrow A'$ be a local homomorphism of complete local noetherian \mathbf{Z}_p -algebras with finite residue field. Then there is a natural base change map $A' \widehat{\otimes}_A \mathbf{N}(M)^{\text{total}} \rightarrow \mathbf{N}(M_{A'})^{\text{total}}$, and its image contains the image of $\pi^h \mathbf{D}^+(M_{A'})$.*

Proof. This follows from Lemma 3.12 by taking limits. \square

We wish to compare this construction with Berger's construction when $A = \mathbf{Z}_p$ and M is a lattice in a crystalline representation with Hodge–Tate weights in $[-h, 0]$. To show they agree, we would have to prove that $\mathbf{N}(M)$ is free over \mathbf{A}_F^+ . There are two problems: A priori, there could be connected components of $\mathcal{L}_M^{\leq h}$ supported on the special fiber and killed by inverting p , which would lead to p -power torsion in $\mathbf{N}(M)^{\text{total}}$. In addition, even after killing p -torsion we would only know that Berger's Wach module is the p -saturation of ours in $\mathbf{D}(M)$.

We now consider only the situation when A is flat over \mathbf{Z}_p , A is integrally closed in $A[1/p]$, and M is a crystalline family of G_F -representations with coefficients in A . In particular, $A \rightarrow \widetilde{A}$ is injective, $A[1/p] \rightarrow \widetilde{A}[1/p]$ is an isomorphism, and $A \rightarrow \widetilde{A}^{\text{t.f.}}$ is an isomorphism.

By construction, there is a natural inclusion $\widehat{\mathcal{N}}^{\leq h}(M) \rightarrow \mathbf{D}(\mathcal{O}_{\widehat{\mathcal{Z}}_M^{\leq h}} \otimes_A M)$, and therefore a natural inclusion $\mathbf{N}(M)^{\text{t.f.}}[1/p] = \mathbf{N}(M)^{\text{total}}[1/p] \rightarrow \mathbf{D}(M)[1/p]$ (we use the fact that $A \rightarrow \widetilde{A}$ is a finite morphism). We define the *saturation* of $\mathbf{N}(M)^{\text{total}}$ by setting $\mathbf{N}(M)^{\text{sat}} := \mathbf{N}(M)^{\text{total}}[1/p] \cap \mathbf{D}(M)$, where the intersection takes place inside $\mathbf{D}(M)[1/p]$. Note that if A is \mathbf{Z}_p -finite and flat, this agrees with Berger's definition of the Wach module.

Lemma 3.22. *Let A and M be as above. Then $\mathbf{N}(M)^{\text{sat}}[1/p] = \mathbf{N}(M)^{\text{total}}[1/p]$.*

Proof. Clearly, $\mathbf{N}(M)^{\text{sat}}[1/p] \subset \mathbf{N}(M)^{\text{total}}[1/p]$. For the other direction, choose $n \in \mathbf{N}(M)^{\text{total}} \subset \mathbf{D}(M_{\widehat{A}})$. Since $A \rightarrow \widetilde{A}$ is finite and $A[1/p] \rightarrow \widetilde{A}[1/p]$ is an isomorphism, there is some integer $k \geq 0$ such that $p^k(\widetilde{A} \widehat{\otimes} \mathbf{A}) \subset \widetilde{A} \widehat{\otimes} \mathbf{A}$. This implies that $p^k \mathbf{D}(M_{\widehat{A}}) \subset \mathbf{D}(M)$, so that $p^k n \in \mathbf{D}(M) \cap \mathbf{N}(M)^{\text{total}} \subset \mathbf{N}(M)^{\text{sat}}$. \square

Lemma 3.23. *The natural map $(A \widehat{\otimes} \mathbf{A}_F) \otimes_{A \widehat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M)^{\text{sat}} \rightarrow \mathbf{D}(M)$ is surjective.*

Proof. In the proofs of Lemma 3.9 and Lemma 3.13 we showed that the natural map $(\widetilde{A} \widehat{\otimes} \mathbf{A}_F) \otimes_{\widetilde{A} \widehat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M)^{\text{total}} \rightarrow \mathbf{D}(M_{\widehat{A}})$ is surjective. It follows that the natural map $(A \widehat{\otimes} \mathbf{A}_F) \otimes_{A \widehat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M)^{\text{t.f.}} \rightarrow \mathbf{D}(M)$ is surjective, and since $\mathbf{N}(M)^{\text{t.f.}} \subset \mathbf{N}(M)^{\text{sat}}$, the result follows. \square

Lemma 3.24. *The action of Γ on $\mathbf{N}(M)^{\text{sat}}/\pi$ is trivial, and the cokernel of $\varphi^* \mathbf{N}(M)^{\text{sat}} \rightarrow \mathbf{N}(M)^{\text{sat}}$ is annihilated by q^h .*

Proof. The kernel of $\mathbf{N}(M)^{\text{sat}} \rightarrow \mathbf{N}(M)^{\text{total}}[1/p]/\pi$ is $\pi \mathbf{N}(M)^{\text{sat}}$ (since π is invertible in $A \widehat{\otimes} \mathbf{A}_F$), so the natural map $\mathbf{N}(M)^{\text{sat}}/\pi \rightarrow \mathbf{N}(M)^{\text{total}}[1/p]/\pi$ is injective. Since the Γ -action on the right is trivial, so is the Γ -action on the left.

For the second assertion, we observe that φ is an isomorphism on $\mathbf{D}(M)$ (and on $\mathbf{D}(M)[1/p]$), and the cokernel of $\varphi^* \mathbf{N}(M)^{\text{total}}[1/p] \rightarrow \mathbf{N}(M)^{\text{total}}[1/p]$ is annihilated by q^h . \square

Lemma 3.25. *Suppose that A is flat over \mathbf{Z}_p . Then there are natural inclusions $\pi^h \mathbf{D}^+(M) \subset \mathbf{N}(M)^{\text{sat}} \subset \mathbf{D}^+(M)$, and natural inclusions $\pi^h ((A \widehat{\otimes} \mathbf{A}^+) \otimes_A M) \subset \mathbf{A}^+ \widehat{\otimes}_{\mathbf{A}_F^+} \mathbf{N}(M)^{\text{sat}} \subset (A \widehat{\otimes} \mathbf{A}^+) \otimes_A M$.*

Proof. The inclusion $\pi^h \mathbf{D}^+(M) \subset \mathbf{N}(M)^{\text{sat}}$ is clear, since $A \rightarrow \widetilde{A}$ is an injection (since it is an isomorphism after inverting p and A is assumed flat over \mathbf{Z}_p), and $\mathbf{D}^+(M)$ is p -torsion-free, and so

$$\pi^h \mathbf{D}^+(M) \subset \pi^h \mathbf{D}^+(M)[1/p] \cap \mathbf{D}(M) \subset \pi^h \mathbf{D}^+(\widetilde{A} \otimes_A M)[1/p] \cap \mathbf{D}(M) \subset \mathbf{N}(M)^{\text{sat}}.$$

where we use $\pi^h \mathbf{D}^+(\tilde{A} \otimes_A M) \subseteq \mathbf{N}(M)^{\text{total}}$, which follows from Lemma 3.19 by taking global sections. More precisely, formal GAGA implies that the injective map of coherent $\mathcal{O}_{\mathcal{Z}_M^{\leq h}} \hat{\otimes} \mathbf{A}^+$ -modules $\pi^h(\mathcal{O}_{\mathcal{Z}_M^{\leq h}} \hat{\otimes} \mathbf{A}^+) \otimes_A M \rightarrow (\mathcal{O}_{\mathcal{Z}_M^{\leq h}} \hat{\otimes} \mathbf{A}^+) \otimes_{\mathcal{O}_{\mathcal{Z}_M^{\leq h}} \hat{\otimes} \mathbf{A}_F^+} \widehat{\mathcal{N}}^{\leq h}$ over $\mathcal{Z}_M^{\leq h} \times_{\text{Spf } A} \text{Spf}(A \hat{\otimes} \mathbf{A}^+)$ arises from an injective map of coherent sheaves over $\mathcal{Z}_M^{\leq h} \times_{\text{Spec } A} \text{Spec}(A \hat{\otimes} \mathbf{A})$, since \mathfrak{m}_A -adic completion is fully faithful. Taking H_F -invariants and pushing forward by Θ_A yields the desired statement.

For the inclusion $\mathbf{N}(M)^{\text{sat}} \subset \mathbf{D}^+(M)$, we first recall from Lemma 3.19 that

$$\widehat{\mathcal{N}}^{\leq h}(M) \subset \mathbf{D}^+(\mathcal{O}_{\mathcal{Z}_M^{\leq h}} \otimes_A M) \subset (\mathcal{O}_{\mathcal{Z}_M^{\leq h}} \hat{\otimes} \mathbf{A}^+) \otimes_A M$$

so $\mathbf{N}(M)^{\text{total}}[1/p] \subset (M \otimes_A (\tilde{A} \hat{\otimes} \mathbf{A}^+))[1/p] = (M \otimes_A (A \hat{\otimes} \mathbf{A}^+))[1/p]$, and $\mathbf{N}(M)^{\text{total}}[1/p]$ is fixed by H_F . On the other hand,

$$\mathbf{D}(M) = ((A \hat{\otimes} \mathbf{A}) \otimes_A M)^{H_F} \subset (A \hat{\otimes} \mathbf{A}) \otimes_A M$$

Therefore, $\mathbf{N}(M)^{\text{sat}} \subset ((A \hat{\otimes} \mathbf{A}^+)[1/p] \cap (A \hat{\otimes} \mathbf{A})) \otimes_A M$, so it suffices to show that $A \hat{\otimes} \mathbf{A}^+ = (A \hat{\otimes} \mathbf{A}^+)[1/p] \cap (A \hat{\otimes} \mathbf{A})$ inside $(A \hat{\otimes} \mathbf{A})[1/p]$, or equivalently, that $p^k(A \hat{\otimes} \mathbf{A}^+) = (A \hat{\otimes} \mathbf{A}^+) \cap p^k(A \hat{\otimes} \mathbf{A})$ inside $A \hat{\otimes} \mathbf{A}$ for every integer $k \geq 0$. It is clear that $p^k(A \hat{\otimes} \mathbf{A}^+) \subset (A \hat{\otimes} \mathbf{A}^+) \cap p^k(A \hat{\otimes} \mathbf{A})$.

For the other direction, we first recall from §2 that the cokernel C of $\mathbf{A}^+ \hookrightarrow \mathbf{A}$ is \mathbf{Z}_p -flat. It follows that $(A/\mathfrak{m}_A^i) \otimes_{\mathbf{Z}_p} C$ is A/\mathfrak{m}_A^i -flat for all i . Now suppose that $x \in A \hat{\otimes}_{\mathbf{Z}_p} C$ satisfies $px = 0$. Then [Mat89, Theorem 7.6] implies that for all i , the image of x in $(A/\mathfrak{m}_A^i) \otimes_{\mathbf{Z}_p} C$ is of the form $\sum_{j=1}^{k_i} a_j y_j$, where $a_j \in A/\mathfrak{m}_A^i$, $y_j \in (A/\mathfrak{m}_A^i) \otimes_{\mathbf{Z}_p} C$, and $pa_j = 0$ for all j .

We claim that for any n , there is some $i \gg 0$ such that the annihilator $\text{ann}_{A/\mathfrak{m}_A^i}(p) \subset \mathfrak{m}_A^n/\mathfrak{m}_A^i$. Assume this for the moment. Then the image of x in $(A/\mathfrak{m}_A^i) \otimes_{\mathbf{Z}_p} C$ is of the form $\sum_{j=1}^{k_i} a_j y_j$, where $a_j \in \mathfrak{m}_A^n A/\mathfrak{m}_A^i$ and $y_j \in (A/\mathfrak{m}_A^i) \otimes_{\mathbf{Z}_p} C$. But then the image of x in $(A/\mathfrak{m}_A^n) \otimes_{\mathbf{Z}_p} C$ is 0. Since this is true for all n , $x = 0$.

To prove the claim, consider the sequence of ideals

$$\cdots \subset \overline{\text{ann}_{A/\mathfrak{m}_A^{i+1}}(p)} \subset \overline{\text{ann}_{A/\mathfrak{m}_A^i}(p)} \subset \cdots \subset A/\mathfrak{m}_A^n$$

over all $i \geq n$, where we consider the image of $\text{ann}_{A/\mathfrak{m}_A^i}(p)$ in A/\mathfrak{m}_A^n . This is a decreasing sequence of ideals and A/\mathfrak{m}_A^n is artinian, so it stabilizes to some ideal I when $i \geq i_0$ for some $i_0 \gg 0$. If $I = 0$, we are done. If not, we may consider the sequence of ideals

$$\cdots \subset \overline{\text{ann}_{A/\mathfrak{m}_A^{i+1}}(p)} \subset \overline{\text{ann}_{A/\mathfrak{m}_A^i}(p)} \subset \cdots \subset \tilde{I} \subset A/\mathfrak{m}_A^{n+1}$$

where $\tilde{I} \subset A/\mathfrak{m}_A^{n+1}$ denotes the pre-image of I in A/\mathfrak{m}_A^{n+1} . Repeating the previous argument, we obtain a non-zero ideal $I' \subset A/\mathfrak{m}_A^{n+1}$ such that the image of $\text{ann}_{A/\mathfrak{m}_A^i}(p)$ in A/\mathfrak{m}_A^{n+1} is I' for $i \geq i_1$, and $I' \equiv I \neq 0 \pmod{\mathfrak{m}_A^n}$. Repeating this argument for all n , we find that the annihilator $\text{ann}_A(p)$ is non-zero. But this contradicts the assumption that A is \mathbf{Z}_p -flat, so $I = 0$.

Now we turn to the second pair of inclusions. By Lemma 3.19

$$\mathbf{N}(M)^{\text{sat}}[1/p] \subset ((A \hat{\otimes} \mathbf{A}^+) \otimes_{A \hat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M)^{\text{sat}})[1/p] \subset ((A \hat{\otimes} \mathbf{A}^+) \otimes_A M)[1/p]$$

In addition,

$$\mathbf{N}(M)^{\text{sat}} \subset \mathbf{D}(M) \subset (A \hat{\otimes} \mathbf{A}) \otimes_{A \hat{\otimes} \mathbf{A}_F} \mathbf{D}(M) \cong (A \hat{\otimes} \mathbf{A}) \otimes_A M$$

Since M is free over A and we have just shown that $(A \hat{\otimes} \mathbf{A}^+) = (A \hat{\otimes} \mathbf{A}^+)[1/p] \cap (A \hat{\otimes} \mathbf{A})$, we see that $\mathbf{N}(M)^{\text{sat}} \subset (A \hat{\otimes} \mathbf{A}^+) \otimes_A M$, and there is a natural map

$$(A \hat{\otimes} \mathbf{A}^+) \otimes_{A \hat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M)^{\text{sat}} \rightarrow (A \hat{\otimes} \mathbf{A}^+) \otimes_A M$$

This map is injective after inverting p , and $(A \hat{\otimes} \mathbf{A}^+) \otimes_{A \hat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M)^{\text{sat}}$ has no p -torsion, so it is an inclusion.

It remains to show that $\pi^h(A \hat{\otimes} \mathbf{A}^+) \otimes_A M \subset (A \hat{\otimes} \mathbf{A}^+) \otimes_{A \hat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M)^{\text{sat}}$. By Lemma 3.19 we know that

$$\pi^h(A \hat{\otimes} \mathbf{A}^+) \otimes_A M \subset (A \hat{\otimes} \mathbf{A}^+) \otimes_{A \hat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M)^{\text{total}}$$

Since $\pi^h(A \hat{\otimes} \mathbf{A}^+) \otimes_A M$ has no p -torsion, it is contained in $(A \hat{\otimes} \mathbf{A}^+) \otimes_{A \hat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M)^{\text{t.f.}} \subset (A \hat{\otimes} \mathbf{A}^+) \otimes_{A \hat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M)^{\text{sat}}$ and we are done. \square

Corollary 3.26. $\mathbf{N}(M)^{\text{sat}}$ is a finite $A \hat{\otimes} \mathbf{A}_F^+$ -module, and $p^{\alpha(h)} \mathbf{N}(M)^{\text{sat}} \subset \mathbf{N}(M)^{\text{total}}$.

Proof. By Lemma 3.19, $\mathbf{D}^+(M \otimes_A \tilde{A})$ is finite over $A \hat{\otimes} \mathbf{A}_F^+$ (since $A \hat{\otimes} \mathbf{A}_F^+$ is noetherian). Since $A \rightarrow \tilde{A}$ is injective, $\mathbf{D}^+(M) \subset \mathbf{D}^+(M \otimes_A \tilde{A})$ and so $\mathbf{N}(M)^{\text{sat}}$ is finite over $A \hat{\otimes} \mathbf{A}_F^+$.

The proof of the second assertion is very similar to the proof of [Ber04, Proposition IV.1.3]. We note that we have a set of inclusions

$$\begin{array}{ccc} & \mathbf{N}(M)^{\text{total}} \hookrightarrow \mathbf{D}^+(\tilde{A} \otimes_A M) & \\ & \nearrow & \uparrow \\ \pi^h \mathbf{D}^+(M) & & \\ & \searrow & \\ & \mathbf{N}(M)^{\text{sat}} \hookrightarrow \mathbf{D}^+(M) & \end{array}$$

with the action of Γ stabilizing $\mathbf{N}(M)^{\text{total}}$ and $\mathbf{N}(M)^{\text{sat}}$ and acting trivially on $\mathbf{N}(M)^{\text{total}}/\pi$ and $\mathbf{N}(M)^{\text{sat}}/\pi$. In particular, $\pi^h \mathbf{N}(M)^{\text{sat}} \subset \mathbf{N}(M)^{\text{total}}$ (inside $\mathbf{D}^+(\tilde{A} \otimes_A M)$). We may then prove by induction on i that

$$\pi^{h-i} \prod_{j=0}^{i-1} (\chi(\gamma)^{h-j} - 1) \mathbf{N}(M)^{\text{sat}} \subset \mathbf{N}(M)^{\text{total}}$$

Taking $i = h$ and observing that $p^{\alpha(h)}$ generates the same ideal in \mathbf{Z}_p as $\prod_{j=0}^{h-1} (\chi(\gamma)^{h-j} - 1)$, we see that $p^{\alpha(h)} \mathbf{N}(M)^{\text{sat}} \subset \mathbf{N}(M)^{\text{total}}$, as desired. \square

We can also study the behavior of $\mathbf{N}(M)^{\text{sat}}$ under change of coefficients. As the formation of $\mathcal{L}_M^{\leq h}$ is clearly functorial in the coefficients, for any homomorphism of complete local noetherian \mathcal{O} -algebras $A \rightarrow A'$, we have a natural base change morphism $A' \hat{\otimes}_A \mathbf{N}(M)^{\text{sat}} \rightarrow \mathbf{N}(M_{A'})^{\text{sat}}$. However, it need not be an isomorphism.

Lemma 3.27. *Let $A \rightarrow A'$ be a homomorphism of complete local noetherian flat \mathbf{Z}_p -algebras with finite residue fields which are integrally closed in $A[1/p]$ and $A'[1/p]$, respectively. Then the image of the natural map $(A' \hat{\otimes}_A \mathbf{A}_F^+) \otimes_{A \hat{\otimes} \mathbf{A}_F^+} \mathbf{N}^{\text{sat}}(M_A) \rightarrow \mathbf{N}^{\text{sat}}(M_{A'})$ contains $\pi^h \mathbf{D}^+(M_{A'})$.*

Proof. By Proposition 3.21, the image of $(A' \hat{\otimes}_A \mathbf{A}_F^+) \otimes_{A \hat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M)^{\text{total}}$ contains the image of $\pi^h \mathbf{D}^+(M_{A'})$, and since $\pi^h \mathbf{D}^+(M_{A'})$ has no p -torsion, the image of $(A' \hat{\otimes}_A \mathbf{A}_F^+) \otimes_{A \hat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M)^{\text{t.f.}}$ contains the image of $\pi^h \mathbf{D}^+(M_{A'})$. Since $\mathbf{N}(M)^{\text{t.f.}} \subset \mathbf{N}(M)^{\text{sat}}$, we are done. \square

Corollary 3.28. *Let $A \rightarrow A'$ be a local homomorphism of rings satisfying the above hypotheses. Then there is a natural base change morphism $(A' \hat{\otimes}_A \mathbf{A}_F^+) \otimes_{A \hat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M)^{\text{sat}} \rightarrow \mathbf{N}(M_{A'})^{\text{sat}}$, and its cokernel is annihilated by $p^{\alpha(h)}$.*

Proof. This follows from Lemma 3.27 by the same argument as in the proof of Lemma 3.26. \square

Suppose in addition that $A = \varprojlim_i A/J^i$, for some ideal $J \subsetneq A$ such that A/J^i is a finite flat \mathbf{Z}_p -module. There is a natural map $\mathbf{N}(M)^{\text{sat}} \rightarrow \varprojlim_i \mathbf{N}(M_{A/J^i})^{\text{sat}}$.

Proposition 3.29. *The natural map $\mathbf{N}(M)^{\text{sat}} \rightarrow \varprojlim_i \mathbf{N}(M_{A/J^i})^{\text{sat}}$ is an isomorphism.*

Proof. Consider an element $(n_i) \in \varprojlim_i \mathbf{N}(M_{A/J^i})^{\text{sat}}$. Then Corollary 3.26 implies that $(p^{\alpha(h)} n_i) \in \varprojlim_i \mathbf{N}(M_{A/J^i}) = \mathbf{N}(M)$, so (n_i) comes from an element $n \in \mathbf{N}(M)[1/p]$. But (n_i) also comes from an element of $\mathbf{D}(M)$, because $\mathbf{D}(M) = \varprojlim_i \mathbf{D}(M_{A/J^i})$, so $n \in \mathbf{N}(M)^{\text{sat}}$. \square

Let $\mathbf{N}(M)^{p\text{-tors}}$ denote the p -power torsion in $\mathbf{N}(M)^{\text{total}}$.

Lemma 3.30. *If $A = \varprojlim_i A/J^i$, for some ideal $J \subsetneq A$ such that A/J^i is \mathbf{Z}_p -flat, $\mathbf{N}(M)^{p\text{-tors}} = \varprojlim_i \mathbf{N}(M_{A/J^i})^{p\text{-tors}}$.*

Proof. Clearly if $n \in \mathbf{N}(M)^{\text{total}}$ is p -power torsion, so is its image in $\mathbf{N}(M_{A/J^i})^{\text{total}}$ for all i .

On the other hand, the natural map $\mathbf{N}(M)^{\text{total}}[1/p] \otimes_{A[1/p]} (A/J^i)[1/p] \rightarrow \mathbf{N}(M_{A/J^i})^{\text{total}}[1/p]$ is an isomorphism for all i , so to prove the converse, we show that $\mathbf{N}(M)^{\text{total}} \cap \bigcap_i (J^i[1/p]) \mathbf{N}(M)$ is p -power torsion. Since A/J is assumed \mathbf{Z}_p -flat, $J[1/p] \subset A[1/p]$ is a proper ideal. Then by a theorem

of Krull [Mat89, Theorem 8.9] there is some $a \in A[1/p]$ which is 1 modulo $J[1/p]$ and which annihilates $\cap_i (J^i[1/p])\mathbf{N}(M)[1/p]$. Again because A/J is assumed \mathbf{Z}_p -flat, $J[1/p] \cap A = J$, so there is some power of p such that $a' := p^n a$ is an element of A which is congruent to p^n modulo J and annihilates $\ker(\mathbf{N}(M)^{\text{total}} \rightarrow \mathbf{N}(M_{A/J^i})^{\text{total}}[1/p])$ for all i . Then a' is congruent to p^n modulo the maximal ideal of A , so multiplying a' by a 1-unit of A , we see that p^n annihilates $\ker(\mathbf{N}(M)^{\text{total}} \rightarrow \mathbf{N}(M_{A/J^i})^{\text{total}}[1/p])$ for all i and we are done. \square

From now on, we assume that R is \mathbf{Z}_p -flat and integrally closed in $R[1/p]$. We refer to $\mathbf{N}(M)^{\text{sat}}$ as the Wach module of M and we drop the superscript, writing $\mathbf{N}(M)$ instead. For the convenience of the reader, we collect some useful properties of $\mathbf{N}(M)$:

thm:wach

Theorem 3.31. *Let M be a crystalline family of G_F -representations over R (of rank d) with Hodge–Tate weights in $[-h, 0]$. Then there exists a canonical finitely generated, φ - and Γ -stable $R \widehat{\otimes} \mathbf{A}_F^+$ -submodule*

$$\mathbf{N}(M) \subseteq \mathbf{D}(M)$$

satisfying the following properties:

- (1) $(R \widehat{\otimes} \mathbf{A}_F) \otimes_{R \widehat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M) \rightarrow \mathbf{D}(M)$, i.e., N generates $\mathbf{D}(M)$ as a $R \widehat{\otimes} \mathbf{A}_F$ -module,
- (2) $\mathbf{N}(M)[1/p]$ is a finitely generated projective $(R \widehat{\otimes} \mathbf{A}_F^+[1/p])$ -module (of rank d),
- (3) $\mathbf{N}(M)$ is p -saturated in $\mathbf{D}(M)$, i.e., $\mathbf{N}(M)[1/p] \cap \mathbf{D}(M) = \mathbf{N}$,
- (4) Γ acts trivially on $\mathbf{N}(M)/\pi\mathbf{N}(M)$,
- (5) q^h annihilates the cokernel of $\varphi^*\mathbf{N}(M) \rightarrow \mathbf{N}(M)$,
- (6) $\pi^h\mathbf{D}^+(M) \subseteq \mathbf{N}(M) \subseteq \mathbf{D}^+(M)$.

Moreover, for any local homomorphism $R \rightarrow R'$ of complete local noetherian flat \mathbf{Z}_p -algebras with finite residue fields $R \rightarrow R'$, integrally closed in $R[1/p]$ and $R'[1/p]$, respectively, $p^{\alpha(h)}$ annihilates the cokernel of $(R' \widehat{\otimes} \mathbf{A}_F^+) \otimes_{R \widehat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M) \rightarrow \mathbf{N}(M_{A'})$.

sec:fibre

4. GENERIC FIBRE

4.1. Families of Wach modules and (φ, Γ) -modules. Given a family of positive crystalline representations M of G_F over R of rank d , we get a family M^{rig} of Galois representations over the Fréchet–Stein algebra R^{rig} given by taking the rigid analytic generic fiber of R . We therefore obtain a family of (φ, Γ) -modules $\mathbf{D}_{\text{rig}}^\dagger(M^{\text{rig}})$ over R^{rig} . More precisely, we exhaust the quasi-Stein space $\text{Spf}(R)^{\text{rig}}$ by a rising sequence of affinoids $\{U_i = \text{Sp}(A_i)\}$ and we note that for each i there is some $s > 0$ such that over each U_i , $\mathbf{D}_{\text{rig}}^\dagger(M^{\text{rig}}|_{U_i})$ is generated by $\mathbf{D}_F^{\dagger, s}(M^{\text{rig}}|_{U_i})$. Letting $C^{(0, s]}$ denote the half-open annulus $\{0 < v_p(\pi) \leq s\}$ over F , we obtain a coherent sheaf on the quasi-Stein space $\text{Spf}(R)^{\text{rig}} \times C^{(0, s]}$. Taking global sections and extending scalars from $R^{\text{rig}} \widehat{\otimes} \mathbf{B}_{\text{rig}, F}^{\dagger, s}$ to $R^{\text{rig}} \widehat{\otimes} \mathbf{B}_{\text{rig}, F}^\dagger$ yields $\mathbf{D}_{\text{rig}}^\dagger(M^{\text{rig}})$.

Similarly, to the Wach module $\mathbf{N}(M)$ of M , we associate a finite module

$$\mathbf{N}(M)^{\text{rig}} := (\mathbf{B}_{\text{rig}, F}^+ \widehat{\otimes} R^{\text{rig}}) \otimes_{(R \widehat{\otimes} \mathbf{A}_F^+)} \mathbf{N}(M)$$

over $\mathbf{B}_{\text{rig}, F}^+ \widehat{\otimes} R^{\text{rig}}$. Indeed, since $\mathbf{N}(M)$ is a finite $R \widehat{\otimes} \mathbf{A}_F^+$ -module the functor $(-)^{\text{rig}}$ associates to it a coherent sheaf on the rigid space $X = \text{Spf}(R \widehat{\otimes} \mathbf{A}_F^+)^{\text{rig}}$ with $\mathcal{O}_X(X) = \mathbf{B}_{\text{rig}, F}^+ \widehat{\otimes} R^{\text{rig}}$; here we identify this sheaf $\mathbf{N}(M)^{\text{rig}}$ with its global sections, see Lemma A.6.

We claim first that if M is positive, then $\mathbf{N}(M)^{\text{rig}} \subset \mathbf{D}_{\text{rig}}^\dagger(M^{\text{rig}})$. Recall that the Wach module $\mathbf{N}(M)$ induces a finite $\mathbf{B}_{\text{rig}, F}^+ \widehat{\otimes} R^{\text{rig}}$ -module $\mathbf{N}(M)^{\text{rig}}$ by extension of scalars (we implicitly use the Mittag–Leffler condition for modules over Fréchet–Stein algebras to commute the tensor products and inverse limit). Equivalently, it induces a vector bundle on $\text{Spf}(\mathbf{A}_{\mathbf{Q}_p}^+ \widehat{\otimes} R)^{\text{rig}}$.

Let $\cup_n \text{Sp}(A_n)$ be a rising admissible covering of $\text{Spf}(R)^{\text{rig}}$, so that $R^{\text{rig}} = \cap_m A_m$, and let A_m° denote the ring of power-bounded elements of A_m . Then for any m and any s , the extension of scalars $\mathbf{N}(M)^{\text{rig}} \otimes_{\mathbf{B}_{\text{rig}, F}^+ \widehat{\otimes} R^{\text{rig}}} (\mathbf{B}_{\text{rig}, F}^{\dagger, s} \widehat{\otimes} A_m)$ remains finite projective. Moreover, since $\mathbf{N}(M) \subset (\mathbf{A}^+ \widehat{\otimes} R) \otimes_R M$, we have a natural map

$$\mathbf{N}(M)^{\text{rig}} \otimes_{\mathbf{B}_{\text{rig}, F}^+ \widehat{\otimes} R^{\text{rig}}} (\mathbf{B}_{\text{rig}, F}^{\dagger, s} \widehat{\otimes} A_m) \rightarrow (\widetilde{\mathbf{B}}^{\dagger, s} \widehat{\otimes} A_m) \otimes_{A_m} (A_m \otimes_R M)$$

On the other hand, the family of Galois representations $M_{A_m} := A_m \otimes_R M$ over A_m induces a family of (φ, Γ) -modules $\mathbf{D}_{\text{rig}}^{\dagger, s}(M_{A_m})$ over $\mathbf{B}_{\text{rig}, \mathbf{Q}_p}^{\dagger, s} \widehat{\otimes} A_m$ for $s \gg 0$. This family of (φ, Γ) -modules is finite projective,

and $\mathbf{D}_{\text{rig}}^{\dagger,s}(M_{A_m})$ is naturally a sub- $\mathbf{B}_{\text{rig},F}^{\dagger,s} \widehat{\otimes} A_m$ -module of $(\widetilde{\mathbf{B}}^{\dagger,s} \widehat{\otimes} A_m) \otimes_{A_m} (A_m \otimes_R M)$. Equivalently, $\mathbf{D}_{\text{rig}}^{\dagger,s}(M_{A_m})$ is a vector bundle on the product X_s of a half-open annulus with $\text{Sp}(A_m)$.

We may therefore consider the sub- $\mathbf{B}_{\text{rig},F}^{\dagger,s} \widehat{\otimes} A_m$ -module $\mathbf{D}_{\text{rig}}^{\dagger,s}(M_{A_m}) + \left(\mathbf{N}(M)^{\text{rig}} \otimes_{\mathbf{B}_{\text{rig},F}^{\dagger,s} \widehat{\otimes} R^{\text{rig}}} (\mathbf{B}_{\text{rig},F}^{\dagger,s} \widehat{\otimes} A_m) \right)$ of $(\widetilde{\mathbf{B}}^{\dagger,s} \widehat{\otimes} A_m) \otimes_{A_m} (A_m \otimes_R M)$ generated by the images of $\mathbf{N}(M)^{\text{rig}}$ and $\mathbf{D}_{\text{rig}}^{\dagger,s}(A_m \otimes_R M)$. For ease of notation, we write it as $\mathbf{D}_{\text{rig}}^{\dagger,s}(M_{A_m}) + \mathbf{N}(M)_{A_m}^{\text{rig}}$. Then $\mathbf{D}_{\text{rig}}^{\dagger,s}(M_{A_m}) + \mathbf{N}(M)_{A_m}^{\text{rig}}$ remains finite over $\mathbf{B}_{\text{rig},\mathbf{Q}_p}^{\dagger,s} \widehat{\otimes} A_m$, and we have an exact sequence of coherent sheaves on the quasi-Stein space X_s

$$\mathbf{D}_{\text{rig}}^{\dagger,s}(M_{A_m}) \rightarrow \mathbf{D}_{\text{rig}}^{\dagger,s}(M_{A_m}) + \mathbf{N}(M)_{A_m}^{\text{rig}} \rightarrow Q \rightarrow 0$$

for some quotient sheaf Q . But the formations of $\mathbf{D}_{\text{rig}}^{\dagger,s}(M_{A_m})$ and $\mathbf{N}(M)^{\text{rig}}$ commute with specialization on A_m , and in the classical case the Wach module is contained in the (φ, Γ) -module. Thus, Q vanishes at every point of X_s , so is trivial, and $\mathbf{D}_{\text{rig}}^{\dagger,s}(M_{A_m}) \rightarrow \mathbf{D}_{\text{rig}}^{\dagger,s}(M_{A_m}) + \mathbf{N}(M)_{A_m}^{\text{rig}}$ is surjective.

4.2. Comparisons between $\mathbf{N}(M)$ and $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$. The paper [Ber02] explains how to relate (φ, Γ) -modules to the functors $\mathbf{D}_{\text{cris}}(-)$ and $\mathbf{D}_{\text{st}}(-)$, and [Ber04] refines this for positive crystalline representations of G_F . Indeed, if V is a finite-dimensional \mathbf{Q}_p -linear representation of G_K , we have

$$\mathbf{D}_{\text{cris}}(V) = \mathbf{D}_{\text{rig}}^{\dagger}(V)[1/t]^{\Gamma}$$

by [Ber02, Théorème 3.6].

If $T \subset V$ is a lattice in a finite-dimensional \mathbf{Q}_p -linear positive crystalline representation of G_F , then the Wach module $\mathbf{N}(V) = \mathbf{N}(T)[1/p]$ is contained in the (φ, Γ) -module $\mathbf{D}_{\text{rig}}^{\dagger}(V)$, and it generates it over $\mathbf{B}_{\text{rig},F}^{\dagger}$. Further:

Proposition 4.1 ([Ber04, Proposition II.2.1], [Ber04, Proposition III.2.1]). *Let $0 \leq r_1 \leq \dots \leq r_d$ be the opposites of the Hodge–Tate weights of V . Then*

- (1) $\mathbf{D}_{\text{cris}}(V) \subset \mathbf{B}_{\text{rig},F}^+ \otimes_{\mathbf{B}_F^+} \mathbf{N}(V)$, and
- (2) the natural map $\mathbf{B}_{\text{rig},F}^+ \otimes_F \mathbf{D}_{\text{cris}}(V) \rightarrow \mathbf{B}_{\text{rig},F}^+ \otimes_{\mathbf{B}^+} \mathbf{N}(V)$ is an injection and its cokernel is isomorphic to $\bigoplus_{i=1}^d \mathbf{B}_{\text{rig},F}^+ / (\lambda_i)$, where $\lambda_i = (t/\pi)^{r_i}$.

This implies that $\mathbf{D}_{\text{cris}}(V) = \left(\mathbf{B}_{\text{rig},F}^+ \otimes_{\mathbf{B}_F^+} \mathbf{N}(V) \right)^{\Gamma}$. But we have another relation:

Theorem 4.2 ([Ber04, Théorème]). *If V is a positive crystalline representation of G_F and we equip the Wach module $\mathbf{N}(V)$ with the filtration*

$$\text{Fil}^i \mathbf{N}(V) := \{x \in \mathbf{N}(V) \mid \varphi(x) \in q^i \mathbf{N}(V)\}$$

then the natural map $\mathbf{D}_{\text{cris}}(V) \rightarrow \mathbf{N}(V)/\pi \mathbf{N}(V)$ is an isomorphism of filtered φ -modules.

In particular, this implies that $\mathbf{D}_{\text{cris}}(T) := \mathbf{N}(T)/\pi \mathbf{N}(T)$ defines a \mathcal{O}_F -lattice in the F -vector space $\mathbf{D}_{\text{cris}}(V)$.

We wish to extend these results to formal families of positive crystalline representations. To precisely define $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$ we must again exhaust $\text{Spf}(R)^{\text{rig}}$ by affinoid subdomains $\{U_i\}$, consider the finite projective modules $\mathbf{D}_{\text{cris}}(M^{\text{rig}}|_{U_i})$ over each U_i , and take the limit to obtain a finite projective filtered φ -module over R^{rig} . We also obtain an explicit description $\mathbf{D}_{\text{cris}}(M^{\text{rig}}) = ((R^{\text{rig}} \widehat{\otimes} \mathbf{B}_{\text{max}}) \otimes_{R^{\text{rig}}} M^{\text{rig}})^{G_F}$. Then [Bel15, Theorem 4.2.9] shows that $\mathbf{D}_{\text{cris}}(M^{\text{rig}}) = \mathbf{D}_{\text{rig}}^{\dagger}(M^{\text{rig}})^{\Gamma}$.

Lemma 4.3. *If M is positive crystalline, then the image of $\mathbf{N}(M)^{\text{rig}}$ inside $\mathbf{D}_{\text{rig}}^{\dagger}(M^{\text{rig}})$ generates it over $\mathbf{B}_{\text{rig},F}^{\dagger} \widehat{\otimes} R^{\text{rig}}$.*

Proof. Let $U = \text{Sp}(A)$ be an affinoid subdomain of $\text{Spf}(R)^{\text{rig}}$. Then there is some $s \gg 0$ such that $\mathbf{D}_{\text{rig}}^{\dagger}(M^{\text{rig}}|_U)$ is generated by $\mathbf{D}_{\text{rig}}^{\dagger,s}(M_U^{\text{rig}})$. After extending scalars on $\mathbf{N}(M)^{\text{rig}}$ from $\mathbf{B}_{\text{rig},F}^{\dagger} \widehat{\otimes} R^{\text{rig}}$ to $\mathbf{B}_{\text{rig},F}^{\dagger,s} \widehat{\otimes} R^{\text{rig}}$, we have a homomorphism of coherent sheaves of modules on the quasi-Stein space attached to $\mathbf{B}_{\text{rig},F}^{\dagger,s} \widehat{\otimes} A$. Its cokernel is likewise a coherent sheaf. But by specializing at points of U and appealing to the classical case, we conclude that the cokernel must vanish. \square

In fact, with this in place, projectivity of $\mathbf{D}_{\text{rig}}^{\dagger,s}(M^{\text{rig}})$ lets us use specialization to conclude that the kernel vanishes at every point, so we actually have an isomorphism $\mathbf{N}(M)^{\text{rig}} \otimes_{\mathbf{B}_{\text{rig},F}^+ \widehat{\otimes} R^{\text{rig}}} (\mathbf{B}_{\text{rig},F}^{\dagger,s} \widehat{\otimes} R^{\text{rig}}) \rightarrow \mathbf{D}_{\text{rig}}^{\dagger,s}(M^{\text{rig}})$.

Note that $\mathbf{D}_{\text{rig}}^{\dagger}(M^{\text{rig}})$ is projective as a $\mathbf{B}_{\text{rig},\mathbf{Q}_p}^+ \widehat{\otimes} R^{\text{rig}}$ -module; we do not know whether it is R^{rig} -locally free. However, it follows from the construction of families of (φ, Γ) -modules that there is some finite extension L/F such that for any affinoid subdomain $\text{Sp}(A) = U \subset \text{Spf}(R)^{\text{rig}}$, $\mathbf{D}_{\text{rig}}^{\dagger,s}(M^{\text{rig}}|_{U, G_L})$ is a free $\mathbf{B}_{\text{rig},L}^{\dagger,s} \widehat{\otimes} A$ -module of rank n .

Proposition 4.4. *If M is positive crystalline, $\mathbf{D}_{\text{cris}}(M^{\text{rig}}) \subset \mathbf{N}(M)^{\text{rig}}$ inside $\mathbf{D}_{\text{rig}}^{\dagger}(M^{\text{rig}})$.*

Proof. Since $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$ is a finite projective module over $R^{\text{rig}} \otimes_{\mathbf{Q}_p} F$ and $\mathbf{N}(M)[1/p]$ can be trivialized locally on $\text{Spec } R[1/p]$ by Lemma 3.18, we can find an affinoid covering $\{U_m = \text{Sp}(A_m)\}_{m \in I}$ of $\text{Spf}(R)^{\text{rig}}$ such that $\mathbf{D}_{\text{cris}}(M^{\text{rig}}|_{U_m})$ and $\mathbf{N}(M)^{\text{rig}}|_{U_m}$ are free for each i . We may also assume that $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$ is free and $\mathbf{D}_{\text{rig}}^{\dagger}(M^{\text{rig}})$ is generated by $\mathbf{D}_{\text{rig}}^{\dagger,s}(M^{\text{rig}})$ for some $s \gg 0$.

Now we choose a basis $\{\vec{e}_1, \dots, \vec{e}_r\}$ of $\mathbf{N}(M)^{\text{rig}}|_{U_m}$ and a basis $\{\vec{f}_1, \dots, \vec{f}_r\}$ of $\mathbf{D}_{\text{cris}}(M^{\text{rig}}|_{U_m})$. Let $G = (g_{ij}) \in \text{Mat}_{r \times r}(\mathbf{B}_{\text{rig},F}^{\dagger,s} \widehat{\otimes} A_m)$ be a matrix carrying $\{\vec{e}_1, \dots, \vec{e}_r\}$ to $\{\vec{f}_1, \dots, \vec{f}_r\}$. We wish to show that G actually has entries in $\mathbf{B}_{\text{rig},F}^+ \widehat{\otimes} A_m$.

A priori, $g_{ij}(\pi)$ is a power series with coefficients in A_m converging for $p^{-1/s} \leq |\pi| < 1$. We write $g_{ij}(\pi) = \sum_{n \in \mathbf{Z}} a_n \pi^n$ and equip A_m with the sup norm. But by the proof of [Ber04, II.2.1], if we specialize G at any point $x \in \text{Sp}(A_m)$, then $g_{ij}(\pi)(x) \in \mathbf{B}_{\text{rig},F}^+$. Since $|a_n| \leq |a_n(x)|$ for all x , it follows that g_{ij} actually converges for $|\pi| < 1$. \square

Now that we have an inclusion $\mathbf{D}_{\text{cris}}(M^{\text{rig}}) \subset \mathbf{N}(M)^{\text{rig}}$, we wish to understand the cokernel of the induced map $(\mathbf{B}_{\text{rig},F}^+ \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) \rightarrow \mathbf{N}(M)^{\text{rig}}$. When $R = \mathbf{Z}_p$, Berger showed [Ber04, §III] that the quotient is isomorphic to $\bigoplus_{i=1}^n \mathbf{B}_{\text{rig},F}^+ / \lambda_i$, where $\lambda_i = (t/\pi)^{r_i}$. However, as such a decomposition is non-canonical, there is no reason to expect it to vary well in families.

However, we can prove the following weaker results:

Proposition 4.5. *Let M be a family of positive crystalline representations of G_F over R . Then:*

- (1) *Let Q denote the cokernel of the map $\text{inc}_M : (\mathbf{B}_{\text{rig},F}^+ \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) \rightarrow \mathbf{N}(M)^{\text{rig}}$. Then Q is annihilated by λ_d . In particular, Q is t -torsion.*
- (2) *The map inc_M is injective.*
- (3) *The map inc_M induces an isomorphism of line bundles*

$$\left(\frac{\pi}{t}\right)^{r_1 + \dots + r_d} \cdot \det(\text{inc}_M) : (\mathbf{B}_{\text{rig},F}^+ \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \text{Det}_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) \xrightarrow{\sim} \text{Det}_{R^{\text{rig}}} \widehat{\otimes}_{\mathbf{B}_{\text{rig},F}^+} \mathbf{N}(M)^{\text{rig}}$$

Proof. For the first part, we consider the corresponding morphism of coherent sheaves on the rigid analytic space corresponding to $\mathbf{B}_{\text{rig},F}^+ [1/t] \widehat{\otimes} R^{\text{rig}}$. As the formations of $\mathbf{N}(M)^{\text{rig}}$ and $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$ are compatible with specialization on R^{rig} , Berger's result implies that $\lambda_d Q = 0$.

We turn to the second part. Since $(\mathbf{B}_{\text{rig},F}^+ \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})$ and $\mathbf{N}(M)^{\text{rig}}$ are finite projective modules of the same rank, the map is an isomorphism after inverting t . Thus, the kernel is also t -torsion. But $(\mathbf{B}_{\text{rig},F}^+ \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})$ is a projective module over $\mathbf{B}_{\text{rig},F}^+ \widehat{\otimes} R^{\text{rig}}$ so has no t -torsion.

For the last part, it suffices to consider the case where M is rank 1. But then compatibility of $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$ and $\mathbf{N}(M)^{\text{rig}}$ with specialization on R^{rig} , combined with Berger's result, yields the desired result. \square

Corollary 4.6. *Let M be a family of positive crystalline representations of G_F over R . Then*

$$\varphi^{-\langle r_1 + \dots + r_d \rangle} \cdot \det(\varphi^* \mathbf{N}(M)[1/p] \rightarrow \mathbf{N}(M)[1/p])$$

is an isomorphism of line bundles over $(R \widehat{\otimes} \mathbf{A}_F^+)[1/p]$.

Proof. Consider the commutative diagram

$$\begin{array}{ccc} (R^{\text{rig}} \widehat{\otimes} \mathbf{B}_{\text{rig},F}^+ \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})) & \longrightarrow & \mathbf{N}(M)^{\text{rig}} \\ \uparrow & & \uparrow \\ \varphi^* \left((R^{\text{rig}} \widehat{\otimes} \mathbf{B}_{\text{rig},F}^+ \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})) \right) & \longrightarrow & \varphi^* \mathbf{N}(M)^{\text{rig}} \end{array}$$

It is clear that $\varphi^* \left((R^{\text{rig}} \widehat{\otimes} \mathbf{B}_{\text{rig},F}^+ \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})) \right) \cong (R^{\text{rig}} \widehat{\otimes} \mathbf{B}_{\text{rig},F}^+ \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}))$, because φ acts bijectively on $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$. Since $(\varphi(\pi)/t)^{r_1+\dots+r_d} \cdot \det \left(\varphi^* \left((\mathbf{B}_{\text{rig},F}^+ \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) \right) \rightarrow \varphi^* \mathbf{N}(M)^{\text{rig}} \right)$ is an isomorphism of line bundles over $R^{\text{rig}} \widehat{\otimes} \mathbf{B}_{\text{rig},F}^+$, we see that $q^{-(r_1+\dots+r_d)} \cdot \det(\varphi^* \mathbf{N}(M)[1/p] \rightarrow \mathbf{N}(M)[1/p])$ becomes an isomorphism after extending scalars to $R^{\text{rig}} \widehat{\otimes} \mathbf{B}_{\text{rig},F}^+$. But since the stalks at closed points of $\text{Spec}(R \widehat{\otimes} \mathbf{A}_F^+[1/p])$ are noetherian and their completions agree with the completed stalks of $(\text{Spf}(R \widehat{\otimes} \mathbf{A}_F^+))^{\text{rig}}$ by [dJ95, Lemma 7.1.9], this implies the desired result. \square

Corollary 4.7. *Let M be a family of positive crystalline representations of G_F over R . Then the inclusion*

$$\left((R \widehat{\otimes} \mathbf{A}^+) \otimes_{R \widehat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M) \right) \subset \left((R \widehat{\otimes} \mathbf{A}^+) \otimes_R M \right)$$

induces an isomorphism of line bundles over $(R \widehat{\otimes} \mathbf{A}^+)[1/p]$

$$q^{-(r_1+\dots+r_d)} \cdot \det \left(((R \widehat{\otimes} \mathbf{A}^+) \otimes_{R \widehat{\otimes} \mathbf{A}_F^+} \mathbf{N}(M))[1/p] \rightarrow ((R \widehat{\otimes} \mathbf{A}^+) \otimes_R M)[1/p] \right)$$

Proof. We use Lemma 3.18 to find an affine cover $\{U_i = \text{Spec } R[1/p, 1/f_i]\}$ of $\text{Spec } R[1/p]$ such that $M[1/p]|_{U_i}$ and $\mathbf{N}(M)[1/p]|_{U_i}$ are free. Choose bases, and let $X_i \in \text{Mat}_d((R \widehat{\otimes} \mathbf{A})[1/p, 1/f_i])$ be a matrix whose columns are the basis elements of $\mathbf{N}(M)[1/p]|_{U_i}$ with respect to the basis of $M[1/p]|_{U_i}$. Let P_i be the matrix of Frobenius with respect to the chosen basis of $\mathbf{N}(M)[1/p]|_{U_i}$. Then $\varphi(X_i) = X_i P_i$ because Frobenius acts trivially on M . This implies that $\varphi(\det(X_i))/\det(X_i) = \det(P_i) = q^{r_1+\dots+r_d}$ up to a unit of $(R \widehat{\otimes} \mathbf{A}_F^+[1/p, 1/f_i])$, by Corollary 4.6. But $\pi^h \left((R \widehat{\otimes} \mathbf{A})[1/p, 1/f_i] \otimes_{R[1/p, 1/f_i]} M[1/p]|_{U_i} \subset ((R \widehat{\otimes} \mathbf{A})[1/p, 1/f_i]) \otimes_{(R \widehat{\otimes} \mathbf{A}_F^+[1/p, 1/f_i])} \mathbf{N}(M)[1/p]|_{U_i} \right)$ by Lemma 3.25, so $(\det(X_i))$ is trivial after inverting π . This implies that $\det(X_i) = \pi^{r_1+\dots+r_d}$ up to a unit of $(R \widehat{\otimes} \mathbf{A}^+)[1/p, 1/f_i]$. \square

We can also use Proposition 4.5 to relate $\mathbf{N}(M)^{\text{rig}}$ and $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$ when M^{rig} is a crystalline family with non-negative Hodge–Tate weights.

Corollary 4.8. *Suppose that M is a crystalline family with non-negative Hodge–Tate weights. Then $\varphi^* \mathbf{N}(M)^{\text{rig}} \subset (\mathbf{B}_{\text{rig},\mathbf{Q}_p}^+ \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})$. Since this is a map of finite modules over a Fréchet–Stein algebra, continuity is automatic.*

Proof. Suppose that the Hodge–Tate weights of M^{rig} lie in the interval $[0, h]$. Then by definition $\mathbf{N}(M)^{\text{rig}} = \pi^{-h} \mathbf{N}(M(-h))^{\text{rig}}$, and $M^{\text{rig}}(-h)$ is a positive crystalline family, so

$$\varphi^* \mathbf{N}(M)^{\text{rig}} = \varphi(\pi)^{-h} \varphi^* \mathbf{N}(M(-h))^{\text{rig}} \subset t^{-h} \varphi^* \left((\mathbf{B}_{\text{rig},\mathbf{Q}_p}^+ \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}(-h)) \right)$$

by Proposition 4.5 (since $\lambda_i = (t/\pi)^{r_i}$ and $r_i \leq h$). But $\varphi^* \mathbf{D}_{\text{cris}}(M^{\text{rig}}) = \mathbf{D}_{\text{cris}}(M^{\text{rig}})$ and $t^{-h} \mathbf{D}_{\text{cris}}(M^{\text{rig}}(-h)) = \mathbf{D}_{\text{cris}}(M^{\text{rig}})$, so we are done. \square

We now turn to the other comparison between Wach modules and \mathbf{D}_{cris} , namely [Ber04, Théorème III.4.4], which states that if V is a positive crystalline representation of $G_{\mathbf{Q}_p}$, then $\mathbf{N}(V)/\pi \mathbf{N}(V) \cong \mathbf{D}_{\text{cris}}(V)$.

Theorem 4.9. *Let M be a family of positive crystalline G_F -representations over R . Then the composition of the natural maps*

$$\mathbf{D}_{\text{cris}}(M^{\text{rig}}) \hookrightarrow \mathbf{N}(M)^{\text{rig}} \twoheadrightarrow \mathbf{N}(M)^{\text{rig}}/\pi \mathbf{N}(M)^{\text{rig}}$$

is an isomorphism.

Proof. The map of interest is a homomorphism of finite projective modules of the same rank over $R^{\text{rig}} \otimes_{\mathbf{Q}_p} F$. But the formations of $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$ and $\mathbf{N}(M)^{\text{rig}}$ commute with base change on R^{rig} and the map becomes an isomorphism after specialization, so we are done. \square

Since $\mathbf{N}(M)$ provides a natural integral structure on $\mathbf{N}(M)^{\text{rig}}$, we obtain an integral version of \mathbf{D}_{cris} . Namely, we define $\mathbf{D}_{\text{cris}}(M) := \mathbf{N}(M)/\pi\mathbf{N}(M)$, which is a finitely $R \otimes_{\mathbf{Z}_p} \mathcal{O}_F$ -module. Note, however, that we do not know whether $\mathbf{D}_{\text{cris}}(M)$ is projective over R (though $\mathbf{D}_{\text{cris}}(M)[1/p]$ is projective over $R[1/p]$).

Corollary 4.10. *Let M be a family of positive crystalline G_F -representations over R . Then*

$$\mathbf{D}_{\text{cris}}(M^{\text{rig}}) \cong R^{\text{rig}} \otimes_R \mathbf{D}_{\text{cris}}(M).$$

Theorem 4.9 defines an isomorphism $\mathbf{N}(M)^{\text{rig}}/\pi\mathbf{N}(M)^{\text{rig}} \xrightarrow{\sim} (\mathbf{N}(M)^{\text{rig}})^{\Gamma}$. Since $\mathbf{N}(M)^{\text{rig}}$ has an underlying R -module structure, it is natural to ask whether this isomorphism can be defined over a smaller ring, such as $R[1/p] \widehat{\otimes} \mathbf{B}_{\text{rig},F}^+$.

Suppose that L/\mathbf{Q}_p is a finite extension, and suppose that T is a positive crystalline representation of G_F of rank d over \mathcal{O}_L , with Hodge–Tate weights in $[-h, 0]$. Then $\mathbf{N}(T)$ is free of rank d over $\mathcal{O}_L \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$; choose a basis $\{e_1, \dots, e_d\}$ of $\mathbf{N}(T)$ and let its image modulo π be a basis of $\mathbf{D}_{\text{cris}}(T)$. Write A for the isomorphism $\mathbf{N}(T)^{\text{rig}}/\pi\mathbf{N}(T)^{\text{rig}} \xrightarrow{\sim} (\mathbf{N}(T)^{\text{rig}})^{\Gamma}$ with respect to these bases.

Recall that $t := \log(1 + \pi) \in \mathbf{B}_{\text{rig},F}^+$.

Lemma 4.11. *Let the notation be as above, and write $A = 1 + \sum_{m \geq 1} A^{(m)} t^m$ where $A^{(m)}$ is a matrix over $\mathcal{O}_L \otimes_{\mathbf{Z}_p} \mathcal{O}_F$ with entries $a_{ij}^{(m)}$. Then $v_p(a_{ij}^{(m)}) \geq -\frac{(2p-1)}{(p-1)^2} m$, and in particular, the entries of A (viewed as elements of $L \otimes_{\mathbf{Q}_p} \mathbf{B}_{\text{rig},F}^+$) are bounded by 1 on the disk $v_p(\pi) > p - 1$.*

Proof. Write $G = 1 + \sum_{m \geq 1} G^{(m)} t^m$ for the matrix of a non-torsion element $\gamma \in \Gamma$, where $G^{(m)}$ has entries $g_{ij}^{(m)}$ in $L \otimes_{\mathbf{Q}_p} F$. Now the exponential series $\pi = \exp(t) - 1$ converges for any t with $v_p(t) > \frac{1}{p-1}$ and defines a bijection between the affinoid disks $v_p(t) \geq \rho$ and $v_p(\pi) \geq \rho$ for any rational $\rho > \frac{1}{p-1}$. Since Γ acts by automorphisms on $\mathbf{N}(M \otimes_R \mathcal{O}_L)$, G must be invertible over $\mathcal{O}_L \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$, and therefore its entries must be bounded by 1 on any such disk. Therefore $v_p(g_{ij}^{(m)}) \geq -\frac{1}{p-1} m$.

Now the image of A is Γ -invariant, so we must have $G\gamma(A) = A$, or equivalently,

$$\sum_{r=0}^{m-1} G^{(m-r)} A^{(r)} \chi(\gamma)^r = (1 - \chi(\gamma)^m) A^{(m)}$$

for all m . By induction on m , this implies that the matrix $A^{(m)}$ can be written as a sum of terms of the form

$$\frac{\prod_{1 \leq a \leq M} G^{(\lambda_a)} \chi(\gamma)^N}{\prod_{b \in B} (1 - \chi(\gamma)^b)}$$

where M and N are integers, the λ_a are positive integers satisfying $\lambda_1 + \dots + \lambda_M = m$, and B is a subset of $\{1, \dots, m\}$. Using the previous inequality, any such term has valuation bounded below by

$$-\frac{1}{p-1} m - \sum_{1 \leq r \leq m} v_p(1 - \chi(\gamma)^m) \geq -\left(\frac{1}{p-1} + \frac{p}{(p-1)^2}\right) m = -\frac{2p-1}{(p-1)^2} m,$$

using the bound $\sum_{1 \leq r \leq m} v_p(1 - \chi(\gamma)^m) \leq \frac{pm}{(p-1)^2}$ given in [Ber04, §IV.1]. \square

Now we consider the φ -action on $\mathbf{N}(T)$. Let $P \in \text{Mat}_{d \times d}(\mathcal{O}_L \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+)$ be the matrix of φ on $\mathbf{N}(T)$ with respect to $\{e_i\}$ and let $\Phi \in \text{Mat}_{d \times d}(\mathcal{O}_L \otimes_{\mathbf{Z}_p} \mathcal{O}_F)$ be the matrix of φ on $\mathbf{N}(T)/\pi$; Φ is the “constant term” of P . Then Φ is invertible over $L \otimes_{\mathbf{Q}_p} F$, and more precisely, $p^h \Phi^{-1} \in \text{Mat}_{d \times d}(\mathcal{O}_L \otimes_{\mathbf{Z}_p} \mathcal{O}_F)$.

Let $\rho > 0$. Since $\varphi(\pi) = (1 + \pi)^p - 1 \in \pi^p + p\mathbf{Z}_p[[\pi]]$, there is some N_ρ such that φ^{N_ρ} carries the disk $v_p(\pi) \geq \rho$ into a sub-disk of $v_p(\pi) > \frac{2p-1}{(p-1)^s}$, and the induced map the other way on rings carries functions converging for $v_p(\pi) > \frac{2p-1}{(p-1)^s}$ to functions converging for $v_p(\pi) \geq \rho$.

Lemma 4.12. *Let the notation be as above. The entries of A have valuation bounded below by $-N_\rho h$ on the disk $v_p(\pi) \geq \rho$.*

Proof. Since A is Frobenius-equivariant, we have the relation $A\Phi = P\varphi(A)$. Iterating N_ρ times and multiplying by $p^{N_\rho h}$, we obtain $p^{N_\rho h} A = (P\varphi(P) \cdots \varphi^{N_\rho}(P)) \varphi^{N_\rho}(A) (\varphi^{N_\rho}(p^h \Phi^{-1}) \cdots p^h \Phi^{-1})$. Thus, $p^{N_\rho h} A$ has entries which are bounded by 1 on the disk $v_p(\pi) \geq \rho$. \square

Now we return to M over R . Let $U = \text{Spec } R[1/f_U, 1/p] \subset \text{Spec } R[1/p]$ with $f_U \in R \setminus pR$ be an open affine subspace such that $\mathbf{N}(M)[1/p]|_U$ is free. Choose a basis $\{e'_1, \dots, e'_d\}$ for $\mathbf{N}(M)[1/p]|_U$ and write A' for the matrix of the isomorphism above with respect to the image of $\{e'_i\}$ in $\mathbf{D}_{\text{cris}}(M^{\text{rig}})|_U$ and $\{e'_i\}$; we may assume $\{e'_i\} \subset \mathbf{N}(M)[1/f_U]$. Writing $\mathbf{N}_{\{e'_i\}}$ for the $R \widehat{\otimes} \mathbf{A}_F^+$ -span of $\{e'_i\}$, there is some integer β such that $p^{-\beta} \mathbf{N}_{\{e'_i\}} \cap \mathbf{N}(M)[1/f_U] = \mathbf{N}(M)[1/f_U]$, since $\mathbf{N}(M)[1/f_U]$ is finitely generated over $(R \widehat{\otimes} \mathbf{A}_{\mathbf{Q}_p}^+)[1/f_U]$ and $\{e'_i\}$ is a basis for $\mathbf{N}(M)[1/f_U, 1/p]$.

Let $x \in U$ be a closed point with residue field L and let $\{e_1, \dots, e_d\}$ be a basis of $\mathbf{N}(M \otimes_R \mathcal{O}_L)$. Recall that $p^{\alpha(h)} \mathbf{N}(M \otimes_R \mathcal{O}_L) \subset \mathbf{N}(M) \otimes_R \mathcal{O}_L$; it follows that $p^{\alpha(h)+\beta} \mathbf{N}(M \otimes_R \mathcal{O}_L) \subset \mathbf{N}_{\{e'_i\}} \otimes_R \mathcal{O}_L$. Thus, if X is the matrix for the image of $\{e'_i\}$ in $\mathbf{N}_{\{e'_i\}} \otimes_R \mathcal{O}_L$ with respect to $\{e_i\}$, X and $p^{\alpha(h)+\beta} X^{-1}$ have coefficients in $\mathcal{O}_L \otimes_{\mathbf{Z}_p} \mathbf{A}_F^+$. It follows that $A' \otimes_{R[1/p]} L$ is a matrix with entries in $L \otimes_{\mathbf{Q}_p} \mathbf{B}_{\text{rig}, F}^+$ that have valuation bounded below by $-(\alpha(h) + \beta + N_\rho h)$ on the disk $v_p(\pi) \geq \rho$.

In other words, there is some integer k such that $f_U^k A'$ has entries in $R^{\text{rig}} \widehat{\otimes} \mathbf{B}_{\text{rig}, F}^+$ which are bounded on closed disks $v_p(\pi) \geq \rho$. Thus, A' has entries in $(R[1/p] \widehat{\otimes} \mathbf{B}_{\text{rig}, F}^+)[1/f_U]$, by Proposition A.12.

Corollary 4.13. *There is an isomorphism of line bundles over $R[1/p] \widehat{\otimes} \mathbf{B}_{\text{rig}, F}^+$*

$$\text{Det}_{R[1/p] \widehat{\otimes} \mathbf{B}_{\text{rig}, F}^+} (R[1/p] \widehat{\otimes} \mathbf{B}_{\text{rig}, F}^+ \otimes_R \mathbf{D}_{\text{cris}}(M)) \xrightarrow{\sim} \text{Det}_{R[1/p] \widehat{\otimes} \mathbf{B}_{\text{rig}, F}^+} (R[1/p] \widehat{\otimes} \mathbf{B}_{\text{rig}, F}^+ \otimes_R \widehat{\otimes}_{\mathbf{Z}_p} \mathbf{A}_F^+ \mathbf{N}(M))$$

which agrees with the isomorphism $(\pi/t)^{r_1+\dots+r_d} \cdot \det(\text{inc}_M)$ of Proposition 4.5.

5. GALOIS COHOMOLOGY

Let K_∞/\mathbf{Q}_p be an abelian p -adic Lie extension with Galois group $G := G(K_\infty/\mathbf{Q}_p)$ and associated Iwasawa algebra $\Lambda := \mathbf{Z}_p[[G]]$. We write J_G for its Jacobson radical. Let R be a complete local noetherian \mathbf{Z}_p -algebra with finite residue field, flat over \mathbf{Z}_p , and let \mathfrak{m}_R be its maximal ideal. Then we define the ring

$$\Lambda_R(G) := \Lambda \widehat{\otimes}_{\mathbf{Z}_p} R$$

which has Jacobson radical $\mathfrak{J} = J_G \Lambda_R(G) + \mathfrak{m}_R \Lambda_R(G)$. If M is a family of $G_{\mathbf{Q}_p}$ -representations over R , we define the $\Lambda_R(G)$ -module

$$\mathbf{T}(M) := \Lambda \widehat{\otimes}_{\mathbf{Z}_p} M$$

where the $\Lambda_R(G)$ -module structure is induced by $(\lambda \otimes r) \cdot (\lambda' \otimes m) := \lambda \lambda' \otimes r m$. In addition we endow the latter with the following action of $G_{\mathbf{Q}_p}$: for σ in the latter group we set $\sigma(\lambda \otimes m) := \lambda \bar{\sigma}^{-1} \otimes \sigma m$. Now we define the Iwasawa cohomology of a family M of $G_{\mathbf{Q}_p}$ -representations as

$$\mathbf{H}_{\text{Iw}}^i(M) := \mathbf{H}_{\text{Iw}}^i(K_\infty, M) := \mathbf{H}^i(\mathbf{Q}_p, \mathbf{T}(M)).$$

Proposition 5.1. *Suppose that $K_\infty = \mathbf{Q}_p(\mu_{p^\infty})$ and M is crystalline with Hodge–Tate weights in $[0, h]$. Then there is a canonical isomorphism*

$$\mathbf{H}_{\text{Iw}}^1(M) \xrightarrow{\sim} (\pi^{-1} \mathbf{N}(M))^{\psi=1}$$

functorial in the coefficients.

Suppose in addition that for every maximal ideal $\mathfrak{m} \subset R[1/p]$ the specialization $(R[1/p]/\mathfrak{m}) \otimes_R M$ has no quotient isomorphic to the trivial representation. Then $\mathbf{H}_{\text{Iw}}^1(M) \xrightarrow{\sim} \mathbf{N}(M)^{\psi=1}$.

Here the map $\psi : \mathbf{N}(M) \rightarrow \mathbf{N}(M)$ is defined to be the composition

$$\mathbf{N}(M) \subset \varphi^* \mathbf{N}(M) \xrightarrow{\psi \otimes 1} \mathbf{N}(M)$$

and we use the non-negativity of the Hodge–Tate weights of M to assume $\mathbf{N}(M) \subset \varphi^* \mathbf{N}(M)$.

We first need a preparatory lemma, which extends the Fontaine isomorphism $\mathbf{D}(M)^{\psi=1} \xrightarrow{\sim} \mathbf{H}_{\text{Iw}}^1(M)$ to formal families.

Lemma 5.2. *There is a natural isomorphism $\mathbf{D}(M)^{\psi=1} \xrightarrow{\sim} \mathbf{H}_{\text{Iw}}^1(M)$.*

Proof. Let $\Delta_K \subset \Gamma$ be the p -torsion subgroup and recall that Herr showed [Her98] that the Galois cohomology of a finite \mathbf{Z}_p -linear G_K -representation T is functorially computed by the complex

$$\mathbf{D}(T)^{\Delta_K} \xrightarrow{(\varphi^{-1}, \gamma_K - 1)} \mathbf{D}(T)^{\Delta_K} \oplus \mathbf{D}(T)^{\Delta_K} \xrightarrow{(\gamma_K - 1) \oplus (1 - \varphi)} \mathbf{D}(T)^{\Delta_K}$$

In particular, this is compatible with corestriction. Here $\Delta_K \subset \Gamma_K$ is the p -torsion subgroup and $\gamma_K \in \Gamma_K/\Delta_K$ is a topological generator. It follows by taking limits that the Galois cohomology of M is computed by the same complex

$$\mathbf{D}(M)^{\Delta_K} \xrightarrow{(\varphi^{-1}, \gamma_K^{-1})} \mathbf{D}(M)^{\Delta_K} \oplus \mathbf{D}(M)^{\Delta_K} \xrightarrow{(\gamma_K^{-1}) \oplus (1-\varphi)} \mathbf{D}(M)^{\Delta_K}$$

Furthermore, the commutative diagram

$$\begin{array}{ccccc} \mathbf{D}(T)^{\Delta_K} & \xrightarrow{(\varphi^{-1}, \gamma_K^{-1})} & \mathbf{D}(T)^{\Delta_K} \oplus \mathbf{D}(T)^{\Delta_K} & \xrightarrow{(\gamma_K^{-1}) \oplus (1-\varphi)} & \mathbf{D}(T)^{\Delta_K} \\ \downarrow \text{id} & & \downarrow -\psi \oplus \text{id} & & \downarrow -\psi \\ \mathbf{D}(T)^{\Delta_K} & \xrightarrow{(\psi^{-1}, \gamma_K^{-1})} & \mathbf{D}(T)^{\Delta_K} \oplus \mathbf{D}(T)^{\Delta_K} & \xrightarrow{(\gamma_K^{-1}) \oplus (1-\psi)} & \mathbf{D}(T)^{\Delta_K} \end{array}$$

defines a quasi-isomorphism between the top and bottom rows. We obtain an exact sequence

$$0 \rightarrow \mathbf{D}(M)^{\Delta_K, \psi=1}/(\gamma_K - 1) \rightarrow H^1(G_K, M) \rightarrow (\mathbf{D}(M)/(\psi - 1))^{\Gamma_K} \rightarrow 0$$

compatible with corestriction. If we take $K = \mathbf{Q}_p(\mu_{p^n})$ and consider the limit over all n , we obtain the desired isomorphism

$$\mathbf{D}(M)^{\psi=1} \xrightarrow{\sim} H_{\text{Iw}}^1(M)$$

since for $n \gg 0$, Δ is trivial. \square

Proof of Proposition 5.1. It suffices to show that $(\pi^{-1}\mathbf{N}(M))^{\psi=1} = \mathbf{D}(M)^{\psi=1}$, and to see this, it suffices to show that $(\pi^{-1}\mathbf{N}(M)[1/p])^{\psi=1} = \mathbf{D}(M)[1/p]^{\psi=1}$. But $\mathbf{D}(M)^{\Delta, \psi=1} = H_{\text{Iw}}^1(M)$ is a finite $\Lambda_R(G)$ -module by [FK06, Proposition 1.6.5(2)], so $\mathbf{D}(M)^{\psi=1}$ and $(\pi^{-1}\mathbf{N}(M))^{\psi=1} \subset \mathbf{D}(M)^{\psi=1}$ are, as well.

Let $x \in \mathbf{D}(M)[1/p]^{\psi=1}$. Then for every maximal ideal $\mathfrak{m} \subset R[1/p]$, the image of x in $\mathbf{D}((R/\mathfrak{m}) \otimes_R M)[1/p]$ lands in $\mathbf{D}((R[1/p]/\mathfrak{m}) \otimes_R M)[1/p]^{\psi=1} \subset \pi^{-1}\mathbf{N}((R[1/p]/\mathfrak{m}) \otimes_R M)[1/p]$. Since the formation of $\mathbf{N}(M)[1/p]$ commutes with base change on $R[1/p]$, it follows that $x \in \pi^{-1}\mathbf{N}(M)[1/p]$.

The second claim follows similarly. \square

Let d be the R -rank of M .

Proposition 5.3. *The Iwasawa cohomology groups $H_{\text{Iw}}^i(K_\infty, M)$ are finite $\Lambda_R(G)$ -modules. In fact, $H_{\text{Iw}}^i(K_\infty, M) = 0$ for $i \notin \{1, 2\}$, $H_{\text{Iw}}^1(K_\infty, M)[1/p]$ has generic rank d , and $H_{\text{Iw}}^2(K_\infty, M)$ is R -finite (and annihilated by an ideal of $\Lambda_{\mathbf{Z}_p}(G)$).*

Proof. The $\Lambda_R(G)$ -finiteness statement follows from [FK06, Proposition 1.6.5(2)]. The vanishing statements follow from their classical analogs.

A consideration of Tate local duality implies that $H_{\text{Iw}}^2(K_\infty, M) \cong (M^\vee(1)^H)^\vee \cong M(-1)_H$, which is R -finite. The perfectness of $R\Gamma_{\text{Iw}}(K_\infty, M) := R\Gamma(\mathbf{Q}_p, \mathfrak{M})$ combined with the fact that $H_{\text{Iw}}^i(K_\infty, M) = 0$ for $i \geq 3$ implies that the formation of $H_{\text{Iw}}^2(K_\infty, M)$ commutes with base change on $\Lambda_R(G)$, and in particular, with base change on R . Choose a set of generators m_1, \dots, m_k for $H_{\text{Iw}}^2(K_\infty, M)$ as an R -module. For a closed point $x \in \text{Spec } R[1/p]$, let \mathcal{O}_{L_x} denote the ring of integers in the residue field at x . Then there is a finitely generated ideal $I_x \subset \Lambda_{\mathbf{Z}_p}(G)$ which annihilates $H_{\text{Iw}}^2(K_\infty, M \otimes_R \mathcal{O}_{L_x}) = H_{\text{Iw}}^2(K_\infty, M) \otimes_R \mathcal{O}_{L_x}$. This implies that I_x annihilates $H_{\text{Iw}}^2(K_\infty, M)$ in a Zariski open neighborhood of x , and since $\text{Spec } \Lambda_R(G)$ is quasi-compact, we can multiply finitely many ideals together to get a finitely generated ideal of $\Lambda_{\mathbf{Z}_p}(G)$ which annihilates $H_{\text{Iw}}^2(K_\infty, M)$.

The preceding paragraph implies that over an open dense subspace of $\text{Spec } \Lambda_R(G)[1/p]$ (obtained by inverting elements coming from $\Lambda_{\mathbf{Z}_p}(G)[1/p]$), the formation of $H_{\text{Iw}}^1(K_\infty, M)[1/p]$ also commutes with base change on $R[1/p]$. But then comparison with the classical case shows that $H_{\text{Iw}}^1(K_\infty, M)[1/p]$ has generic rank d . \square

For our later calculations we need the convergent cohomological spectral sequence for any (R', R) -bimodule Y :

$$(2) \quad E_2^{i,j} := \text{Tor}_{-i}^R(Y, H_{\text{Iw}}^j(M)) \Rightarrow H_{\text{Iw}}^{i+j}(Y \otimes_R M),$$

which is induced from the isomorphism

$$Y \otimes_R^{\mathbf{L}} R\Gamma_{\text{Iw}}(M) \cong R\Gamma_{\text{Iw}}(Y \otimes_R M),$$

proved in [FK06]. We obtain the following exact sequence of terms in lower degree

(3)

$$0 \longrightarrow \mathrm{Tor}_2^R(Y, H_{I_w}^2(M)) \longrightarrow Y \otimes_R H_{I_w}^1(M) \longrightarrow H_{I_w}^1(Y \otimes_R M) \longrightarrow \mathrm{Tor}_1^R(Y, H_{I_w}^2(M)) \longrightarrow 0$$

as well as the isomorphisms

$$Y \otimes_R H_{I_w}^2(M) \cong H_{I_w}^2(Y \otimes_R M) \quad \text{and} \quad \mathrm{Tor}_i^R(Y, H_{I_w}^1(M)) \cong \mathrm{Tor}_{i+2}^R(Y, H_{I_w}^2(M))$$

for $i \geq 1$. By Proposition 5.3 we obtain pseudo-isomorphisms, i.e., isomorphisms up to pseudo-null modules,

(4)

$$Y \otimes_R H_{I_w}^1(M) \sim H_{I_w}^1(Y \otimes_R M) \quad \text{and} \quad \mathrm{Tor}_i^R(Y, H_{I_w}^1(M)) \sim 0$$

for $i \geq 1$.

In the Fontaine-Lafaille range we obtain the following result which is not needed elsewhere in this article.

prop:phiN

Proposition 5.4. *If M is crystalline with Hodge-Tate weights in $[a, b]$ satisfying $h = b - a < p - 1$, then $(\varphi^* \mathbf{N}(M))^{\psi=0}$ is a free $R \hat{\otimes} \Lambda(G)$ -module of rank r .*

Proof. From [SRV13] we obtain the isomorphisms

$$(\varphi^* \mathbf{N}(M/I^n M))^{\psi=0} = \bigoplus_{j=1}^{p-1} (1 + \pi)^j \otimes \mathbf{N}(M/I^n M)$$

which induces after taking projective limits an isomorphism of $R \hat{\otimes} \Lambda(G)$ -modules

$$(\varphi^* \mathbf{N}(M))^{\psi=0} = \bigoplus_{j=1}^{p-1} (1 + \pi)^j \otimes \mathbf{N}(M).$$

Now choose lifts n_1, \dots, n_r in $\mathbf{N}(M)$ of any $\mathbf{A}_{\mathbf{Q}_p}^+$ -basis $\bar{n}_1, \dots, \bar{n}_r$ of $\mathbf{N}(M/IM)$. Since

$$\begin{aligned} (\varphi^* \mathbf{N}(M))^{\psi=0} / I(\varphi^* \mathbf{N}(M))^{\psi=0} &= \bigoplus_{j=1}^{p-1} (1 + \pi)^j \otimes \mathbf{N}(M) / I\mathbf{N}(M) \\ &\cong \bigoplus_{j=1}^{p-1} (1 + \pi)^j \otimes \mathbf{N}(M/IM) \end{aligned}$$

is isomorphic to $\Lambda(G)^r$ with basis $\bar{n}_1, \dots, \bar{n}_r$ by (loc. cit.) and the fact that by assumption the base change map for \mathbf{N} is an isomorphism, we see by the Nakayama lemma that the map

$$(R \hat{\otimes} \Lambda(G))^r \rightarrow (\varphi^* \mathbf{N}(M))^{\psi=0}$$

induced by the choice of n_1, \dots, n_r is surjective with kernel, say, C . By induction on d and using torsionfreeness of $\varphi^* \mathbf{N}(M)$ we conclude that $C/X_d C = 0$ whence $C = 0$ by the Nakayama lemma again. The proposition follows and we even get an explicit basis. \square

We now specialize to the following setting (which will be crucial for the construction of the regulator maps): Let F_∞ be the unramified \mathbf{Z}_p -extension of \mathbf{Q}_p . We set $K_\infty = F_\infty(\mu_{p^\infty})$, $U = \mathrm{Gal}(F_\infty/\mathbf{Q}_p) \cong \mathrm{Gal}(K_\infty/\mathbf{Q}_{p,\infty})$ and $G = \mathrm{Gal}(K_\infty/\mathbf{Q}_p)$. We regard Γ as a subgroup of G , by identifying it with $\mathrm{Gal}(K_\infty/F_\infty)$, so we have $G = U \times \Gamma$.

Following [LZ14] we define

$$\mathbf{N}_\infty(M) := \mathbf{N}(M) \hat{\otimes}_{\mathbf{Z}_p} S_\infty$$

Here $S_\infty := \varprojlim_n \mathcal{O}_{F_n}$ with the transition maps being the trace maps. This is called the *Yager module*, and it is free of rank one over $\Lambda(U)$ and naturally endowed with a Frobenius action. Moreover, S_∞ is naturally contained in $\Lambda_{\mathcal{O}_{\hat{F}_\infty}}(U)$, and the induced subspace topology coincides with the inverse limit topology (see the discussion in §3 in loc. cit.). Indeed, setting

$$\Lambda_{\mathbf{Z}_p}(U)_{\tau_p} := \{x \in \widehat{\mathbf{Z}_p^{\mathrm{nr}}} \otimes_{\mathbf{Z}_p} \Lambda_{\mathbf{Z}_p}(U) \mid (\phi \otimes 1)x = (1 \otimes \bar{\tau}_p)x\},$$

where $\bar{\tau}_p$ denotes the image of the unique lift τ_p in $G(\mathbf{Q}_p^{ab}/\mathbf{Q}_p(\mu(p)))$ of the arithmetic Frobenius $\sigma \in G(\mathbf{Q}_p^{\mathrm{nr}}/\mathbf{Q}_p)$ one checks ([LZ14, Prop. 3.6]) that

(5)

$$S_\infty \cong \Lambda_{\mathbf{Z}_p}(U)_{\tau_p}.$$

Similarly, we consider

$$\mathcal{H}_{\mathbf{Q}_p}(U)_{\tau_p} := \{x \in \widehat{\mathbf{Z}_p^{\mathrm{nr}}} \otimes_{\mathbf{Z}_p} \mathcal{H}_{\mathbf{Q}_p}(U) \mid (\sigma \otimes 1)x = (1 \otimes \bar{\tau}_p)x\}$$

f:Yager

and

$$\mathcal{H}_{\mathbf{Q}_p}(G)_{\tau_p} := \{x \in \widehat{\mathbf{Z}_p^{nr}} \otimes_{\mathbf{Z}_p} \mathcal{H}_{\mathbf{Q}_p}(G) \mid (\sigma \otimes 1)x = (1 \otimes \overline{\tau_p})x\},$$

where in the latter case $\overline{\tau_p}$ denotes the image of τ_p in $G := \Gamma \times U$ with $\Gamma \cong \mathbf{Z}_p$.

Lemma 5.5. (1) $\Lambda_{\mathbf{Z}_p}(U)_{\tau_p}$ and $\mathcal{H}_{\mathbf{Q}_p}(U)_{\tau_p}$ are free rank one modules over $\Lambda_{\mathbf{Z}_p}(U)$ and $\mathcal{H}_{\mathbf{Q}_p}(U)$, respectively.

(2) $\Lambda_{\mathbf{Z}_p}(G)_{\tau_p}$ and $\mathcal{H}_{\mathbf{Q}_p}(G)_{\tau_p}$ are free rank one modules over $\Lambda_{\mathbf{Z}_p}(G)$ and $\mathcal{H}_{\mathbf{Q}_p}(G)$, respectively. Moreover, we have

$$\mathcal{H}_{\mathbf{Q}_p}(U)_{\tau_p} \widehat{\otimes}_{\mathbf{Q}_p} \mathcal{H}_{\mathbf{Q}_p}(\Gamma) = \mathcal{H}_{\mathbf{Q}_p}(G)_{\tau_p}.$$

Proof. (1) Let \mathcal{O} be either \mathbf{Z}_p or $\widehat{\mathbf{Z}_p^{nr}}$ and set $L := \mathcal{O}[\frac{1}{p}]$. For the purpose of the proof we identify $\Lambda_{\mathcal{O}}(U)$ and $\mathcal{H}_L(U)$ with $\mathcal{O}[[T]]$ and the subring of $L[[T]]$ consisting of power series which converge on the open unit disk, respectively. Then the defining condition of e.g. $\Lambda_{\mathbf{Z}_p}(U)_{\tau_p}$ is equivalent to $(\sigma - 1)\lambda = T\lambda$ whence we obtain the isomorphism

$$\Lambda_{\mathbf{Z}_p}(U)_{\tau_p} \cong \{(a_i) \in \varprojlim_{i \geq 0, \sigma-1} \widehat{\mathbf{Z}_p^{nr}} \mid a_0 \in \mathbf{Z}_p\}$$

sending $\lambda = \sum a_i T^i$ to (a_i) . Fix any such λ_0 with $a_0 = 1$, the existence of which stems from the exact sequence

$$0 \rightarrow \mathbf{Z}_p \rightarrow \widehat{\mathbf{Z}_p^{nr}} \xrightarrow{\sigma-1} \widehat{\mathbf{Z}_p^{nr}} \rightarrow 0.$$

Then λ_0 is a unit in $\Lambda_{\widehat{\mathbf{Z}_p^{nr}}}(U)$ and for any λ in $\Lambda_{\mathbf{Z}_p}(U)_{\tau_p}$ or $\mathcal{H}_{\mathbf{Q}_p}(U)_{\tau_p}$ we have that

$$\sigma(\lambda \lambda_0^{-1}) = \sigma(\lambda) \sigma(\lambda_0)^{-1} = \overline{\tau_p} \lambda \overline{\tau_p}^{-1} \lambda_0^{-1} = \lambda \lambda_0^{-1},$$

i.e., $\lambda \lambda_0^{-1}$ belongs to $\Lambda_{\mathbf{Z}_p}(U)$ or $\mathcal{H}_{\mathbf{Q}_p}(U)$, respectively. This implies that λ_0 forms a basis in both cases.

(2) Since $\lambda_0 \in \Lambda_{\mathbf{Z}_p}(U)_{\tau_p} \subseteq \Lambda_{\mathbf{Z}_p}(G)_{\tau_p}$ the last argument applies in the same way. From this the last statement is clear, too. \square

Using the same argument as in the proof of [LZ14, Theorem 4.4, Proposition 4.5] we obtain the following sharpening of Proposition 5.1:

Proposition 5.6. *If M is crystalline with Hodge-Tate weights in $[0, h]$, then there is a canonical isomorphism*

$$H_{Iw}^1(K_\infty, M) \cong \mathbf{N}_\infty(M)^{\psi=1}.$$

6. REGULATOR MAPS FOR FAMILIES

We have to extend the various regulator maps to families and thus generalize the main result from [LZ14]. Let F be any finite unramified extension of \mathbf{Q}_p , and let F_∞ be the unramified \mathbf{Z}_p -extension of F . We set $K_\infty = F_\infty(\mu_{p^\infty})$, $U = \text{Gal}(K_\infty/\mathbf{Q}_p)$ and $G = \text{Gal}(K_\infty/\mathbf{Q}_p)$. We regard Γ as a subgroup of G , by identifying it with $\text{Gal}(K_\infty/F_\infty)$, so we have $G = U \times \Gamma$.

6.1. Γ -regulator. Let M now be a crystalline family with non-negative Hodge-Tate weights and not containing the trivial representation as a quotient (of any M_n). Then we have the following map:

$$\mathcal{L}_{\mathbf{Q}_p, M, \xi}^\Gamma : H_{Iw}^1(\mathbf{Q}_p(\mu_{p^\infty})/\mathbf{Q}_p, M) \cong \mathbf{N}(M)^{\psi=1} \xrightarrow{1-\varphi} (\varphi^* \mathbf{N}(M))^{\psi=0} \hookrightarrow (\mathcal{H}_{\mathbf{Q}_p}(\Gamma) \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})$$

and the last inclusion follows from Corollary 4.8 and the identification

$$(6) \quad (\mathbf{B}_{\text{rig}, \mathbf{Q}_p}^+ \widehat{\otimes} R^{\text{rig}})^{\psi=0} \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) \cong (\mathcal{H}_{\mathbf{Q}_p}(\Gamma) \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}).$$

Note that the latter is a projective $\mathcal{H}_{\mathbf{Q}_p}(\Gamma) \widehat{\otimes} R^{\text{rig}}$ -module because M is assumed crystalline.

For any finite extension E/\mathbf{Q}_p contained in F_∞ , we may apply the above construction to the induced representation $M' := \text{Ind}_E^{\mathbf{Q}_p} M$. Using Shapiro's Lemma and Frobenius reciprocity (see §4.3 in [LZ14] for a similar construction), we then obtain the following map:

$$\mathcal{L}_{E, M, \xi}^\Gamma : H_{Iw}^1(E(\mu_{p^\infty})/E, M) \longrightarrow (\mathcal{H}_E(\Gamma) \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}).$$

6.2. G -regulator. Assume first that $F = \mathbf{Q}_p$, i.e., $U \cong \mathbf{Z}_p$. Extending the observations from (loc. cit.) one easily shows that

$$(\varphi^*(\mathbf{N}_\infty(M))^{\psi=0} \cong (\varphi^*\mathbf{N}(M))^{\psi=0} \widehat{\otimes}_{\mathbf{Z}_p} S_\infty.$$

Using Proposition 5.6 we obtain the following map

$$\mathcal{L}_{M,\xi}^G : H_{\text{Iw}}^1(K_\infty, M) \cong \mathbf{N}_\infty(M)^{\psi=1} \xrightarrow{1-\varphi} (\varphi^*\mathbf{N}_\infty(M))^{\psi=0} \hookrightarrow (\mathcal{H}_{\bar{L}}(G) \widehat{\otimes}_{\mathbf{Q}_p} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})$$

generalizing definition 4.6 in (loc. cit.). More precisely, as in §4.2 of (loc. cit.) one has an embedding

$$(7) \quad S_\infty \hookrightarrow \mathcal{H}_{\mathbf{Q}_p}(U)_{\tau_p} \hookrightarrow \mathcal{H}_{\bar{L}}(U)$$

which is continuous with respect to the inverse limit topology on S_∞ and the Fréchet topology on the target. Hence the natural map on the algebraic tensor product

$$(\varphi^*\mathbf{N}(M))^{\psi=0} \otimes_{\mathbf{Z}_p} S_\infty \rightarrow (\mathcal{H}_{\mathbf{Q}_p}(\Gamma) \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) \otimes_{\mathbf{Q}_p} \mathcal{H}_{\bar{L}}(U)$$

induced from the embedding in Corollary 4.8 plus (6) and (7) extends to a continuous $R \widehat{\otimes}_{\mathbf{Z}_p} \Lambda(U \times \Gamma)$ -equivariant map

$$(\varphi^*\mathbf{N}(M))^{\psi=0} \widehat{\otimes}_{\mathbf{Z}_p} S_\infty \rightarrow (\mathcal{H}_{\bar{L}}(U) \widehat{\otimes}_{\mathbf{Q}_p} \mathcal{H}_{\mathbf{Q}_p}(\Gamma) \widehat{\otimes}_{\mathbf{Q}_p} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) \cong (\mathcal{H}_{\bar{L}}(G) \widehat{\otimes}_{\mathbf{Q}_p} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})$$

The image of this map is contained in

$$(\mathcal{H}_{\mathbf{Q}_p}(U)_{\tau_p} \widehat{\otimes}_{\mathbf{Q}_p} \mathcal{H}_{\mathbf{Q}_p}(\Gamma) \widehat{\otimes}_{\mathbf{Q}_p} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) = (\mathcal{H}_{\mathbf{Q}_p}(G)_{\tau_p} \widehat{\otimes}_{\mathbf{Q}_p} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})$$

Since $\mathcal{H}_{\mathbf{Q}_p}(G)_{\tau_p} \widehat{\otimes}_{\mathbf{Q}_p} R^{\text{rig}}$ is a free rank one $\mathcal{H}_{\mathbf{Q}_p}(G) \widehat{\otimes}_{\mathbf{Q}_p} R^{\text{rig}}$ -module by Lemma 5.5, the source and target are finite $\mathcal{H}_{\mathbf{Q}_p}(G) \widehat{\otimes}_{\mathbf{Q}_p} R^{\text{rig}}$ -modules of the same generic rank.

The extension to finite unramified extensions F/\mathbf{Q}_p (which we may and do assume to be linearly disjoint from the unramified \mathbf{Z}_p -extension of \mathbf{Q}_p) is carried out as usual by applying the above construction to the induced representation $M' := \text{Ind}_F^{\mathbf{Q}_p} M$ and using Shapiro's lemma and Frobenius reciprocity; see §4.3 in (loc. cit.) for the details.

The first part of the following theorem generalizes Thm. 4.7 in (loc. cit.) to families:

Theorem 6.1. *Assume that M is crystalline with Hodge-Tate weights ≥ 0 . Then the following hold:*

- (1) *For any finite extension E/\mathbf{Q}_p contained in F_∞ , we have a commutative diagram*

$$\begin{array}{ccc} H_{\text{Iw}}^1(K_\infty, M) & \xrightarrow{\mathcal{L}_{M,\xi}^G} & (\mathcal{H}_{\bar{L}}(G) \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) \\ \downarrow & & \downarrow \\ H_{\text{Iw}}^1(E(\mu_{p^\infty}), M) & \xrightarrow{\mathcal{L}_{M,\xi}^{G'}} & (\mathcal{H}_{\bar{L}}(G') \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}). \end{array}$$

Here $G' = \text{Gal}(E(\mu_{p^\infty})/\mathbf{Q}_p)$, the right-hand vertical arrow is the map on distributions corresponding to the projection $G \rightarrow G'$, and the map $\mathcal{L}_{M,\xi}^{G'}$ is defined by

$$\mathcal{L}_{M,\xi}^{G'} = \sum_{\sigma \in \text{Gal}(E/\mathbf{Q}_p)} [\sigma] \cdot \mathcal{L}_{E,M,\xi}^\Gamma(\sigma^{-1} \circ x),$$

where $\mathcal{L}_{E,M,\xi}^\Gamma$ is the cyclotomic regulator map for $E(\mu_{p^\infty})/E$. Moreover, the map $\mathcal{L}_{M,\xi}^G$ is injective.

- (2) *For any ideal $P \subset R$ such that R/P is flat over \mathbf{Z}_p , there is a commutative diagram*

$$\begin{array}{ccc} H_{\text{Iw}}^1(K_\infty, M) & \xrightarrow{\mathcal{L}_{M,\xi}^G} & (\mathcal{H}_{\bar{L}}(G) \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) \\ \downarrow & & \downarrow \\ H_{\text{Iw}}^1(K_\infty, M/P) & \xrightarrow{\mathcal{L}_{M/P,\xi}^G} & (\mathcal{H}_{\bar{L}}(G) \widehat{\otimes} R^{\text{rig}}/PR^{\text{rig}}) \otimes_{R^{\text{rig}}/PR^{\text{rig}}} \mathbf{D}_{\text{cris}}((M/P)^{\text{rig}}). \end{array}$$

(3) For any $x \in H_{\text{Iw}}^1(K_\infty, M)$, we have

$$\sigma(\mathcal{L}_{M,\xi}^G(x)) = \tau_p \cdot \mathcal{L}_{M,\xi}^G(x)$$

where σ denotes the arithmetic Frobenius.

By abuse of notation we also write $\mathcal{L}_{M,\xi}^G$ for the induced map with source $(\mathcal{H}_{\tilde{L}}(G) \widehat{\otimes} R^{\text{rig}}) \otimes H_{\text{Iw}}^1(K_\infty, M)$.

Proof. The proof of Thm. 4.7 in (loc. cit.) carries over almost literally to show the first part apart from the injectivity statement, which is proved in the same way as Prop. 4.12 in (loc. cit.).

The commutativity of the diagram in 2 is just the fact that the construction of $\mathcal{L}_{M,\xi}^G$ is functorial in the coefficients of M : More precisely, we have to check that the following diagrams are commutative.

HIwN

$$(8) \quad \begin{array}{ccc} H_{\text{Iw}}^1(K_\infty, M) \otimes_R R/P & \xrightarrow{\sim} & \mathbf{N}_\infty(M)^{\psi=1} \otimes_R R/P \\ \downarrow & & \downarrow \\ H_{\text{Iw}}^1(K_\infty, M \otimes_R R/P) & \xrightarrow{\sim} & \mathbf{N}_\infty(M \otimes_R R/P)^{\psi=1} \end{array}$$

(where the right vertical map is induced from the natural base change map $N(M) \otimes_R R/P \rightarrow N(M/PM)$ from section 3 and the R -linearity of ψ while the left vertical map comes from the obvious base change spectral sequence using the flatness of M over R , see e.g. [Nek06, §8.4.8.3] or (2), (3)),

NtoNpsi=0

$$(9) \quad \begin{array}{ccc} \mathbf{N}_\infty(M)^{\psi=1} \otimes_R R/P & \xrightarrow{1-\varphi} & (\varphi^* \mathbf{N}_\infty(M))^{\psi=0} \otimes_R R/P \\ \downarrow & & \downarrow \\ \mathbf{N}_\infty(M \otimes_R R/P)^{\psi=1} & \xrightarrow{1-\varphi} & (\varphi^* \mathbf{N}_\infty(M \otimes_R R/P))^{\psi=0} \end{array}$$

and

Npsi=0Dcris

$$(10) \quad \begin{array}{ccc} (\varphi^* \mathbf{N}_\infty(M))^{\psi=0} \otimes_R R/P & \hookrightarrow & (\mathcal{H}_{\tilde{L}}(G) \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) \otimes_R R/P \\ \downarrow & & \downarrow \\ (\varphi^* \mathbf{N}_\infty(M \otimes_R R/P))^{\psi=0} & \hookrightarrow & (\mathcal{H}_{\tilde{L}}(G) \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}((M \otimes_R R/P)^{\text{rig}}) \end{array}$$

The commutativity of diagram (8) follows from the functoriality of the Fontaine isomorphism. For diagram (9), commutativity follows immediately from the R -linearity of φ . The commutativity of (10) follows from the commutativity of the diagram

$$\begin{array}{ccc} (\varphi^* \mathbf{N}(M))^{\psi=0} \otimes_R R/P & \longrightarrow & (\mathcal{H}_{\mathbf{Q}_p}(\Gamma) \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) \otimes_R R/P \\ \downarrow & & \downarrow \cong \\ (\varphi^* \mathbf{N}(M \otimes_R R/P))^{\psi=0} & \longrightarrow & (\mathcal{H}_{\mathbf{Q}_p}(\Gamma) \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}((M \otimes_R R/P)^{\text{rig}}) \end{array}$$

But this follows from the functoriality of the maps $\mathbf{N}(M) \rightarrow \mathbf{D}_{\text{rig}}^\dagger(M^{\text{rig}})$ and $\mathbf{D}_{\text{cris}}(M^{\text{rig}}) \rightarrow \mathbf{D}_{\text{rig}}^\dagger(M^{\text{rig}})$, which is clear.

The third part is clear from the construction. \square

7. CONSTRUCTION OF THE ISOMORPHISM

The aim of this section is as follows. Let M be a crystalline R -linear Galois representation with R a complete local noetherian \mathbf{Z}_p -algebra with finite residue field, which is Cohen–Macaulay, normal, and flat over \mathbf{Z}_p , such that R^{rig} is an integral domain. Let $\tilde{R} := \widehat{\mathbf{Z}_p^{\text{nr}}} \widehat{\otimes}_{\mathbf{Z}_p} R$ and let

f:tilde

$$(11) \quad \Lambda_R(G) := \Lambda_{\mathbf{Z}_p}(G) \widehat{\otimes}_{\mathbf{Z}_p} R \quad \text{and} \quad \Lambda_{\tilde{R}}(G) := \Lambda_{\mathbf{Z}_p}(G) \widehat{\otimes}_{\mathbf{Z}_p} \tilde{R}.$$

We shall construct a canonical isomorphism of determinants over the ring $\Lambda_{R,\mathbf{Q}_p}(G) := \Lambda_{\tilde{R}}(G)[1/p]$:

$$\Theta_{\Lambda_{\tilde{R},\mathbf{Q}_p}(G),\xi}(T) : \text{Det}_{\Lambda_{\tilde{R},\mathbf{Q}_p}(G)}(0) \xrightarrow{\sim} \Lambda_{\tilde{R},\mathbf{Q}_p}(G) \otimes_{\Lambda_R(G)} \left\{ \text{Det}_{\Lambda_R(G)} R\Gamma_{\text{Iw}}(K_\infty, M) \cdot \text{Det}_{\Lambda_R(G)} (\Lambda_R(G) \otimes_R \mathbf{D}_{\text{cris}}(M)) \right\},$$

where $\mathbf{D}_{\text{cris}}(M) = N(M)/\pi N(M)$.

Our construction is to define the isomorphism over the total quotient ring $\mathcal{K}_{\tilde{R}}(G)$ of the distribution algebra $\mathcal{H}_{\tilde{R}}(G) := \mathcal{H}_{\mathbf{Q}_p}(G) \widehat{\otimes} \tilde{R}^{\text{rig}}$, and then descend it to $\Lambda_{\tilde{R}, \mathbf{Q}_p}(G)$.

In order to carry out our calculations, we will need to specialize at closed points of $\text{Spec } R[1/p]$ and utilize the results of [LVZ15]. However, as we are interested in finite modules over $\Lambda_{\tilde{R}}(G)[1/p]$, which has many more closed points than $\Lambda_R(G)[1/p]$, we record the following useful lemmas:

Lemma 7.1. *The natural map $\tilde{R}[1/p] \rightarrow \prod_x \widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} K_x$ is injective, where the product runs over closed points of $\text{Spec } R[1/p]$ and K_x denotes the residue field at x .*

Proof. According to Appendix A.1 we can exhaust the generic fiber $\text{Spf}(R)^{\text{rig}}$ of $\text{Spf } R$, whose points corresponds to the closed points of $\text{Spec } R[1/p]$, by affinoid subdomains $\text{Sp } A_m$, which are reduced since by one of our running hypotheses $R[1/p]$ is reduced. Since $\widehat{\mathbf{Z}}_p^{\text{nr}}$ is unramified over \mathbf{Z}_p , the maximal ideal of definition I of R generates the maximal ideal of definition of $\widehat{\mathbf{Z}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Z}_p} R$. For every $m \geq 0$, there is a natural map

$$\widehat{\mathbf{Z}}_p^{\text{nr}} \otimes_{\mathbf{Z}_p} (R[I^m/p])^\wedge \rightarrow (\widehat{\mathbf{Z}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Z}_p} R)[I^m/p]$$

which is an isomorphism after p -adically completing (since it is an isomorphism modulo every power of p).

Thus, we obtain a map

$$\tilde{R}[\frac{1}{p}] \hookrightarrow \tilde{R}^{\text{rig}} \cong \varprojlim_m (\tilde{R}[I^m/p])^\wedge [1/p] \cong \varprojlim_m (\widehat{\mathbf{Z}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Z}_p} (R[I^m/p])^\wedge) [1/p] \cong \varprojlim_m \widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} A_m$$

So it is enough to consider the analogous map $\widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes} A \rightarrow \prod_x \widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes} K_x$ when A is a reduced \mathbf{Q}_p -affinoid algebra and x runs over the points of $\text{Sp}(A)$. Then the subring $A^\circ \subset A$ of power-bounded elements is topologically finite type over \mathbf{Z}_p , and its reduction modulo p is a finite type \mathbf{F}_p -algebra.

If A is a Tate algebra, this is clear, since points of $\text{Sp}(\widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} T_n)$ lying over points of $\text{Sp}(T_n)$ are dense (with respect to the canonical topology), since $\text{Sp}(T_n)$ is a polydisk and \mathbf{Q}_p^{nr} is dense in $\widehat{\mathbf{Q}}_p^{\text{nr}}$. To treat the general case, we may assume that A is irreducible. Then Noether normalization implies that there is a finite (torsion-free) monomorphism $T_n \rightarrow A$, and since \mathbf{Q}_p is discretely valued, [BGR84, Corollary 6.4.1/6] implies that the induced map $T_n^\circ \rightarrow A^\circ$ on subrings of power-bounded elements is also finite.

Let $f \in \widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} A$ be a function such that for every homomorphism $T_n \rightarrow L$, L/\mathbf{Q}_p a finite extension, the image of f in $L' \otimes_{T_n} (\widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} A)$ is trivial; since every point of $\text{Sp}(A)$ lies over a point of $\text{Sp}(T_n)$, this holds for every f satisfying the hypotheses of this lemma. We wish to show that $f = 0$. After multiplying by a power of p , we may assume that $f \in \widehat{\mathbf{Z}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Z}_p} A^\circ$. Consider a point $x : \widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} T_n \rightarrow L$, where L is a finite extension of $\widehat{\mathbf{Q}}_p^{\text{nr}}$. Under this homomorphism, $\widehat{\mathbf{Z}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Z}_p} T_n^\circ$ is carried to the ring of integers \mathcal{O}_L of L . For each integer $n \geq 0$, we choose another homomorphism $x_n : T_n \rightarrow \overline{\mathbf{Q}}_p$ such that $x|_{\widehat{\mathbf{Z}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Z}_p} T_n^\circ} \equiv \widehat{\mathbf{Z}}_p^{\text{nr}} \otimes_{\mathbf{Z}_p} x_n|_{T_n^\circ} \pmod{p^n}$. Thus, $x(f) \equiv 0 \pmod{p^n}$ for all n . Since $\widehat{\mathbf{Z}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Z}_p} A^\circ$ is p -adically separated and complete, $x(f) = 0$. Since $\widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} A$ is a finite $\widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} T_n$ -module and $\widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} T_n$ is reduced, we are done. \square

Lemma 7.2. *Let A be a reduced \mathbf{Q}_p -affinoid algebra, and suppose that for every $x : A \rightarrow L$, L/\mathbf{Q}_p a finite extension with ring of integers \mathcal{O}_L , the induced map $\widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} A \rightarrow \widehat{\mathbf{Q}}_p^{\text{nr}} \otimes_{\mathbf{Q}_p} L$ carries $f \in \widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} A$ to $\widehat{\mathbf{Z}}_p^{\text{nr}} \otimes_{\mathbf{Z}_p} \mathcal{O}_L$. Then $|f(x)| \leq 1$ for all points $x \in \text{Sp}(\widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} A)$.*

Proof. If $A = T_n$, the result is clear because points of $\text{Sp}(\widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} T_n)$ lying over points of $\text{Sp}(T_n)$ are dense with respect to the canonical topology. To treat the general case, we will again assume $\text{Sp}(A)$ is irreducible and use Noether normalization to find finite torsion-free monomorphisms $T_n \rightarrow A$ and $\widehat{\mathbf{Z}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Z}_p} T_n^\circ \rightarrow \widehat{\mathbf{Z}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Z}_p} A^\circ$.

Since the supremum norm on A and $\widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} A$ is power-multiplicative, we may replace f by a power of itself and assume that $|f|_{\text{sup}} \in |\mathbf{Q}_p|$.

Suppose that $f \notin \widehat{\mathbf{Z}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Z}_p} A^\circ$. Then there is some $k > 0$ such that $p^k f \in \widehat{\mathbf{Z}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Z}_p} A^\circ$ and $|p^k f(x)| \leq p^{-1}$ for every point x lying over a point of $\text{Sp}(A)$. Choose the minimal such k . The first condition

implies that $|p^k f|_{\text{sup}} \leq 1$, and the minimality of k implies that $|p^k f|_{\text{sup}} > p^{-1}$. Indeed, if $\{e_i\}$ is a basis for $\widehat{\mathbf{F}}_p$ over \mathbf{F}_p , then the Teichmüller lifts $\{[e_i]\}$ form a basis for $\widehat{\mathbf{Q}}_p^{\text{nr}}$ over \mathbf{Q}_p , and we may write $p^k f$ uniquely as $p^k f = \sum_i f_i [e_i]$ with $f_i \in A^\circ$. If $|p^k f|_{\text{sup}} \leq p^{-1}$, then for every point $x \in \text{Sp}(A)$ the reduction $\overline{p^k f(x)} = \sum_i \overline{e_i f_i(x)}$ modulo p is 0. But since the e_i are linearly independent, we see that $\overline{f_i(x)} = 0$ for all i , and therefore $f_i \in pA^\circ$ for all i , contradicting the minimality of k .

The maximum modulus principal implies that f actually attains its supremum at a point of $\text{Sp}(\widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} A)$, which lies over a point $x \in \text{Sp}(\widehat{\mathbf{Q}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Q}_p} T_n)$. As in the proof of the previous lemma, we may p -adically approximate x by a series of points $x_n \in \text{Sp}(T_n)$. As before, $x(p^k f) \equiv (\widehat{\mathbf{Z}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Z}_p} x_n)(p^k f) \pmod{p^n}$. But $|x(p^k f)|_{\text{sup}} = 1$ (because $|p^k f|_{\text{sup}} = 1$) whereas $|(\widehat{\mathbf{Z}}_p^{\text{nr}} \widehat{\otimes}_{\mathbf{Z}_p} x_n)(p^k f)|_{\text{sup}} \leq p^{-1}$ (by our assumption on f at points over points of $\text{Sp}(A)$), which is a contradiction. Thus, $k = 0$ and we are done. \square

We can think of these lemmas as saying that “classical points, i.e., those lying above points of $\text{Spf}(R)^{\text{rig}}$, are dense in $\text{Spf}(\widetilde{R})^{\text{rig}}$ ”.

7.1. Construction of Θ over $\mathcal{K}_{\widetilde{R}}(G)$. Over $\mathcal{K}_{\widetilde{R}}(G)$, the construction of the isomorphism Θ is very simple:

Proposition 7.3. *The regulator $\mathcal{L}_{M,\xi}^G$ induces an isomorphism*

$$\text{Det}_{\mathcal{K}_{\widetilde{R}}(G)}(0) \xrightarrow{\sim} \text{Det}_{\mathcal{K}_{\widetilde{R}}(G)}((\mathcal{K}_{\widetilde{R}}(G)) \otimes_{\Lambda_R(G)} R\Gamma_{\text{Iw}}(K_\infty, M)) \cdot \text{Det}_{\mathcal{K}_{\widetilde{R}}(G)}((\mathcal{K}_{\widetilde{R}}(G)) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})).$$

Proof. By Proposition 5.3 and property B.h) in Appendix B,

$$\text{Det}_{\Lambda_{\mathcal{O},R}(G)}(R\Gamma_{\text{Iw}}(K_\infty, M)) \cong \text{Det}_{\Lambda_{\mathcal{O},R}(G)}(\mathbf{H}_{\text{Iw}}^1(K_\infty, M))^{-1}$$

up to $\Lambda_{\mathbf{Z}_p}(G)$ -torsion. We therefore consider the map

$$\mathcal{L}_{M,\xi}^G : \mathcal{H}_{\widetilde{R}}(G) \otimes_{\Lambda_R(G)} \mathbf{H}_{\text{Iw}}^1(K_\infty, M) \rightarrow \mathcal{H}_{\widetilde{R}}(G) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}).$$

This map is defined by base extension from a map $\mathbf{H}_{\text{Iw}}^1(K_\infty, M) \rightarrow (\mathcal{H}_{\mathbf{Q}_p}(G)_{\tau_p} \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})$; choosing an $\mathcal{H}_{\mathbf{Q}_p}(G)$ -basis of $\mathcal{H}_{\mathbf{Q}_p}(G)_{\tau_p}$, it suffices to show that the map of $\mathcal{H}_{\mathbf{Q}_p}(G) \widehat{\otimes} R^{\text{rig}}$ -modules

$$\mathcal{L}_{M,\xi}^G : \mathcal{H}_R(G) \otimes_{\Lambda_R(G)} \mathbf{H}_{\text{Iw}}^1(K_\infty, M) \rightarrow (\mathcal{H}_{\mathbf{Q}_p}(G)_{\tau_p} \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})$$

is generically an isomorphism.

Since $\mathbf{H}_{\text{Iw}}^1(K_\infty, M)$ is generically free of rank d and $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$ is projective of rank d , it suffices to prove that this map is generically surjective. But after specializing at any closed point x of $\text{Spf}(R)^{\text{rig}}$, [LVZ15, Proposition 3.4.1] implies that the image of the regulator map contains a basis of $\text{Frac } \mathcal{H}_{\mathbf{Q}_p}(G) \otimes_{\mathbf{Q}_p} \mathbf{D}_{\text{cris}}(M_x)$, so generic surjectivity follows. \square

Definition 7.4. Let $\Theta_{\mathcal{K}_{\widetilde{R}}(G),\xi}(M)$ be the isomorphism

$$\text{Det}_{\mathcal{K}_{\widetilde{R}}(G)}(0) \xrightarrow{\sim} \text{Det}_{\mathcal{K}_{\widetilde{R}}(G)}((\mathcal{K}_{\widetilde{R}}(G)) \otimes_{\Lambda_R(G)} R\Gamma_{\text{Iw}}(K_\infty, M)) \cdot \text{Det}_{\mathcal{K}_{\widetilde{R}}(G)}((\mathcal{K}_{\widetilde{R}}(G)) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}))$$

given by

$$\Theta_{\mathcal{K}_{\widetilde{R}}(G),\xi}(M) = \ell(M)^{-1} \text{Det}(\mathcal{L}_{M,\xi}^G)$$

where $\ell(M) \in \mathcal{H}_{\mathbf{Q}_p}(G)$ is defined below.

Definition 7.5.

(1) Let $\gamma \in \Gamma$ be any non-torsion element. Then we define

$$\ell_j := \frac{\log \gamma}{\log \chi(\gamma)} - j$$

(2) For $n \in \mathbf{Z}$, define the element $\mu_n \in \text{Frac } \mathcal{H}_{\mathbf{Q}_p}(\Gamma)$ by

$$\mu_n = \begin{cases} \ell_0 \cdots \ell_{n-1} & \text{if } n \geq 1 \\ 1 & \text{if } n = 0 \\ (\ell_{-1} \cdots \ell_n)^{-1} & \text{if } n \leq -1. \end{cases}$$

(3) For V a Hodge–Tate representation of $G_{\mathbf{Q}_p}$, with Hodge–Tate weights n_1, \dots, n_d , let

$$\ell(V) = \prod_{i=1}^d \mu_{n_i}.$$

The p -adic Hodge type of M is locally constant on $\mathrm{Spf}(R)^{\mathrm{rig}}$. Since we assume that $R[1/p]$ is integral, the Hodge–Tate weights are constant. If the Hodge–Tate weights are n_1, \dots, n_d , we set $\ell(M) := \prod_{i=1}^d \mu_{n_i}$. This is an element of $\mathcal{H}_{\mathbf{Q}_p}(G) \subset \mathcal{H}_{\bar{R}}(G)$.

Since $\mathbf{D}_{\mathrm{cris}}(M^{\mathrm{rig}}) = R^{\mathrm{rig}} \otimes_R \mathbf{D}_{\mathrm{cris}}(M)$ by Corollary 4.10, we can rewrite the above isomorphism as

$$\mathrm{Det}_{\mathcal{K}_{\bar{R}}(G)}(0) \xrightarrow{\sim} \mathcal{K}_{\bar{R}}(G) \otimes_{\Lambda_R(G)} \left\{ \mathrm{Det}_{\Lambda_R(G)}(R\Gamma_{\mathrm{Iw}}(K_\infty, M)) \cdot \mathrm{Det}_{\Lambda_R(G)}(\Lambda_R(G) \otimes_R \mathbf{D}_{\mathrm{cris}}(M)) \right\}$$

7.2. Definition of Θ over $\Lambda_R(G)[1/p]$. We wish to show that the isomorphism $\Theta_{\mathcal{K}_{\bar{R}}(G), \xi}(M)$ can be defined over $\Lambda_{\bar{R}, \mathbf{Q}_p}(G)$. More precisely, we want to show the following:

Proposition 7.6. *There is a trivialization of line bundles*

$$\Theta_{\Lambda_{R, \mathbf{Q}_p}(G), \xi} : \mathrm{Det}_{\Lambda_R(G)[1/p]}(0) \xrightarrow{\sim} \mathrm{Det}_{\Lambda_{\bar{R}, \mathbf{Q}_p}(G)}(R\Gamma_{\mathrm{Iw}}(K_\infty, M)[1/p]) \cdot \mathrm{Det}_{\Lambda_{\bar{R}, \mathbf{Q}_p}(G)}\left(\Lambda_{\bar{R}, \mathbf{Q}_p}(G) \otimes_{R[1/p]} \mathbf{D}_{\mathrm{cris}}(M)[1/p]\right)$$

which agrees with $\Theta_{\mathcal{K}_{\bar{R}}(G), \xi}(M)$ after extending scalars. Moreover, for every specialization $R[1/p] \rightarrow L$, the base change $\Theta_{\Lambda_{R, \mathbf{Q}_p}(G), \xi} \otimes_{R[1/p]} L$ agrees with the trivialization

$$\Theta_{\Lambda_L(G), \xi} : \mathrm{Det}_{\Lambda_L(G)}(0) \rightarrow \mathrm{Det}_{\Lambda_L(G)}\left(\tilde{L} \otimes_L R\Gamma_{\mathrm{Iw}}(K_\infty, M_L)\right) \cdot \mathrm{Det}_{\Lambda_L(G)}\left(\Lambda_L(G) \otimes_L \mathbf{D}_{\mathrm{cris}}(M_L)\right)$$

defined in [LVZ15, Theorem 4.2.1].

We first recall that $\Theta_{\mathcal{K}_{\bar{R}}(G), \xi}(M)$ is defined via base extension from an isomorphism over the total ring of fractions of $\mathcal{H}_{\mathbf{Q}_p}(G)_{\tau_p} \widehat{\otimes} R^{\mathrm{rig}}$. We therefore descend the latter isomorphism to an isomorphism over $\Lambda_{\mathbf{Q}_p}(G)_{\tau_p} \widehat{\otimes} R[1/p]$ (using the fact that $\Lambda_{\mathbf{Q}_p}(G)_{\tau_p}$ is free of rank 1 over $\Lambda_{\mathbf{Q}_p}(G)$).

To construct a homomorphism of line bundles

$$\mathrm{Det}_{\Lambda_{\bar{R}, \mathbf{Q}_p}(G)} R\Gamma(K_\infty, M)[1/p]^{-1} \rightarrow \mathrm{Det}_{\Lambda_{\bar{R}, \mathbf{Q}_p}(G)}\left(\Lambda_{\bar{R}, \mathbf{Q}_p}(G) \otimes_{R[1/p]} \mathbf{D}_{\mathrm{cris}}(M)[1/p]\right)$$

it suffices to construct homomorphisms on an affine cover and check that they agree on overlaps. Thus, our strategy is to work locally on $\mathrm{Spec} \Lambda_{\bar{R}, \mathbf{Q}_p}(G)$ (to trivialize $\mathrm{Det}_{\Lambda_{\bar{R}, \mathbf{Q}_p}(G)}(R\Gamma(K_\infty, M)[1/p])$ and $\mathrm{Det}_{\Lambda_{\bar{R}, \mathbf{Q}_p}(G)}\left(\Lambda_{\bar{R}, \mathbf{Q}_p}(G) \otimes_{R[1/p]} \mathbf{D}_{\mathrm{cris}}(M)[1/p]\right)$), choose integral generators on both sides, and calculate that the determinant actually lands in $\Lambda_{\bar{R}, \mathbf{Q}_p}(G)$.

However, we will have to be a bit careful, for two reasons: First, our previous computations concerned $\mathrm{Det}_{\Lambda_{\bar{R}, \mathbf{Q}_p}(G)} H_{\mathrm{Iw}}^1(K_\infty, M)[1/p]$, not $\mathrm{Det}_{\Lambda_{\bar{R}, \mathbf{Q}_p}(G)}(R\Gamma_{\mathrm{Iw}}(K_\infty, M)[1/p])^{-1}$ and they are only canonically isomorphic when $H_{\mathrm{Iw}}^2(K_\infty, M)[1/p] = 0$. Second, we can only compute with $\mathrm{Det}_{\Lambda_{\bar{R}, \mathbf{Q}_p}(G)} H_{\mathrm{Iw}}^1(K_\infty, M)$, and it is only compatible with base change on R when $H_{\mathrm{Iw}}^2(K_\infty, M) = 0$. Thus, we will first work away from the support of $H_{\mathrm{Iw}}^2(K_\infty, M)$ and then use the Cohen–Macaulay-ness of R to extend over this locus.

First of all, $\det \mathcal{L}_{M, \xi}^G$ is defined by a homomorphism between modules over $\mathcal{H}_{\bar{R}}(G)$. If we can show that the valuation of $\ell(V)^{-1} \det \mathcal{L}_{M, \xi}^G$ is bounded (with respect to suitable choices of local generators), then normality of $\Lambda_{\bar{R}}(G)$ will imply that it actually lives in $\Lambda_{\bar{R}, \mathbf{Q}_p}(G)$, as desired. We therefore study specializations of $\ell(V)^{-1} \det \mathcal{L}_{M, \xi}^G$.

We first work over open affine subspaces $U = \mathrm{Spec} A_U \subset \Lambda_{\bar{R}}(G)$ where $H_{\mathrm{Iw}}^2(K_\infty, M)$ vanishes; on such subspaces, $\mathrm{Det}_{A_U} R\Gamma_{\mathrm{Iw}}(K_\infty, M) \otimes_{R A_U} \cong \mathrm{Det}_{A_U} (H_{\mathrm{Iw}}^1(K_\infty, M))^{-1} \otimes_{R A_U}$ by property B.h) in Appendix B.

Over these subspaces, we will consider specializations of the determinant of $\mathcal{L}_{M, \xi}^G$ along maps $R[1/p] \rightarrow L$, where L/\mathbf{Q}_p is a finite field extension. A priori, such base changes involve derived tensor products and thus higher Tor groups. However, Theorem 6.1 (2) together with the spectral sequence (2) (for $Y = L$) shows that when $H_{\mathrm{Iw}}^2 = 0$, these Tor groups vanish.

In general, we will use (4) to check that no higher Tor-groups are involved in the descent calculation. Indeed, for a Cohen–Macaulay ring pseudo-null modules possess a canonical trivialisation by the same proof as for [Ven13, Lemma 2.2]. From this comment and by construction below, it will be clear that — once $\Theta_{\Lambda_{R, \mathbf{Q}_p}(G), \xi}$ exists — it satisfies the desired compatibility with specialisations.

Lemma 7.7. *Let $U = \text{Spec } A_U \subset \text{Spec } \Lambda_R(G)$ be an open affine subspace where $H_{\text{Iw}}^2(K_\infty, M) = 0$ and $\text{Det}_{\Lambda_R(G)}(H_{\text{Iw}}^1(K_\infty, M))$ and $\text{Det}_{\Lambda_R(G)}(\Lambda_R(G) \otimes_R \mathbf{D}_{\text{cris}}(M))$ are trivial. Then $\ell(V)^{-1} \det \mathcal{L}_{M,\xi}^G$ can be defined over the pre-image of U in $\text{Spec } \Lambda_{\bar{R}}(G)$.*

Proof. Write $A_U = \Lambda_R(G)[1/f_U]$ for some $f_U \in \Lambda_R(G)$. Choosing generators of $\text{Det}_{\Lambda_{\bar{R}}(G)}(H_{\text{Iw}}^1(K_\infty, M))|_U$ and $\text{Det}_{\Lambda_{\bar{R}}(G)}(\Lambda_{\bar{R}}(G) \otimes_R \mathbf{D}_{\text{cris}}(M))|_U$, we may view $\ell(V)^{-1} \det \mathcal{L}_{M,\xi}^G$ as multiplication by an element $a_U := a'_U/f_U^k$, where $a'_U \in \mathcal{H}_{\bar{R}}(G)$. In fact, since $\mathcal{L}_{M,\xi}^G$ is the base extension of a morphism of $\mathcal{H}_R(G)$ -modules

$$\mathcal{H}_R(G) \otimes_{\Lambda_R(G)} H_{\text{Iw}}^1(K_\infty, M) \rightarrow (\mathcal{H}_{\mathbf{Q}_p}(G)_{\tau_p} \widehat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}})$$

of the same rank, we may assume that $a'_U \in \mathcal{H}_R(G)$.

The formation of $H_{\text{Iw}}^1(K_\infty, M)$ commutes with specialization on U , since $H_{\text{Iw}}^2(K_\infty, M)|_U = 0$ and $R\Gamma_{\text{Iw}}(K_\infty, M)$ does. Let L/\mathbf{Q}_p be a finite extension with ring of integers \mathcal{O}_L , and let $x : R[1/p] \rightarrow L$ be a closed point of $\text{Spec } R[1/p]$ such that $A_U[1/p] \otimes_{R[1/p]} L$ is non-zero. Then the natural homomorphism $R \rightarrow \mathcal{O}_L$ induces a map $\mathbf{D}_{\text{cris}}(M) \rightarrow \mathbf{D}_{\text{cris}}(M \otimes_R \mathcal{O}_L)$ with cokernel annihilated by $p^{\alpha(h)}$. Therefore, there is a natural homomorphism $\text{Det}_R \mathbf{D}_{\text{cris}}(M) \rightarrow \text{Det}_{\mathcal{O}_L} \mathbf{D}_{\text{cris}}(M \otimes_R \mathcal{O}_L)$ with cokernel annihilated by $p^{d\alpha(h)}$. Now [LVZ15, Corollary 4.3.8] implies that the specialization of $\ell(V)^{-1} \det \mathcal{L}_{M,\xi}^G$ at x is given by multiplication by an element of $\Lambda_{\bar{\mathcal{O}}_L}(G)$ with respect to generators of $H_{\text{Iw}}^1(K_\infty, M \otimes_R \mathcal{O}_L)$ and $\mathbf{D}_{\text{cris}}(M \otimes_R \mathcal{O}_L)$.

This implies that for every such x , $p^{d\alpha(h)} f_U(x)^k a_U(x) \in \Lambda_{\bar{\mathcal{O}}_L}(G)$. But if x is a closed point of $\text{Spec } R[1/p]$ where $A_U[1/p] \otimes_{R[1/p]} L$ is zero, then $f_U(x) = 0$ and $p^{d\alpha(h)} f_U(x)^k a_U(x) \in \Lambda_{\bar{\mathcal{O}}_L}(G)$. Thus, for every point $x' \in \text{Spf}(\Lambda_R(G))^{\text{rig}}$,

$$|p^{d\alpha(h)} a'_U(x')| = |p^{d\alpha(h)} f_U(x')^k a_U(x')| \leq 1$$

Now by Lemma 7.2 (and exhausting $\text{Spf}(R)^{\text{rig}}$ by affinoid subdomains)

$$|p^{d\alpha(h)} a'_U(x')| = |p^{d\alpha(h)} f_U(x')^k a_U(x')| \leq 1$$

for all points $x \in \text{Spf}(\Lambda_{\bar{R}}(G))^{\text{rig}}$. Since R is assumed normal, Proposition [dJ95, Proposition 7.3.6] implies that $a'_U \in \Lambda_{\bar{R}, \mathbf{Q}_p}(G)$.

It remains to check that $\ell(V)^{-1} \det \mathcal{L}_{M,\xi}^G|_{U[1/p]}$ is an isomorphism, i.e., that a_U has no zeroes. But by construction, $\ell(V)^{-1} \det \mathcal{L}_{M,\xi}^G|_{U[1/p]}$ is compatible with specialization on $R[1/p]$, so comparison with the classical case shows that it cannot vanish. \square

If $U, U' \subset \text{Spec } \Lambda_{\bar{R}}(G)$ are open affine subspaces as above, then $\ell(V)^{-1} \det \mathcal{L}_{M,\xi}^G|_U$ and $\ell(V)^{-1} \det \mathcal{L}_{M,\xi}^G|_{U'}$ agree on $U \cap U'$, because they agree after extending scalars to $\mathcal{K}_{\bar{R}}(G)$.

Lemma 7.8. *Let $U = \text{Spec } A_U \subset \text{Spec } \Lambda_R(G)$ be an open affine subspace where $\text{Det}_{\Lambda_R(G)}(R\Gamma_{\text{Iw}}(K_\infty, M))$ and $\text{Det}_{\Lambda_R(G)}(\Lambda_R(G) \otimes_R \mathbf{D}_{\text{cris}}(M))$ are trivial. Then $\ell(V)^{-1} \det \mathcal{L}_{M,\xi}^G$ can be defined over the pre-image $\tilde{U} = \text{Spec } A_{\tilde{U}}$ of U in $\text{Spec } \Lambda_{\bar{R}}(G)$.*

Proof. Choose bases of $\text{Det}_{\Lambda_R(G)}(R\Gamma_{\text{Iw}}(K_\infty, M))^{-1}$ and $\text{Det}_{\Lambda_R(G)}(\Lambda_R(G) \otimes_R \mathbf{D}_{\text{cris}}(M))$ as in the proof of the previous lemma; an isomorphism

$$\text{Det}_{A_{\tilde{U}}[1/p]}(A_{\tilde{U}}[1/p] \otimes_{\Lambda_R(G)} R\Gamma_{\text{Iw}}(K_\infty, M))^{-1} \xrightarrow{\sim} \text{Det}_{A_{\tilde{U}}[1/p]}(A_{\tilde{U}}[1/p] \otimes_R \mathbf{D}_{\text{cris}}(M))$$

corresponds to multiplication by a unit of $A_{\tilde{U}}[1/p]$. The support of $H_{\text{Iw}}^2(K_\infty, M)$ is a closed subscheme $V \subset U$ of codimension at least 2; for each open affine subspace $U' = \text{Spec } A_{U'} \subset U \setminus V$ we obtain a unit $a_{U'} \in A_{U'}[1/p]$ by Lemma 7.7 and these units agree on overlaps. Therefore, the $a_{U'}$ glue to a section a'_U on all of $(U \setminus V)[1/p]$. But $\text{Spec } \Lambda_{\bar{R}}(G)$ is Cohen–Macaulay, so a'_U extends to a section a_U over all of $U[1/p]$.

It remains to check that $a_U \in A_U[1/p]^\times$. Let L/\mathbf{Q}_p be a finite extension with ring of integers \mathcal{O}_L , and let $x : R[1/p] \rightarrow L$ be a closed point of $\text{Spec } R[1/p]$ such that $A_U[1/p] \otimes_{R[1/p]} L$ is non-zero. Then we know that the image of a_U in $A_U[1/p] \otimes_{R[1/p]} L$ extends to a unit of $\Lambda_{\bar{L}}(G)$ by the classical case, since the formation of $\text{Det}_{\Lambda_R(G)}(R\Gamma_{\text{Iw}}(K_\infty, M))^{-1}$ commutes with specialization on R and $\mathbf{D}_{\text{cris}}(M)[1/p]$ commutes with specialization on $R[1/p]$. Thus the claim follows from Lemma 7.14. \square

It is again clear that if $U, U' \subset \text{Spec } \Lambda_{\bar{R}}(G)$ are open affine subspaces as above, then $\ell(V)^{-1} \det \mathcal{L}_{M,\xi}^G|_U$ and $\ell(V)^{-1} \det \mathcal{L}_{M,\xi}^G|_{U'}$ agree on $U \cap U'$. This proves Proposition 7.6.

Remark 7.9. Although the proofs of Lemma 7.7 and Lemma 7.8 construct isomorphisms over affine subspaces of $\Lambda_R(G)[1/p]$, we only claim to have descended $\ell(M)^{-1}\mathcal{L}_{M,\xi}^G$ to $\Lambda_{\tilde{R}}(G)[1/p]$. This is because the proofs implicitly involved making a non-canonical choice of a generator of $\Lambda_{\mathbf{Z}_p}(G)_{\tau_p}$ over $\Lambda_{\mathbf{Z}_p}(G)$.

7.3. Definition of the epsilon-isomorphism. Recall from [LVZ15, §2.3, §2.4] that if V is a d -dimensional L -linear crystalline representation of $G_{\mathbf{Q}_p}$, then the ε -factor of $\mathbf{D}_{\text{pst}}(V)$ is 1. Further, multiplication by $t^{-m(V)}\varepsilon(\mathbf{D}_{\text{pst}}(V), \xi)$ defines an isomorphism $\tilde{L} \otimes_L \text{Det}_L(\mathbf{D}_{\text{dR}}(V)) \rightarrow \tilde{L} \otimes_L \text{Det}_L V$, where $m(V) = r_1 + \dots + r_d$ is the sum of the opposites of the Hodge–Tate weights. This multiplication takes place inside the canonical isomorphism $\mathbf{B}_{\text{dR}} \otimes_{\mathbf{Q}_p} \text{Det}_L \mathbf{D}_{\text{dR}}(V) \xrightarrow{\sim} \mathbf{B}_{\text{dR}} \otimes_{\mathbf{Q}_p} \text{Det}_L V$.

If M is a crystalline family of Galois representations over R , then $\mathbf{D}_{\text{dR}}(M^{\text{rig}})$ and $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$ are well-defined and the Hodge–Tate weights are locally constant on $\text{Spf}(R)^{\text{rig}}$.

Theorem 7.10. *Let M be a crystalline family of Galois representations over R . Then there is a unique isomorphism*

$$\varepsilon_{\tilde{R}, \mathbf{Q}_p, \xi, \text{dR}}(M) : \text{Det}_{\tilde{R}[1/p]}(\tilde{R}[1/p] \otimes_R \mathbf{D}_{\text{cris}}(M)) \xrightarrow{\sim} \text{Det}_{\tilde{R}[1/p]}(\tilde{R}[1/p] \otimes_R M)$$

whose image under specialization $x : R \rightarrow \mathcal{O}_L$ is the isomorphism defined above.

Remark 7.11. We cannot hope to construct $\varepsilon_{\tilde{R}, \xi, \text{dR}}(M)$ such that its *integral* specializations agree with the isomorphisms $\varepsilon_{\mathcal{O}_L, \xi, \text{dR}}(M)$ constructed in [LVZ15], because M commutes with base change on R but $\mathbf{D}_{\text{cris}}(M)$ does not.

Proof. By [Bel15, thm. 1.1.14] and Lemma A.6 there is a canonical isomorphism

$$\text{can}_M : (R^{\text{rig}} \hat{\otimes}_{\mathbf{Q}_p} \mathbf{B}_{\text{max}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) \cong (R^{\text{rig}} \hat{\otimes}_{\mathbf{Q}_p} \mathbf{B}_{\text{max}}) \otimes_{R^{\text{rig}}} \Gamma(\text{Spf}(R)^{\text{rig}}, M^{\text{rig}}) = (R^{\text{rig}} \hat{\otimes}_{\mathbf{Q}_p} \mathbf{B}_{\text{max}}) \otimes_R M.$$

Both sides have natural R -linear structures, given by $\mathbf{D}_{\text{cris}}(M)$ and M , respectively. We wish to show that the determinant of can_M with respect to these R -modules is an element of $t^{m(M)}\tilde{R}[1/p]^\times$. For this, we work locally on $\text{Spec } R[1/p]$ to trivialize $M[1/p]$ and $\mathbf{D}_{\text{cris}}(M)[1/p]$ and show that for any $R[1/p]$ -bases of $M[1/p]$ and of $\mathbf{D}_{\text{cris}}(M)[1/p]$, the matrix of can_M has determinant in $t^{m(M)}\tilde{R}[1/p]^\times$ (viewed as a subgroup of $\mathbf{B}_{\text{max}}^\times$).

In order to show this we follow the argument of [Ber04, Proposition V.1.2]. After twisting, we may assume that M is positive.

By Corollary 4.13 and Proposition A.14, locally on $\text{Spec } R[1/p]$ the determinant of the (injective) map $(\mathbf{B}_{\text{rig}, \mathbf{Q}_p}^+ \hat{\otimes} R^{\text{rig}}) \otimes_{R^{\text{rig}}} \mathbf{D}_{\text{cris}}(M^{\text{rig}}) \rightarrow \mathbf{N}(M)^{\text{rig}}$ (with respect to any choice of bases) is an element of $(t/\pi)^{m(M)}(R[1/p] \hat{\otimes} \mathbf{B}_{\text{rig}, F}^+)^\times = (t/\pi)^{m(M)}(R \hat{\otimes} \mathbf{A}_{\mathbf{Q}_p}^+)[1/p]^\times$.

Next, we have an inclusion $(R \hat{\otimes} \mathbf{A}^+) \otimes_{R \hat{\otimes} \mathbf{A}_{\mathbf{Q}_p}^+} \mathbf{N}(M) \subset (R \hat{\otimes} \mathbf{A}^+) \otimes_R M$. Then Corollary 4.7 tells us that after inverting p , locally on $\text{Spec } R[1/p]$ the determinant of the inclusion is the ideal $(\pi^{m(M)}) \subset (R \hat{\otimes} \mathbf{A}^+)[1/p]$.

Extending scalars to $R^{\text{rig}} \hat{\otimes} \mathbf{B}_{\text{max}}$ and composing the two maps, we see that the determinant of the canonical isomorphism can_M is an element of $t^{m(M)}(R \hat{\otimes} \mathbf{A}^+)[1/p]^\times$. We wish to show that it actually lives in $t^{m(M)}\tilde{R}[1/p]^\times$.

Since the formation of $\mathbf{D}_{\text{cris}}(M^{\text{rig}})$ is compatible with taking exterior powers, we may replace M with $\det(M)$ and consider only families of characters over R . Furthermore, by twisting, we may assume that M is unramified (so has Hodge–Tate weight 0), so that the determinant of can_M is φ -invariant up to a unit of R , and we may assume $\det(\text{can}_M) \in R \hat{\otimes} \mathbf{A}^+$. We consider the equation $\varphi(\det(\text{can}_M)) = r \cdot \det(\text{can}_M)$ modulo successive powers of \mathfrak{m}_R . For each n , there is some s such that $\varphi^s(\det(\text{can}_M)) = \det(\text{can}_M)$ modulo \mathfrak{m}_R^n ; viewing this as an equation inside $(R/\mathfrak{m}_R^n) \otimes_{\mathbf{Z}_p} \tilde{\mathbf{A}}^+$ implies that $\det(\text{can}_M) \in (R/\mathfrak{m}_R^n) \hat{\otimes} W(\mathbf{F}_{p^s})$. Passing to the limit yields the desired result. \square

Definition 7.12. We define

$$\varepsilon_{\Lambda_R, \mathbf{Q}_p}(G, \xi)(M) : \text{Det}_{\Lambda_{\tilde{R}, \mathbf{Q}_p}(G)}(0) \xrightarrow{\sim} \left[\text{Det}_{\Lambda_{\tilde{R}, \mathbf{Q}_p}(G)} \left(\Lambda_{\tilde{R}, \mathbf{Q}_p}(G) \otimes_{\Lambda_R(G)} R\Gamma_{\text{Iw}}(K_\infty, M) \right) \right] \left[\text{Det}_{\Lambda_{\tilde{R}, \mathbf{Q}_p}(G)} \left(\Lambda_{\tilde{R}, \mathbf{Q}_p}(G) \otimes_R M \right) \right]$$

to be the isomorphism given by

$$(-\gamma_{-1})^d (-1)^{m(M)} \cdot \Theta_{\Lambda_{\tilde{R}, \mathbf{Q}_p}(G), \xi}(M) \cdot \varepsilon_{\tilde{R}, \mathbf{Q}_p, \xi, \text{dR}}(M),$$

thm:Dcris-M

def:epsilon

where we regard $\varepsilon_{\tilde{R}, \mathbf{Q}_p, \xi, \text{dR}}(M)$ as an isomorphism

$$\text{Det}_{\Lambda_{\tilde{R}, \mathbf{Q}_p}(G)}(\Lambda_{\tilde{R}, \mathbf{Q}_p}(G) \otimes_R \mathbf{D}_{\text{cris}}(M)) \xrightarrow{\sim} \text{Det}_{\Lambda_{\tilde{R}, \mathbf{Q}_p}(G)}(\Lambda_{\tilde{R}, \mathbf{Q}_p}(G) \otimes_R M)$$

via base-extension.

Theorem 7.13. *There is an isomorphism of line bundles*

$$\varepsilon_{\Lambda_R(G), \xi}(M) : \text{Det}_{\Lambda_{\tilde{R}}(G)}(0) \xrightarrow{\sim} \left[\text{Det}_{\Lambda_{\tilde{R}}(G)}(\Lambda_{\tilde{R}}(G) \otimes_{\Lambda_R(G)} R\Gamma_{\text{Iw}}(K_\infty, M)) \right] \left[\text{Det}_{\Lambda_{\tilde{R}}(G)}(\Lambda_{\tilde{R}}(G) \otimes_R M) \right]$$

such that $\varepsilon_{\Lambda_R(G), \xi}(M)$ agrees with $\varepsilon_{\Lambda_{R, \mathbf{Q}_p}(G), \xi}(M)$ after inverting p . The specializations $\varepsilon_{\Lambda_R(G), \xi}(M) \otimes_R \mathcal{O}_L$ agree with the isomorphisms $\varepsilon_{\Lambda_{\mathcal{O}_L}(G), \xi}(M \otimes_R \mathcal{O}_L)$ of [LVZ15] by construction.

Proof. There is some integer m such that $p^m \varepsilon_{\Lambda_{R, \mathbf{Q}_p}(G), \xi}(M)$ is a homomorphism of line bundles over $\Lambda_{\tilde{R}}(G)$. But the construction of $\varepsilon_{\Lambda_{R, \mathbf{Q}_p}(G), \xi}(M)$ is compatible with base change $R[1/p] \rightarrow L$, and $\varepsilon_{\Lambda_{R, \mathbf{Q}_p}(G), \xi}(M) \otimes_R \mathcal{O}_L$ is an isomorphism of line bundles over $\Lambda_{\tilde{\mathcal{O}}_L}(G)$ by [LVZ15, Corollary 4.3.8]. This implies that $m = 0$ and $\varepsilon_{\Lambda_{R, \mathbf{Q}_p}(G), \xi}(M)$ is an isomorphism of line bundles over $\Lambda_{\tilde{R}}(G)$. \square

7.4. Properties of the epsilon-isomorphism. In the following we are going to apply a principle of specialization based on:

Lemma 7.14. *The canonical map*

$$K_1(\Lambda_{\tilde{R}}) \rightarrow \prod_{R \rightarrow \mathcal{O}} K_1(\Lambda_{\tilde{\mathcal{O}}}),$$

where $R \rightarrow \mathcal{O}$ runs through all \mathbf{Z}_p -algebra homomorphisms with \mathcal{O} the valuation ring of any finite extension of \mathbf{Q}_p (and such that the image has the same quotient field as \mathcal{O}) is injective.

Cp. also Corollary 7.22 below for a similar statement.

Proof. Since $SK_1(\Lambda_{\tilde{R}}) = SK_1(\Lambda_{\tilde{\mathcal{O}}}) = 1$ the statement is equivalent to the injectivity of

$$\Lambda_{\tilde{R}}^\times \rightarrow \prod_{R \rightarrow \mathcal{O}} \Lambda_{\tilde{\mathcal{O}}}^\times,$$

or - because $\bigcap_{R \rightarrow \mathcal{O}} (1 + \ker(\Lambda_{\tilde{R}} \rightarrow \Lambda_{\tilde{\mathcal{O}}})) = 1 + \bigcap_{R \rightarrow \mathcal{O}} \ker(\Lambda_{\tilde{R}} \rightarrow \Lambda_{\tilde{\mathcal{O}}})$ - to the injectivity of

$$(12) \quad \Lambda_{\tilde{R}} \rightarrow \prod_{R \rightarrow \mathcal{O}} \Lambda_{\tilde{\mathcal{O}}}.$$

Since $\Lambda_{\tilde{R}}$ is a product of rings of the type of R dealt with in Lemma 7.1 and as $\tilde{\Lambda}_R = \Lambda_{\tilde{R}}$, we obtain an injective map $\Lambda_{\tilde{R}} \hookrightarrow \prod_x \widehat{\mathbf{Q}_p^{\text{nr}}} \widehat{\otimes}_{\mathbf{Q}_p} K_x$, where the closed points x corresponds to \mathbf{Z}_p -algebra homomorphisms $\Lambda_R \rightarrow K_x$ (such that the image has quotient field K_x). One easily sees that this map factorises as

$$\begin{array}{ccc} & \prod_{R \rightarrow \mathcal{O}} \Lambda_{\tilde{\mathcal{O}}} & \\ \nearrow & & \searrow \\ \Lambda_{\tilde{R}} & \xrightarrow{\quad} & \prod_x \widehat{\mathbf{Q}_p^{\text{nr}}} \widehat{\otimes}_{\mathbf{Q}_p} K_x, \end{array}$$

because each \mathbf{Z}_p -algebra homomorphisms $\Lambda_R \rightarrow K_x$ factorizes as $\Lambda_R \rightarrow \Lambda_{\mathcal{O}_x} \rightarrow K_x$ with the first map being induced on the coefficients by a \mathbf{Z}_p -algebra homomorphisms $R \rightarrow \mathcal{O}_x$ where \mathcal{O}_x is the valuation ring of K_x . The injectivity of (12), whence the claim, follows. \square

We note for later use some properties of the isomorphisms $\varepsilon_{\Lambda_R(G), \xi}(M)$:

Proposition 7.15 (Compatibility with short exact sequences). *Let*

$$0 \longrightarrow M' \longrightarrow M \longrightarrow M'' \longrightarrow 0$$

be a short exact sequence of R -linear crystalline representations of $G_{\mathbf{Q}_p}$. Then

$$\varepsilon_{\Lambda_R(G), \xi}(M) = \varepsilon_{\Lambda_R(G), \xi}(M') \cdot \varepsilon_{\Lambda_R(G), \xi}(M'').$$

Proof. We first observe that the determinants

$$\left[\text{Det}_{\Lambda_R, \mathbf{Q}_p}(G) (R\Gamma_{\text{Iw}}(K_\infty, M)) \right] \cdot \left[\text{Det}_{\Lambda_R, \mathbf{Q}_p}(G) (R\Gamma_{\text{Iw}}(K_\infty, M')) \right]^{-1} \cdot \left[\text{Det}_{\Lambda_R, \mathbf{Q}_p}(G) (R\Gamma_{\text{Iw}}(K_\infty, M'')) \right]^{-1}$$

and

$$\left[\text{Det}_{\Lambda_{\tilde{R}, \mathbf{Q}_p}(G)} \left(\Lambda_{\tilde{R}, \mathbf{Q}_p}(G) \otimes_R M \right) \right] \cdot \left[\text{Det}_{\Lambda_{\tilde{R}, \mathbf{Q}_p}(G)} \left(\Lambda_{\tilde{R}, \mathbf{Q}_p}(G) \otimes_R M' \right) \right]^{-1} \cdot \left[\text{Det}_{\Lambda_{\tilde{R}, \mathbf{Q}_p}(G)} \left(\Lambda_{\tilde{R}, \mathbf{Q}_p}(G) \otimes_R M'' \right) \right]^{-1}.$$

have canonical trivializations, so their product (with scalars extended to $\Lambda_{\tilde{R}, \mathbf{Q}_p}(G)$) does, as well.

On the other hand, the product $\varepsilon_{\Lambda_R(G), \xi}(M) \cdot \varepsilon_{\Lambda_R(G), \xi}(M')^{-1} \cdot \varepsilon_{\Lambda_R(G), \xi}(M'')^{-1}$ provides another trivialization of the product of these determinants. But since

$$\varepsilon_{\Lambda_R(G), \xi}(\mathcal{O} \otimes_R M) = \varepsilon_{\Lambda_R(G), \xi}(\mathcal{O} \otimes_R M') \varepsilon_{\Lambda_R(G), \xi}(\mathcal{O} \otimes_R M'')$$

for every map $R \rightarrow \mathcal{O}$ where \mathcal{O} is finite flat over \mathbf{Z}_p , the proposition follows. \square

Proposition 7.16 (Change of coefficient ring). *Let R' be an R -algebra which is again a complete local noetherian \mathbf{Z}_p -algebra which is Cohen–Macaulay, normal, and \mathbf{Z}_p -flat with finite residue field. Then*

$$\varepsilon_{\Lambda_{R'}(G), \xi}(R' \otimes_R M) = R' \otimes_R \varepsilon_{\Lambda_R(G), \xi}(M).$$

Proof. Clear. \square

The next compatibility property takes a little more notation to state. For brevity let us write Λ for $\Lambda_R(G)$. For η a continuous R -valued character of G , we have a twisting homomorphism $\text{Tw}_\eta : \Lambda \rightarrow \Lambda$ which maps a group element $g \in G$ to $\eta(g)g$. Hence we obtain a pullback functor $(\text{Tw}_\eta)^*$ from the category of Λ -modules to itself:

$$(\text{Tw}_\eta)^* \mathfrak{M} := \Lambda \otimes_{\Lambda, \text{Tw}_\eta} \mathfrak{M}.$$

This can also be described in terms of tensoring with the Λ -bimodule $\Lambda \otimes_{\Lambda, \text{Tw}_\eta} \Lambda$, which is free of rank one as a Λ -module. Hence the twisting functor extends to a functor from the category $\underline{\text{Det}}(\Lambda)$ to itself, and is compatible with the functor Det .

Note that we have an isomorphism

$$(\text{Tw}_\eta)^*(\Lambda \otimes_R M) \cong \Lambda \otimes_R M(\eta^{-1}), \quad a \otimes b \otimes v \mapsto a \text{Tw}_\eta(b) \otimes (v \otimes m_{\eta^{-1}})$$

as $(\Lambda, G_{\mathbf{Q}_p})$ -modules, if Λ acts on $\Lambda \otimes_{\mathcal{O}} T$ via left multiplication on the left factor, while $g \in G_{\mathbf{Q}_p}$ sends $\lambda \otimes v$ to $\lambda \bar{g}^{-1} \otimes gv$ where \bar{g} denotes the image of g in G (and analogously for the action on $\Lambda \otimes_R M(\eta^{-1})$). Here $m_{\eta^{-1}}$ a (fixed) basis of $M(\eta^{-1})$.

We clearly have $(\text{Tw}_\eta)^* \circ (\text{Tw}_{\eta^{-1}})^* = \text{id}$. Similar definitions apply to other coefficient rings than $\Lambda_{\mathcal{O}}(G)$, including $\Lambda_L(G)$, $\mathcal{H}_L(G)$ or $\Lambda_{\tilde{R}}(G)$.

Finally note that for a Λ -module \mathfrak{M} we have a canonical isomorphism

$$\Lambda \otimes_{\Lambda, \text{Tw}_{\eta^{-1}}} \mathfrak{M} = \mathfrak{M} \otimes_R Rm_\eta, \quad \lambda \otimes m \mapsto \text{Tw}_\eta(\lambda)m \otimes m_\eta,$$

of Λ -modules, where the Λ -module structure on the right hand side is induced by the diagonal action of G upon it.

Proposition 7.17 (Invariance under crystalline twists). *If $M' = M(\eta)$ for a crystalline character η with values in R , then*

$$\text{Tw}_{\eta^{-1}}(\varepsilon_{\Lambda_R(G), \xi}(M)) \cong \varepsilon_{\Lambda_R(G), \xi}(M(\eta))$$

Proof. This again follows by noting that $\text{Tw}_{\eta^{-1}}(\varepsilon_{\Lambda_R(G), \xi}(M))$ and $\varepsilon_{\Lambda_R(G), \xi}(M(\eta))$ both provide trivializations of

$$\left[\text{Det}_{\Lambda_{\tilde{R}, \mathbf{Q}_p}(G)} \left(\Lambda_{\tilde{R}, \mathbf{Q}_p}(G) \otimes_{\Lambda_R(G)} R\Gamma_{\text{Iw}}(K_\infty, M(\eta)) \right) \right] \left[\text{Det}_{\Lambda_{\tilde{R}, \mathbf{Q}_p}(G)} \left(\Lambda_{\tilde{R}, \mathbf{Q}_p}(G) \otimes_R M(\eta) \right) \right]$$

Since they agree after extending scalars along $R \rightarrow \mathcal{O}$ when \mathcal{O} is finite flat over \mathbf{Z}_p , they are the same. \square

We use these results to extend the definition of $\varepsilon_{\Lambda_R(G), \xi}(M)$ to lattices M a crystalline family of Galois representations with arbitrary (bounded) Hodge–Tate weights, by tensoring the corresponding maps for $M(j)$ with $R(-j)$, where $j \gg 0$ is such that $M(j)$ has non-negative Hodge–Tate weights. Clearly, $\varepsilon_{\Lambda_R(G), \xi}(M)$ is again compatible with specializations in R .

7.5. Epsilon-isomorphisms for more general modules. As before we let L denote a finite extension of \mathbf{Q}_p . We recall the following definition from [FK06, §1.4].

Definition 7.18. A ring A is of

- (**type 1**): if there exists a two sided ideal I of A such that A/I^n is finite of order a power of p for any $n \geq 1$, and such that $A \cong \varprojlim_n A/I^n$.
(**type 2**): if A is the matrix-algebra $M_n(E)$ of some finite extension E over \mathbf{Q}_p and some dimension $n \geq 1$.

By Lemma 1.4.4 in (loc. cit.), A is of type 1 if and only if the defining condition above holds with I equal to the Jacobson ideal $J = J(A)$. Such rings are always semi-local and A/J is a finite product of matrix algebras over finite fields. For a ring A of type (1) or (2) we define

$$\tilde{A} := \widehat{\mathbf{Z}_p^{\text{nr}}} \widehat{\otimes}_{\mathbf{Z}_p} A$$

where $\widehat{\mathbf{Z}_p^{\text{nr}}}$ denotes the completion of the ring of integers of the maximal unramified extension of \mathbf{Q}_p .

Now let M be a crystalline family of representation of $G_{\mathbf{Q}_p}$ with coefficients in R . We set $\mathbf{T}(M) := \Lambda_R(G) \otimes_R M$, which we consider as $\Lambda(G)$ -module by multiplication on the left tensor factor and as $G_{\mathbf{Q}_p}$ -module via $g(\lambda \otimes t) = \lambda \bar{g}^{-1} \otimes gt$. The following isomorphism (essentially a version of Shapiro's lemma) is well known:

Proposition 7.19. *We have*

$$R\Gamma(\mathbf{Q}_p, \mathbf{T}(M)) \cong R\Gamma_{\text{Iw}}(K_\infty, M)$$

as $\Lambda(G)$ -modules.

Proof. See e.g. [Nek06, 8.4.4.2 Proposition]. □

Let $\Lambda = \Lambda_R(G)$, which is a ring of type 1, with $\tilde{\Lambda} = \Lambda_{\tilde{R}}(G)$. Then we have constructed an isomorphism

$$\varepsilon_{\Lambda, \xi}(M) : \text{Det}_{\tilde{\Lambda}}(0) \xrightarrow{\cong} \tilde{\Lambda} \otimes_{\Lambda} \{\text{Det}_{\Lambda} R\Gamma(\mathbf{Q}_p, \mathbf{T}(M)) \cdot \text{Det}_{\Lambda} \mathbf{T}(M)\}.$$

We shall establish that this satisfies the properties predicted by [FK06, Conjecture 3.4.3] for the module $\mathbf{T}(M)$. For that purpose it is convenient to write also $\varepsilon_{\Lambda, \xi}(\mathbf{T}(M))$ for the above ε -isomorphism, and to extend it to a slightly more general class of modules.

We consider quadruples (A, Y, M, ξ) where

- A is a p -torsion-free R -algebra which is also a ring of type (1) or (2) above,
- ξ is a compatible system of p^n -th roots of unity (as before),
- M is a crystalline family of representation of $G_{\mathbf{Q}_p}$ with coefficients in R ,
- Y is a finitely-generated projective left A -module, equipped with a continuous A -linear action of G .

Given such a quadruple, we define $\mathbf{T} = Y \otimes_R M$, which we equip with the obvious left A -module structure and an action of $G_{\mathbf{Q}_p}$ via $g \cdot (y \otimes t) = y \bar{g}^{-1} \otimes gt$. Then (A, \mathbf{T}, ξ) is a triple satisfying the conditions of [FK06, §3.4.1]. Moreover, the action of G on Y extends to a Λ -module structure, and we have

$$\mathbf{T} = Y \otimes_{\Lambda} \mathbf{T}(M)$$

where $\mathbf{T}(M)$ is as above. So we may define

$$\varepsilon_{A, \xi}(\mathbf{T}) := Y \otimes_{\Lambda} \varepsilon_{\Lambda, \xi}(\mathbf{T}(M)),$$

which is an isomorphism

$$\text{Det}_{\tilde{A}}(0) \xrightarrow{\cong} \tilde{A} \otimes_A \{\text{Det}_A R\Gamma(\mathbf{Q}_p, \mathbf{T}) \cdot \text{Det}_A \mathbf{T}\};$$

here we have used the fact that

$$Y \otimes_{\Lambda}^L R\Gamma(\mathbf{Q}_p, \mathbf{T}(M)) \cong R\Gamma(\mathbf{Q}_p, Y \otimes_{\Lambda} \mathbf{T}(M))$$

by [FK06, 1.6.5].

Remark 7.20. Note that A need not be commutative, and Y need not be either projective or finitely-generated as a Λ -module.

prop:compat2

Proposition 7.21. *Suppose $A = \mathcal{O}_E$ for some finite extension E/\mathbf{Q}_p , and the finite-dimensional E -vector space $E \otimes_A Y$ is de Rham as a representation of G . Then $E \otimes_{\mathcal{O}_E} \mathbf{T}$ is also de Rham, and $E \otimes_{\mathcal{O}_E} \varepsilon_{A,\xi}(\mathbf{T})$ coincides with the canonical isomorphism $\varepsilon_{E,\xi}(E \otimes_{\mathcal{O}_E} \mathbf{T})$ of [LVZ15, §2.4], the ‘standard trivialisation’.*

Proof. We may assume that E is sufficiently large that all the Jordan–Hölder constituents of $E \otimes_A Y$ are one-dimensional. By the compatibility with short exact sequences, it suffices to assume $E \otimes_A Y$ is itself one-dimensional, so $E \otimes_A Y = E(\eta)$ for a de Rham character η of G . Moreover the R -algebra structure of A provides a homomorphism $R \rightarrow \mathcal{O}_E$, whence $\Lambda_R \rightarrow \Lambda_{\mathcal{O}_E}$, such that

$$\begin{aligned} E \otimes_{\mathcal{O}_E} \varepsilon_{A,\xi}(\mathbf{T}) &= E \otimes_{\mathcal{O}_E} Y \otimes_{\Lambda_R} \varepsilon_{\Lambda_R,\xi}(\mathbf{T}(M)) \\ &\cong E \otimes_{\mathcal{O}_E} Y \otimes_{\Lambda_{\mathcal{O}_E}} \Lambda_{\mathcal{O}_E} \otimes_{\Lambda_R} \varepsilon_{\Lambda_R,\xi}(\mathbf{T}(M)) \\ &\cong E \otimes_{\mathcal{O}_E} Y \otimes_{\Lambda_{\mathcal{O}_E}} \varepsilon_{\Lambda_{\mathcal{O}_E},\xi}(\Lambda_{\mathcal{O}_E} \otimes_{\Lambda_R} \mathbf{T}(M)) \\ &\cong E \otimes_{\mathcal{O}_E} Y \otimes_{\Lambda_{\mathcal{O}_E}} \varepsilon_{\Lambda_{\mathcal{O}_E},\xi}(\mathbf{T}(\mathcal{O}_E \otimes_R M)) \\ &\cong E \otimes_{\mathcal{O}_E} \varepsilon_{A,\xi}(\mathbf{T}) \\ &\cong \varepsilon_{E,\xi}(E \otimes_{\mathcal{O}_E} \mathbf{T}). \end{aligned}$$

Here the first identity is the definition of $\varepsilon_{A,\xi}(\mathbf{T})$, the second comes from the fact that $\Lambda_R \rightarrow \mathcal{O}_E$ factors through $\Lambda_R \rightarrow \Lambda_{\mathcal{O}_E}$, the third is the compatibility from Theorem 7.13 (and its extension to arbitrary crystalline families after Proposition 7.17) of $\varepsilon_{\Lambda_R,\xi}(\mathbf{T}(M))$ with respect to specialization in R , the fourth follows from the functoriality of the definition of $\mathbf{T}(M)$, while the fifth uses the definition of $\varepsilon_{A,\xi}(\mathbf{T})$ and the fact that

$$\mathbf{T} = Y \otimes_{\Lambda_R} \mathbf{T}(M) \cong Y \otimes_{\Lambda_{\mathcal{O}_E}} \mathbf{T}(\mathcal{O}_E \otimes_R M).$$

Finally, the last identification is just [LVZ15, Prop. 4.6.4]. \square

specializations

Corollary 7.22. *Suppose that the pair (A, \mathbf{T}) satisfies the following condition:*

- if $\Phi_{\mathbf{T}}$ is the set of all R -algebra homomorphisms $\rho : A \rightarrow M_n(E)$ (where E/L is a finite extension and n an integer, both depending on ρ) such that $E^n \otimes_{A,\rho} \mathbf{T}$ is de Rham, then

$$K_1(A) \rightarrow \prod_{\rho \in \Phi_{\mathbf{T}}} E^\times$$

is injective.

Then $\varepsilon_{A,\xi}(\mathbf{T})$ depends only on ξ and on the isomorphism class of \mathbf{T} as an $A[G_{\mathbf{Q}_p}]$ -module.

Proof. Clear from the preceding proposition, since the isomorphism $\varepsilon_{A,\xi}(\mathbf{T})$ must be consistent with the de Rham ε -isomorphisms $\varepsilon_{E,\xi}(E^n \otimes_{A,\rho} \mathbf{T})$, which are uniquely determined by (A, \mathbf{T}, ξ) . \square

Remark 7.23. We suspect that the uniqueness statement of the corollary is true for arbitrary type 1 R -algebras A , but this is much more difficult to prove in general. For instance, if M_1, M_2 are two crystalline families of representations over R such that $M_1/p^n \cong M_2/p^n$ for some $n \geq 1$, then on taking $A = R/p^n$ this would imply that the ε -isomorphisms for M_1 and M_2 are congruent modulo p^n . This should certainly be true, and actually one motivation for this paper is to show it at least for $R = \mathcal{O}_E$ with E any finite extension of \mathbf{Q}_p , see section 8 below.

We shall now show that the association $(A, Y, M, \xi) \rightarrow \varepsilon_{A,\xi}(\mathbf{T})$ satisfies properties corresponding to conditions (i)–(iv) and (vi) of [FK06, Conjecture 3.4.3].

Property (i) (additivity). The first condition of *op.cit.* states that for any three triples (A, \mathbf{T}_i, ξ) , $i = 1, 2, 3$, with common A and ξ , and an exact sequence

$$0 \longrightarrow \mathbf{T}_1 \longrightarrow \mathbf{T}_2 \longrightarrow \mathbf{T}_3 \longrightarrow 0,$$

we have

$$\varepsilon_{A,\xi}(\mathbf{T}_2) = \varepsilon_{A,\xi}(\mathbf{T}_1) \varepsilon_{A,\xi}(\mathbf{T}_3).$$

By assumption our \mathbf{T}_i are of the form $Y_i \otimes M_i$, for crystalline R -representations M_i and A -modules Y_i with G -action. We shall consider only the cases when the exact sequence arises from an exact sequence of Y_i ’s with a common M , or an exact sequence of M_i ’s with a common Y . The first case is obvious from the construction of $\varepsilon_{A,\xi}(-)$. The latter case follows from the additivity of the standard trivialisation (see [LVZ15, Prop. 2.4.3]) or from the additivity statement (property (i) in §4.6 in (*loc. cit.*)).

Property (ii) (base change). The second condition is a compatibility with base-change in A ; this is immediate from our construction.

Property (iii) (change of ξ). Let $c \in \mathbf{Z}_p^\times$, and let γ_c be any element of $G_{\mathbf{Q}_p}$ acting trivially on \mathbf{Q}_p^{nr} and such that $\chi(\gamma_c) = c$. Then we must show that

$$\varepsilon_{A,c\xi}(\mathbf{T}) = [M, \gamma_c] \varepsilon_{A,\xi}(\mathbf{T}),$$

where $[M, \gamma_c]$ is the class in $K_1(A)$ of the A -linear automorphism of \mathbf{T} given by γ_c . (This is well-defined, as γ_c is uniquely determined up to conjugation in $G_{\mathbf{Q}_p}$.) It suffices to check this when $A = \Lambda$ and $\mathbf{T} = \mathbf{T}(M)$; but this is immediate from the corresponding property of the specializations (property (iii) in §4.6 in (loc. cit.)).

Property (iv) (Galois equivariance). Let φ denote the arithmetic Frobenius automorphism of $\widehat{\mathbf{Z}}_p^{\text{nr}}$. Then we must show that

$$\varepsilon_{A,\xi}(\mathbf{T}) \in \text{Isom}(\text{Det}_A(0), \text{Det}_A R\Gamma(\mathbf{Q}_p, \mathbf{T}) \cdot \text{Det}_A \mathbf{T}) \times^{K_1(A)} \left\{ x \in K_1(\tilde{A}) : \varphi(x) = [\mathbf{T}, \sigma_p]^{-1} x \right\}$$

where σ_p is the arithmetic Frobenius element of $\text{Gal}(\mathbf{Q}_p^{\text{ab}}/\mathbf{Q}_p)$. Again, it suffices to assume $(A, \mathbf{T}) = (\Lambda, \mathbf{T}(M))$ and the result is now clear from the Galois-equivariance properties of the specializations (property (iv) in §4.6 in (loc. cit.)).

Property (v) (compatibility with de Rham ε -isomorphisms). If A is the ring of integers of a finite extension F/L , and $F \otimes_A \mathbf{T}$ is de Rham, we must check that $\varepsilon_{A,\xi}(\mathbf{T})$ is consistent with the standard trivialisation. This is exactly Proposition 7.21 above.

Property (vi) (local duality). Let \mathbf{T} be a free A -module with compatible $G_{\mathbf{Q}_p}$ -action as above. Then

$$\mathbf{T}^* := \text{Hom}_A(\mathbf{T}, A)$$

is a free A° -module – for the action $h \mapsto h(-)r$, r in the opposite ring A° of A – with compatible $G_{\mathbf{Q}_p}$ -action given by $h \mapsto h \circ \sigma^{-1}$. Recall that in Iwasawa theory we have the canonical involution $\iota : \Lambda^\circ \rightarrow \Lambda$, induced by $g \mapsto g^{-1}$, which allows to consider (left) Λ° -modules again as (left) Λ -modules, e.g. one has $\mathbf{T}^*(M)^\iota \cong \mathbf{T}(M^*)$ as $(\Lambda, G_{\mathbf{Q}_p})$ -module, where $M^\iota := \Lambda \otimes_{\iota, \Lambda^\circ} M$ denotes the Λ -module with underlying abelian group M , but on which $g \in G$ acts as g^{-1} for any Λ° -module M .

Given $\varepsilon_{A^\circ, -\xi}(\mathbf{T}^*(1))$ we may apply the dualising functor $-^*$ to obtain an isomorphism

$$\varepsilon_{A^\circ, -\xi}(\mathbf{T}^*(1))^* : (\text{Det}_{A^\circ}(R\Gamma(\mathbf{Q}_p, \mathbf{T}^*(1)))_{\tilde{A}^\circ})^* (\text{Det}_{A^\circ}(\mathbf{T}^*(1))_{\tilde{A}^\circ})^* \rightarrow \mathbf{1}_{\tilde{A}^\circ},$$

while the local Tate duality isomorphism [FK06, §1.6.12]

$$\psi(\mathbf{T}) : R\Gamma(\mathbf{Q}_p, \mathbf{T}) \cong R\text{Hom}_{A^\circ}(R\Gamma(\mathbf{Q}_p, \mathbf{T}^*(1)), A^\circ)[-2]$$

induces an isomorphism

$$\overline{\text{Det}_A(\psi(\mathbf{T}))}_{\tilde{A}}^{-1} : ((\text{Det}_{A^\circ}(R\Gamma(\mathbf{Q}_p, \mathbf{T}^*(1)))_{\tilde{A}^\circ})^*)^{-1} \cong \text{Det}_A(R\text{Hom}_{A^\circ}(R\Gamma(\mathbf{Q}_p, \mathbf{T}^*(1)), A^\circ))_{\tilde{A}}^{-1} \rightarrow \text{Det}_A(R\Gamma(\mathbf{Q}_p, \mathbf{T}))_{\tilde{A}}^{-1}.$$

Here, for a map $f : A \rightarrow B$ in $\underline{\text{Det}}(A)$, we write $\bar{f} : B \rightarrow A$ for its inverse with respect to composition, while $f^{-1} =: \overline{\text{id}_{B^{-1}} \cdot f \cdot \text{id}_{A^{-1}}} : A^{-1} \rightarrow B^{-1}$ for its inverse with respect to the multiplication in $\underline{\text{Det}}(A)$, i.e., $f \cdot f^{-1} = \text{id}_{\text{Det}_A(0)}$.

Consider the product

$$\varepsilon_{A,\xi}(\mathbf{T}) \cdot \varepsilon_{A^\circ, -\xi}(\mathbf{T}^*(1))^* \cdot \overline{\text{Det}_A(\psi(\mathbf{T}))}_{\tilde{A}}^{-1} : \text{Det}_A(\mathbf{T}(-1))_{\tilde{A}} \cong \text{Det}_A(\mathbf{T}^*(1)^*)_{\tilde{A}} \rightarrow \text{Det}_A(\mathbf{T})_{\tilde{A}}$$

and the isomorphism $\mathbf{T}(-1) \xrightarrow{\cdot \xi} \mathbf{T}$ which sends $t \otimes \xi^{\otimes -1}$ to t .

duality

Proposition 7.24 (Duality). *Let \mathbf{T} be as above such that $\mathbf{T} \cong Y \otimes_\Lambda \mathbf{T}(M)$ for some (A, Λ) -bimodule Y , which is projective as A -module. Then*

$$\varepsilon_{A,\xi}(\mathbf{T}) \cdot \varepsilon_{A^\circ, -\xi}(\mathbf{T}^*(1))^* \cdot \overline{\text{Det}_A(\psi(\mathbf{T}))}_{\tilde{A}}^{-1} = \text{Det}_A \left(\mathbf{T}(-1) \xrightarrow{\cdot \xi} \mathbf{T} \right)_{\tilde{A}}.$$

Proof. First note that the statement is stable under applying $Y' \otimes_A -$, for some (A', A) -bimodule Y' which is projective as a A' -module, by the functoriality of local Tate duality and the lemma below. Thus we are reduced to the case $(A, \mathbf{T}) = (\Lambda, \mathbf{T}(M))$ where M is a crystalline family of representation of $G_{\mathbf{Q}_p}$ with coefficients in R . Thus the claim follows again by specialisation and (property (vi) in §4.6 in (loc. cit.)) using Lemma 7.14. \square

Lemma 7.25. *Let Y be a (A', A) -bimodule such that $Y \otimes_A \mathbf{T} \cong \mathbf{T}'$ as $(A', G_{\mathbf{Q}_p})$ -module and let $Y^* = \text{Hom}_{A'}(Y, A')$ the induced (A°, A°) -bimodule. Then there is a natural*

(1) *equivalence of functors*

$$Y \otimes_A \text{Hom}_{A^\circ}(-, A^\circ) \cong \text{Hom}_{A^\circ}(Y^* \otimes_{A^\circ} -, A^\circ)$$

on $P(A^\circ)$;

(2) *isomorphism $Y^* \otimes_{A^\circ} \mathbf{T}^* \cong (\mathbf{T}')^*$ of $(A^\circ, G_{\mathbf{Q}_p})$ -modules.*

Proof. This is easily checked using the adjointness of Hom and \otimes . \square

8. APPLICATION TO DEFORMATION RINGS

Consider the following situation: For $i = 1, 2$ let T_i be a crystalline G_F -representation over \mathcal{O} with Hodge-Tate weights in $[a, b]$ (with integers a, b), such that there exist some $n_0 \geq 1$ such that

$$T_1/\varpi^n T_1 \cong T_2/\varpi^n T_2 =: T_{(n)}$$

for $n_0 \geq n \geq 1$. Now let $R_{T_{(1)}}^{\square, \text{cris}, [a, b]}$ be universal framed deformation ring together with the universal G_F -representation $T_{T_{(1)}}^{\square, \text{cris}, [a, b]}$ parameterising crystalline lifts T of $T_{(1)}$ over \mathcal{O} -algebras R with Hodge-Tate weights in $[a, b]$.

Proposition 8.1. *If $R_{T_{(1)}}^{\square, \text{cris}, [a, b]}$ is a \mathbf{Z}_p -flat, normal, Cohen-Macaulay ring such that $R_{T_{(1)}}^{\square, \text{cris}, [a, b]}[\frac{1}{p}]$ is reduced, then, for all $n_0 \geq n \geq 0$, we have*

$$\mathcal{O}/\varpi^n \mathcal{O} \otimes_{\mathcal{O}} \varepsilon_{\mathcal{O}, \xi}(T_1) = \mathcal{O}/\varpi^n \mathcal{O} \otimes_{\mathcal{O}} \varepsilon_{\mathcal{O}, \xi}(T_2).$$

In particular, this allows us to define $\varepsilon_{p, \mathcal{O}/\varpi^n \mathcal{O}}(T_{(n)})$.

Proof. Since T_i and $T_{(n)}$ are lifts of $T_{(1)}$, for $i = 1, 2$, they corresponds to \mathcal{O} -algebra homomorphisms $\pi_i : R_{T_{(1)}}^{\square, \text{cris}, [a, b]} \rightarrow \mathcal{O}$ and $\pi_{(n)} : R_{T_{(1)}}^{\square, \text{cris}, [a, b]} \rightarrow \mathcal{O}/\varpi^n \mathcal{O}$, respectively, giving rise to a commutative diagram for each i

$$\begin{array}{ccc} R_{T_{(1)}}^{\square, \text{cris}, [a, b]} & \xrightarrow{\pi_i} & \mathcal{O} \\ & \searrow \pi_{(n)} & \downarrow pr \\ & & \mathcal{O}/\varpi^n \mathcal{O}. \end{array}$$

Thus we obtain for $i = 1, 2$

$$\begin{aligned} \mathcal{O}/\varpi^n \mathcal{O} \otimes_{\mathcal{O}} \varepsilon_{\mathcal{O}, \xi}(T_i) &= \mathcal{O}/\varpi^n \mathcal{O} \otimes_{\mathcal{O}} \left(\mathcal{O} \otimes_{R_{T_{(1)}}^{\square, \text{cris}, [a, b]}, \pi_i} \varepsilon_{R_{T_{(1)}}^{\square, \text{cris}, [a, b]}, \xi}(T_{T_{(1)}}^{\square, \text{cris}, [a, b]}) \right) \\ &= \mathcal{O}/\varpi^n \mathcal{O} \otimes_{R_{T_{(1)}, \pi_{(n)}}^{\square, \text{cris}, [a, b]}} \varepsilon_{R_{T_{(1)}}^{\square, \text{cris}, [a, b]}, \xi}(T_{T_{(1)}}^{\square, \text{cris}, [a, b]}), \end{aligned}$$

whence the claim. \square

Remark 8.2. (1) It follows from [CHT08, §2.4] that $R_{T_{(1)}}^{\square, \text{cris}, [a, b]}$ satisfies the conditions of the proposition, if $b - a < p - 1$. Indeed, in that case it is isomorphic to a power series ring over \mathcal{O} .

(2) A similar statement holds also for $\varepsilon_{\Lambda \mathcal{O}, \xi}(\mathbf{T}(T_i))$, of course.

9. RELATION TO NAKAMURA'S WORK

Let M be a crystalline family of representations of $G_{\mathbf{Q}_p}$ with coefficients in R and let $\mathbf{T}(M) := \Lambda_R(\Gamma) \otimes_R M$, with $G_{\mathbf{Q}_p}$ action given by $g(x \otimes y) := g(x) \otimes [\bar{g}]^{-1}y$. Then the generic fiber M^{rig} is a family of Galois representations over $Y := \text{Spf}(R)^{\text{rig}}$, and by [KL10, Thm. 3.11] there is a (φ, Γ) -module $\mathbf{D}_{\text{rig}}(M^{\text{rig}})$ over the relative Robba ring \mathcal{R}_Y attached to M^{rig} . We also define its universal cyclotomic deformation $\mathcal{M} := \mathbf{D}_{\text{rig}}(M^{\text{rig}}) \widehat{\otimes}_{R^{\text{rig}}} \Lambda_R(\Gamma)^{\text{rig}}$, where φ is taken to act trivially on $\Lambda_R(\Gamma)^{\text{rig}}$ and Γ acts via $\gamma(d \otimes x) = \gamma(d) \otimes [\gamma]^{-1}x$. Since M^{rig} is a crystalline family, it is trianguline after making a finite extension of R , so \mathcal{M} is trianguline as well; let \mathcal{F} denote a triangulation of \mathcal{M} . In this situation, Nakamura [Nak13, Cor. 3.12] has constructed an ε -isomorphism

$$(13) \quad \varepsilon_{\mathcal{F}, \mathcal{H}_R(\Gamma), \xi}^N(\mathcal{M}) : \text{Det}_{\mathcal{H}_R(\Gamma)}(0) \xrightarrow{\sim} \Delta_{\Lambda_R(\Gamma)^{\text{rig}}}(\mathcal{M})$$

where $\Delta_{\Lambda_R(\Gamma)^{\text{rig}}}(\mathcal{M})$ is a (graded) invertible $\text{Spf}(\Lambda_R(\Gamma)^{\text{rig}})$ -module such that

$$(14) \quad \mathcal{H}_{\tilde{R}}(\Gamma) \otimes_{\mathcal{H}_R(\Gamma)} (\Delta_{\Lambda_R(\Gamma)^{\text{rig}}}(\mathcal{M})) \cong \mathcal{H}_{\tilde{R}}(\Gamma) \otimes_{\Lambda_R(\Gamma)} \{ \text{Det}_{\Lambda_R(\Gamma)} R\Gamma(\mathbf{Q}_p, \mathbf{T}(M)) \cdot \text{Det}_{\Lambda_R(\Gamma)} \mathbf{T}(M) \}$$

by Cor. 3.2 in (loc. cit.).

Corollary 9.1. *Under the isomorphism (14) we have*

$$\mathcal{H}_{\tilde{R}}(\Gamma) \otimes_{\Lambda_{\tilde{R}}(G)} \varepsilon_{\Lambda_{\tilde{R}}(G), \xi}(M) = \mathcal{H}_{\tilde{R}}(\Gamma) \otimes_{\mathcal{H}_R(\Gamma)} \varepsilon_{\mathcal{F}, \mathcal{H}_R(\Gamma), \xi}^N(\mathcal{M}).$$

In particular, the isomorphism $\varepsilon_{\mathcal{F}, \mathcal{H}_R(\Gamma), \xi}^N(\mathcal{M})$ does not depend on the choice of \mathcal{F} .

Proof. Recall that $\text{Spf}(\Lambda_{\tilde{R}}(\Gamma))^{\text{rig}} \xrightarrow{\sim} \text{Spf}(\tilde{R})^{\text{rig}} \times \text{Spf}(\Lambda_{\tilde{Z}_p}(\Gamma))^{\text{rig}}$, so that a point of $\text{Spf}(\Lambda_{\tilde{R}}(\Gamma))^{\text{rig}}$ is a pair (x, η) , where $x \in \text{Spf}(\tilde{R})^{\text{rig}}$ and η corresponds to a character $G \rightarrow \overline{\mathbf{Q}_p^{\text{nr}}}^\times$ (which we also write η). If (x, η) lies over a point $(x', \eta') \in \text{Spf}(\Lambda_R(\Gamma))^{\text{rig}}$ and η' is a de Rham character, we can compute the specializations of both sides at (x, η) . Indeed, both sides specialize to $\widehat{\mathbf{Q}_p^{\text{nr}}} \otimes_{\mathbf{Q}_p} \varepsilon_{\kappa(x'), \xi}(M \otimes_R \kappa(x')^\circ)[1/p](\eta'^{-1})$, that is, the base-change to $\widehat{\kappa(x')}$ of Fukaya–Kato's ε -isomorphism for de Rham representations.

Points of $\text{Spf}(\Lambda_{\tilde{Z}_p}(\Gamma))^{\text{rig}}$ corresponding to de Rham characters are Zariski dense in the rigid analytic space. Combined with Lemma 7.1, we conclude that the two maps agree. \square

APPENDIX A. GENERIC FIBERS

A.1. Quasi-Stein spaces.

Definition A.1. A rigid analytic space Y over K is said to be *quasi-Stein* if it admits an admissible covering by a rising union of affinoid subdomains $Y_0 \subset Y_1 \subset \dots$ such that the transition maps $\Gamma(Y_{m+1}, \mathcal{O}_{Y_{m+1}}) \rightarrow \Gamma(Y_m, \mathcal{O}_{Y_m})$ are flat with dense image.

Example A.2. Fix $s > 0$, and let X be the coordinate on the closed unit disk. Then the half-open annulus $0 < v_p(X) \leq 1/s$ is a quasi-Stein space, as it is the rising union of the closed annuli $1/s' \leq v_p(X) \leq 1/s$ as $s' \rightarrow \infty$.

By Kiehl's results on coherent sheaves on rigid analytic spaces, a coherent sheaf \mathcal{F} on Y is simply a compatible system of coherent sheaves $\{\mathcal{F}_m\}$ on $\{Y_m\}$.

Quasi-Stein spaces behave much as affine schemes do in algebraic geometry. In particular, Kiehl proved the following theorem on the cohomology of coherent sheaves on quasi-Stein spaces.

Theorem A.3 ([Kie67, Satz 2.4]). *Let Y be a quasi-Stein space, and let \mathcal{F} be a coherent sheaf on Y . Then*

- (1) $H^i(Y, \mathcal{F}) = 0$ for $i > 0$,
- (2) the image of $\mathcal{F}(Y)$ in $\mathcal{F}(Y_m)$ is dense for all m ,
- (3) for every point $y \in Y$, the image of $\mathcal{F}(Y)$ in \mathcal{F}_y generates \mathcal{F}_y .

In particular, $A_\infty := \Gamma(Y, \mathcal{O}_Y) = \varprojlim_m \Gamma(Y_m, \mathcal{O}_{Y_m})$ is a Fréchet-Stein algebra and $F_\infty := \Gamma(Y, \mathcal{F}) = \varprojlim_m \Gamma(Y_m, \mathcal{F}_m)$ is a coadmissible module over A_∞ , in the sense of [ST03]. By [ST03, §3], the natural morphisms $F_\infty \rightarrow \Gamma(Y_m, \mathcal{F}_m)$ have dense image, and $\mathbf{R}^i \varprojlim_m \Gamma(Y_m, \mathcal{F}_m) = 0$ for $i > 0$.

There is no *a priori* reason for F_∞ to be a finite module over A_∞ . For example, let $X_m = \text{Sp}(\prod_{i=0}^m \mathbf{Q}_p(\zeta_{p^i}))$, where $\text{Sp}(A)$ denotes the affinoid space associated to A , and let \mathcal{F}_m be the sheaf on Y_m associated to

$\prod_{i=0}^n \mathbf{Q}_p(\zeta_{p^i})^{\oplus i}$. Then F_∞ is not A_∞ -finite, because the fiber of F_∞ at $\mathrm{Sp}(\mathbf{Q}_p(\zeta_{p^n}))$ is a $\mathbf{Q}_p(\zeta_{p^n})$ -vector space of dimension n . Happily, this is the only thing that can go wrong.

fibrals-ranks

Lemma A.4 ([Bel15, Lemma 2.4.4]). *Let \mathcal{F} be a coherent sheaf over a finite-dimensional quasi-Stein space Y over E . Then $H^0(Y, \mathcal{F})$ is finitely generated as an $H^0(Y, \mathcal{O}_Y)$ -module if and only if there is some integer d such that $\dim_{\kappa(y)} \mathcal{F}(y) \leq d$ for all $y \in Y$.*

fibrals-ranks

Corollary A.5 ([Bel15, Corollary 2.4.5]). *Suppose that \mathcal{F} is flat over \mathcal{O}_Y , where Y is a finite-dimensional quasi-Stein space. Then $H^0(Y, \mathcal{F})$ is projective of rank d over $H^0(Y, \mathcal{O}_Y)$ if and only if $\dim_{\kappa(y)} \mathcal{F}(y) = d$ for all $y \in Y$.*

An important family of examples of quasi-Stein spaces are those arising as the generic fibers of affine formal schemes of the form $\mathrm{Spf}(R)$ with $R = \mathcal{O}(Y_1, \dots, Y_{d_1})[[X_1, \dots, X_{d_2}]]/I$. Recall that this is a construction due to Berthelot (and expositied in [dJ95, §7]). Briefly, we construct the generic fiber $\mathrm{Spf}(R)^{\mathrm{rig}}$ of $\mathrm{Spf}(R)$ by giving a cover, as follows: If J is an ideal of definition of R , then for each $m \geq 0$, we consider the ϖ -adic completion R_m of $R[J^m/\varpi]$ inside $R[1/p]$. It turns out that $R_m[1/p]$ is K -affinoid and $\{\mathrm{Sp}(R_m[1/p])\}$ is a rising union, with $\mathrm{Sp}(R_m[1/p])$ identified with an affinoid subdomain of $\mathrm{Sp}(R_{m+1}[1/p])$. The global sections of the structure sheaf on $\mathrm{Spf}(R)^{\mathrm{rig}}$ will be the ring $R^{\mathrm{rig}} := \varprojlim_m R_m[1/p]$. In general, R^{rig} will be non-Noetherian (unless $d_2 = 0$).

If R is a formal power series ring, this construction corresponds to exhausting the open unit polydisk by closed balls of radius $|\varpi|^{1/m}$.

There is a canonical homomorphism $R \rightarrow R^{\mathrm{rig}}$, and it follows from [dJ95, Lemma 7.1.9] that this map is flat. More precisely, maximal ideals of $R[1/p]$ are in bijection with points of $\mathrm{Spf}(R)^{\mathrm{rig}}$, and the canonical homomorphism induces isomorphisms on complete local rings. Thus, $R[1/p] \rightarrow R^{\mathrm{rig}}$ is faithfully flat.

Given a finite R -module M , we can construct a coherent sheaf M^{rig} on $\mathrm{Spf}(R)^{\mathrm{rig}}$. Explicitly, $M_{\mathrm{Sp}(R_m[1/p])}^{\mathrm{rig}} := M \otimes_R R_m[1/p]$ (in fact, since the homomorphisms $R \rightarrow R_m[1/p]$ factor through $R[1/p]$, we can construct a coherent sheaf on $\mathrm{Spf}(R)^{\mathrm{rig}}$ from any finite $R[1/p]$ -module). This is sufficient to define M^{rig} on any admissible open subset of $\mathrm{Spf}(R)^{\mathrm{rig}}$. Moreover, the global sections of M^{rig} behave exactly as we would hope:

lemma:rig

Lemma A.6. $\Gamma(\mathrm{Spf}(R)^{\mathrm{rig}}, M^{\mathrm{rig}}) = M \otimes_R R^{\mathrm{rig}}$.

Proof. We have a natural homomorphism $M \otimes_R R^{\mathrm{rig}} \rightarrow \varprojlim_m (M \otimes_R R_m[1/p])$, and we need to show it is an isomorphism. This is clear if M is finite free. Next, we check that if $R^{\oplus n_1} \rightarrow M$ finitely presented, $(R^{\mathrm{rig}})^{\oplus n_1} \rightarrow \Gamma(\mathrm{Spf}(R)^{\mathrm{rig}}, M^{\mathrm{rig}})$.

So suppose we have an exact sequence

$$0 \rightarrow K_1 \rightarrow R^{\oplus n_1} \rightarrow M \rightarrow 0$$

for some R -finite module K . Since $R \rightarrow R_m[1/p]$ is flat, for each m we have an exact sequence

$$0 \rightarrow K_1 \otimes_R R_m[1/p] \rightarrow (R_m[1/p])^{\oplus n_1} \rightarrow M \otimes_R R_m[1/p] \rightarrow 0$$

But then $\mathbf{R}^1 \varprojlim_m (K_1 \otimes_R R_m[1/p]) = 0$, so $(R^{\mathrm{rig}})^{\oplus n_1} \rightarrow \Gamma(\mathrm{Spf}(R)^{\mathrm{rig}}, M^{\mathrm{rig}})$.

Now we choose a presentation of K_1 , yielding an exact sequence

$$0 \rightarrow K_2 \rightarrow R^{\oplus n_2} \rightarrow K_1 \rightarrow 0$$

The same argument as above, applied to K_1 , shows that

$$0 \rightarrow \varprojlim_m K_2 \otimes_R R_m[1/p] \rightarrow (R^{\mathrm{rig}})^{\oplus n_2} \rightarrow \varprojlim_m K_1 \otimes_R R_m[1/p] \rightarrow 0$$

remains exact.

In particular, we have an commutative diagram with exact rows

$$\begin{array}{ccccccc} (R^{\mathrm{rig}})^{\oplus n_2} & \longrightarrow & (R^{\mathrm{rig}})^{\oplus n_1} & \longrightarrow & M \otimes_R R^{\mathrm{rig}} & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ (R^{\mathrm{rig}})^{\oplus n_2} & \longrightarrow & (R^{\mathrm{rig}})^{\oplus n_1} & \longrightarrow & \Gamma(\mathrm{Spf}(R)^{\mathrm{rig}}, M^{\mathrm{rig}}) & \longrightarrow & 0 \end{array}$$

The two left vertical arrows are isomorphisms, so it follows that $M \otimes_R R^{\mathrm{rig}} \xrightarrow{\sim} \Gamma(\mathrm{Spf}(R)^{\mathrm{rig}}, M^{\mathrm{rig}})$. \square

Now we restrict to the case when $R = \mathcal{O}[[X_1, \dots, X_d]]$, so that R^{rig} is the ring of analytic functions on the open unit d -dimensional polydisk.

Definition A.7. For $0 < \rho < 1$, the ρ -Gauss norm on R^{rig} is defined by $\|\sum_{I \in \mathbf{Z}_{\geq 0}^{\oplus d}} a_I \underline{X}^I\|_{\rho} = \sup_I |a_I| \rho^{|I|}$.

It turns out that $\|f\|_{\rho} = \sup_{x \in B[0, \rho]} |f(x)|$ where the supremum is taken over points $x \in \overline{K}^{\oplus d}$ with $|x| \leq \rho$, and the supremum is actually attained at some point. Moreover, the ρ -Gauss norms are multiplicative, i.e., $\|f_1 f_2\|_{\rho} = \|f_1\|_{\rho} \|f_2\|_{\rho}$. This is a special property of polydisks; in general the sup norm on an affinoid is only submultiplicative.

bdd-fncns

Lemma A.8. Suppose $f \in R^{\text{rig}}$ and there is a constant C such that $|f(x)| \leq C$ for all $x \in (\overline{K})^{\oplus d}$ with $|x| < 1$. Then $f \in R[1/p]$.

Proof. If $|f(x)|$ is bounded by C on the open polydisk, then $\|f\|_{\rho} \leq C$ for all $\rho < 1$. Then

$$\|p^{\log_p C} f\|_{\rho} = \sup_I \{C^{-1} |a_I| \rho^{|I|}\} \leq 1$$

It follows that $|p^{\log_p C} a_I| \rho^{|I|} \leq 1$ for all I and all $\rho < 1$, and therefore $|p^{\log_p C} a_I| \leq 1$ for all I . Therefore, $p^{\log_p C} f \in R$. \square

Although R^{rig} is much bigger than $R[1/p]$, we can exploit the multiplicativity of the ρ -Gauss norms to prove that the units of the two rings are the same:

units

Proposition A.9. $(R^{\text{rig}})^{\times} = (R[1/p])^{\times}$

Proof. We need to show that if $f \in (R^{\text{rig}})^{\times}$, then f is bounded. So suppose f is invertible on $\text{Spf}(R)^{\text{rig}}$, and consider a sequence of closed balls with rational radii exhausting $\text{Spf}(R)^{\text{rig}}$, i.e., their radii form a sequence $\{\rho_i\}_{i \geq 1}$ of rational numbers which is strictly increasing and goes to 1. Let $c_i := \|f\|_{\rho_i}$, so that c_i is the maximum of $|f(x)|$ on the closed ball of radius ρ_i centered at the origin.

Since f is assumed invertible and $ff^{-1} = 1$, multiplicativity of the ρ -Gauss norm implies that $\|f^{-1}\|_{\rho_i} = 1/c_i$. But if $i < j$, then $\rho_i < \rho_j$ so

$$1/c_i = \|f^{-1}\|_{\rho_i} \leq \|f^{-1}\|_{\rho_j} = 1/c_j$$

This implies $c_j \leq c_i$. In particular, $c_j \leq c_1$ for all j and so $f \in R[1/p]$ by Lemma A.8. \square

In fact, we can prove more.

localize-units

Proposition A.10. Suppose $f, f' \in R^{\text{rig}}$ and $0 \neq ff' = g \in R[1/p]$. Then $f, f' \in R[1/p]$.

Proof. We may assume that $g \in R$, so that $|g(x)| \leq 1$ for every closed point $x \in \text{Spec } R[1/p]$; let $C := \sup_{x \in B[0, 1]} |g(x)|$, where the supremum is taken over points $x \in \overline{K}^{\oplus d}$ with $|x| < 1$. Let $0 < \rho < \rho' < 1$ be rational numbers.

Let $c := \|f\|_{\rho}$, so that c is the maximum of $|f(x)|$ on the closed ball of radius ρ centered at the origin, and let $c' := \|f\|_{\rho'}$. Then multiplicativity of the ρ -Gauss norm implies that $\|f'\|_{\rho} = \|g\|_{\rho}/c$ and $\|f'\|_{\rho'} = \|g\|_{\rho'}/c'$. But $\rho < \rho'$ so

$$\|g\|_{\rho}/c = \|f'\|_{\rho} \leq \|f'\|_{\rho'} = \|g\|_{\rho'}/c'$$

This implies that

$$c' \leq (\|g\|_{\rho'}/\|g\|_{\rho})c \leq (C/\|g\|_{\rho})c$$

This holds for every $\rho' > \rho$, so $f \in R[1/p]$ by Lemma A.8. \square

partial-gen-fib

A.2. Partial generic fibers. Given two complete local noetherian \mathcal{O} -algebras R_1 and R_2 , we wish to define a ring of functions on the rigid analytic space $\text{Spf}(R_1 \widehat{\otimes} R_2)^{\text{rig}} = \text{Spf}(R_1)^{\text{rig}} \times \text{Spf}(R_2)^{\text{rig}}$ consisting of those functions which are ‘‘bounded along $\text{Spf}(R_2)^{\text{rig}}$ ’’, but not necessarily ‘‘along $\text{Spf}(R_1)^{\text{rig}}$ ’’. By way of contrast, $(R_1 \widehat{\otimes} R_2)[1/p]$ roughly consists of bounded functions, whereas the ring of all analytic functions on $\text{Spf}(R_1 \widehat{\otimes} R_2)^{\text{rig}}$ imposes no boundedness conditions.

Definition A.11. If $I_1 \subset R_1$ is the biggest ideal of definition of R_1 , recall that we defined $R_{1,m}$ to be the ϖ -adic completion of $R_1[I_1^m/\varpi]$ inside $R_1[1/p]$. If $R_{1,m} \widehat{\otimes} R_2$ is the completed tensor product with respect to the profinite topology on R_2 , we define

$$R_1^{\text{rig}} \widehat{\otimes} R_2[1/p] := \varprojlim_m (R_{1,m} \widehat{\otimes} R_2[1/p])$$

We will actually only need this construction when $R_1 = \mathcal{O}[\underline{X}_1, \dots, \underline{X}_k]$, so we work out a more concrete description of $\mathcal{O}[\underline{X}]^{\text{rig}} \widehat{\otimes} R$.

Proposition A.12. *The ring $\mathcal{O}[\underline{X}]^{\text{rig}} \widehat{\otimes} R[1/p]$ defined above is equal to the ring*

$$\left\{ \sum_{\underline{n} \in \mathbf{Z}_{\geq 0}^{\oplus k}} r_{\underline{n}} \underline{X}^{\underline{n}} \mid r_{\underline{n}} \in R[1/p] \text{ and for all } \underline{\rho} \in \mathbf{Q}_{>0}^{\oplus k}, \varpi^{n_{\rho}} r_{\underline{n}} \in R \text{ for } n \gg 0 \right\}$$

as subrings of the ring of analytic functions on the generic fiber $\text{Spf}(R[\underline{X}])^{\text{rig}}$.

Proof. Both rings are subrings of the ring of functions on $\text{Spf}(R[\underline{X}])^{\text{rig}}$; every such function f has a unique representative of the form $f = \sum_{\underline{n} \in \mathbf{Z}_{\geq 0}^{\oplus k}} r_{\underline{n}} \underline{X}^{\underline{n}}$, where $r_{\underline{n}} \in R^{\text{rig}}$ and for every k -tuple of positive rational numbers $\underline{\rho}$, $\varpi^{n_{\rho}} r_{\underline{n}} \rightarrow 0$ in R^{rig} .

Suppose that $f = \sum_{\underline{n} \in \mathbf{Z}_{\geq 0}^{\oplus k}} r_{\underline{n}} \underline{X}^{\underline{n}}$, where $r_{\underline{n}} \in R[1/p]$ and for every k -tuple of positive rational numbers $\underline{\rho}$, $\varpi^{n_{\rho}} r_{\underline{n}} \in R$ for sufficiently large \underline{n} . Taking $\underline{\rho} = (1/2s, \dots, 1/2s)$ for $s \in \mathbf{Z}_{>0}$, our assumption implies that $f \in (\mathcal{O}(\underline{X}, \frac{\underline{X}^s}{\varpi}) \widehat{\otimes} R)[1/p]$. Since this holds for every such s , we see that $f \in \mathcal{O}[\underline{X}]^{\text{rig}} \widehat{\otimes} R[1/p]$.

On the other hand, if $f \in (\mathcal{O}(\underline{X}, \frac{\underline{X}^s}{\varpi}) \widehat{\otimes} R)[1/p]$, there is some $k \in \mathbf{Z}$ such that $p^k f \in \mathcal{O}(\underline{X}, \frac{\underline{X}^s}{\varpi}) \widehat{\otimes} R$. This implies that $\varpi^{k+n/s} r_{\underline{n}} \in R$ for all \underline{n} , which implies that for $n \geq ks$, $\varpi^{2n/s} r_{\underline{n}} \in R$. Thus, if $f \in \mathcal{O}[\underline{X}]^{\text{rig}} \widehat{\otimes} R[1/p]$, the coefficients satisfy the desired growth condition. \square

Remark A.13. This alternate characterization of $\mathcal{O}[\underline{X}]^{\text{rig}} \widehat{\otimes} R[1/p]$ makes it clear that if $\mathfrak{m} \subset R[1/p]$ is a maximal ideal, then $\mathcal{O}[\underline{X}]^{\text{rig}} \widehat{\otimes} (R[1/p]/\mathfrak{m}) \cong (\mathcal{O}[\underline{X}]^{\text{rig}} \widehat{\otimes} R[1/p])/\mathfrak{m}$.

Proposition A.14. *Suppose that R is normal and $R[1/p]$ is reduced. If $f, f' \in R[1/p] \widehat{\otimes} \mathcal{O}[\underline{X}]^{\text{rig}}$ satisfy $ff' = g \in R[1/p]$ and g is not identically zero on any irreducible component of $\text{Spec } R[1/p]$, then $f, f' \in (R \widehat{\otimes} \mathcal{O}[\underline{X}])[1/p]$.*

Proof. We may assume that $g \in R$, so that $|g(x)| \leq 1$ for every closed point $x \in \text{Spec } R[1/p]$. For a closed point $x \in \text{Spec } R[1/p]$ and a rational number $0 < \rho < 1$, let $\|f\|_{x, \rho}$ denote the supremum of $|f|$ on the subset $\{x\} \times B[0, \rho] \subset \text{MaxSpec } R[1/p] \times (\text{Spf } \mathcal{O}[\underline{X}])^{\text{rig}}$. Let $\|f\|_{\rho}$ denote the supremum of $|f|$ over all of $\text{MaxSpec } R[1/p] \times B[0, \rho]$.

If $x \in \text{Spec } R[1/p]$ is a closed point such that $g(x) \neq 0$, then the proof of Proposition A.9 shows that for every pair of rational numbers $0 < \rho < \rho' < 1$,

$$\|f\|_{x, \rho'} \leq \|f\|_{x, \rho} \leq \|f\|_{\rho}$$

Thus, f is bounded on the complement of the zero locus of g ; suppose $|f(x)| \leq C$ for all $x \in \text{Spec } R[1/p, 1/g] \times (\text{Spf } \mathcal{O}[\underline{X}])^{\text{rig}}$.

Now let $x \in \text{Spec } R[1/p]$ be a closed point with $g(x) = 0$, and let $\{x_i\}_i$ be a sequence of closed points of $\text{Spec } R[1/p, 1/g]$ converging to x . Then for any $x' \in \text{Spf } \mathcal{O}[\underline{X}]^{\text{rig}}$, we have shown that $|f(x_i, x')| \leq C$. But this implies that $|f(x, x')| \leq C$, so $\|f\|_{\rho} \leq C$ for all $\rho < 1$ and we are done. \square

APPENDIX B. DETERMINANT FUNCTORS

In this appendix we recall some details of the formalism of determinant functors introduced by Fukaya and Kato in [FK06], see also [Del87].

We fix an associative unital noetherian ring R . We write $B(R)$ for the category of bounded complexes of (left) R -modules, $C(R)$ for the category of bounded complexes of finitely generated (left) R -modules, $P(R)$ for the category of finitely generated projective (left) R -modules, $C^{\text{P}}(R)$ for the category of bounded (cohomological) complexes of finitely generated projective (left) R -modules. By $D^{\text{P}}(R)$ we denote the category of perfect complexes as full triangulated subcategory of the derived category $D^{\text{b}}(R)$ of $B(R)$. We write $(C^{\text{P}}(R), \text{quasi})$ and $(D^{\text{P}}(R), \text{is})$ for the subcategory of quasi-isomorphisms of $C^{\text{P}}(R)$ and isomorphisms of $D^{\text{P}}(R)$, respectively.

For each complex $C = (C^{\bullet}, d_C^{\bullet})$ and each integer r we define the r -fold shift $C[r]$ of C by setting $C[r]^i = C^{i+r}$ and $d_{C[r]}^i = (-1)^r d_C^{i+r}$ for each integer i .

We first recall that there exists a Picard category \mathcal{C}_R and a determinant functor $\text{Det}_R : (C^{\text{P}}(R), \text{quasi}) \rightarrow \mathcal{C}_R$ with the following properties (for objects C, C' and C'' of $C^{\text{P}}(R)$)

n-fiber-desc

localize-units

determinants

- B.a) \mathcal{C}_R has an associative and commutative product structure $(M, N) \mapsto M \cdot N$ (which we often write more simply as MN) with canonical unit object $\mathbf{1}_R = \text{Det}_R(0)$. If P is any object of $P(R)$, then in \mathcal{C}_R the object $\text{Det}_R(P)$ has a canonical inverse $\text{Det}_R(P)^{-1}$. Every object of \mathcal{C}_R is of the form $\text{Det}_R(P) \cdot \text{Det}_R(Q)^{-1}$ for suitable objects P and Q of $P(R)$.
- B.b) All morphisms in \mathcal{C}_R are isomorphisms and elements of the form $\text{Det}_R(P)$ and $\text{Det}_R(Q)$ are isomorphic in \mathcal{C}_R if and only if P and Q correspond to the same element of the Grothendieck group $K_0(R)$. There is a natural identification $\text{Aut}_{\mathcal{C}_R}(\mathbf{1}_R) \cong K_1(R)$ and if $\text{Mor}_{\mathcal{C}_R}(M, N)$ is non-empty, then it is a $K_1(R)$ -torsor where each element α of $K_1(R) \cong \text{Aut}_{\mathcal{C}_R}(\mathbf{1}_R)$ acts on $\phi \in \text{Mor}_{\mathcal{C}_R}(M, N)$ to give $\alpha\phi : M = \mathbf{1}_R \cdot M \xrightarrow{\alpha \cdot \phi} \mathbf{1}_R \cdot N = N$.
- B.c) Det_R preserves the product structure: specifically, for each P and Q in $P(R)$ one has $\text{Det}_R(P \oplus Q) = \text{Det}_R(P) \cdot \text{Det}_R(Q)$.
- B.d) If $C' \rightarrow C \rightarrow C''$ is a short exact sequence of complexes, then there is a canonical isomorphism $\text{Det}_R(C) \cong \text{Det}_R(C') \text{Det}_R(C'')$ in \mathcal{C}_R (which we usually take to be an identification).
- B.e) If C is acyclic, then the quasi-isomorphism $0 \rightarrow C$ induces a canonical isomorphism $\mathbf{1}_R \rightarrow \text{Det}_R(C)$.
- B.f) For any integer r one has $\text{Det}_R(C[r]) = \text{Det}_R(C)^{(-1)^r}$.
- B.g) the functor Det_R factorises over the image of $C^p(R)$ in $D^p(R)$ and extends (uniquely up to unique isomorphisms) to $(D^p(R), \text{is})$. Moreover, if R is regular, also property B.d) extends to all distinguished triangles.
- B.h) For each C in $D^b(R)$ we write $H(C)$ for the complex which has $H(C)^i = H^i(C)$ in each degree i and in which all differentials are 0. If $H(C)$ belongs to $D^p(R)$ (in which case one says that C is *cohomologically perfect*), then C belongs to $D^p(R)$ and there are canonical isomorphisms

$$\text{Det}_R(C) \cong \text{Det}_R(H(C)) \cong \prod_{i \in \mathbf{Z}} \text{Det}_R(H^i(C))^{(-1)^i}.$$

(For an explicit description of the first isomorphism see [KM76, §3].)

- B.i) If R' is another (associative unital noetherian) ring and Y an (R', R) -bimodule that is both finitely generated and projective as an R' -module, then the functor $Y \otimes_R - : P(R) \rightarrow P(R')$ extends to a commutative diagram

$$\begin{array}{ccc} (D^p(R), \text{is}) & \xrightarrow{\text{Det}_R} & \mathcal{C}_R \\ Y \otimes_R^L - \downarrow & & \downarrow Y \otimes_R - \\ (D^p(R'), \text{is}) & \xrightarrow{\text{Det}_{R'}} & \mathcal{C}_{R'} \end{array}$$

In particular, if $R \rightarrow R'$ is a ring homomorphism and C is in $D^p(R)$, then we often simply write $\text{Det}_R(C)_{R'}$ in place of $R' \otimes_R \text{Det}_R(C)$.

- Aj) B.j) Let R° be the opposite ring of R . Then the functor $\text{Hom}_R(-, R)$ induces an anti-equivalence between \mathcal{C}_R and \mathcal{C}_{R° with quasi-inverse induced by $\text{Hom}_{R^\circ}(-, R^\circ)$; both functors will be denoted by $-^*$. This extends to give a diagram

$$\begin{array}{ccc} (D^p(R), \text{is}) & \xrightarrow{\text{Det}_R} & \mathcal{C}_R \\ \text{RHom}_R(-, R) \downarrow & & \downarrow -^* \\ (D^p(R^\circ), \text{is}) & \xrightarrow{\text{Det}_{R^\circ}} & \mathcal{C}_{R^\circ} \end{array}$$

which commutes (up to unique isomorphism); similarly we have such a commutative diagram for $\text{RHom}_{R^\circ}(-, R^\circ)$.

If R denotes any commutative ring, then by [FK06, §1.2.4 (3)] the Fukaya-Kato/Deligne determinant functor is closely related to Knudsen's and Mumford's which takes values in the category of line bundles plus ranks over $\text{Spec}(R)$. In case $SK_1(R) = 1$ both determinants can be naturally identified. Since in this article we often use geometric (specialisation) arguments we thus take frequently this point of view without indicating this in the notation.

Remark B.1. Recall from [Wei13, III.Lem. 1.4] that for a commutative semi-local ring R we have $SK_1(R) = 1$ and $K_1(R) \cong R^\times$. This applies in particular to $\Lambda_R(G)$ and $\Lambda_{\bar{R}}(G)$ as defined in (11).

We end this section by considering the example where $R = K$ is a field and V a finite dimensional vector space over K . Then, according to [FK06, 1.2.4], $\text{Det}_K(V)$ can be identified with the highest exterior product $\bigwedge^{\text{top}} V$ of V and for an automorphism $\phi : V \rightarrow V$ the determinant $\text{Det}_K(\phi) \in K^\times = K_1(K)$ can be identified with the usual determinant $\det_K(\phi)$. In particular, we identify $\text{Det}_K = K$ with canonical basis 1. Then a map $\mathbf{1}_K \xrightarrow{\psi} \mathbf{1}_K$ corresponds uniquely to the value $\psi(1) \in K^\times$.

Example B.2. Note that every finite \mathbf{Z}_p -module A possesses a free resolution C , i.e. $\text{Det}_{\mathbf{Z}_p}(A) \cong \text{Det}_{\mathbf{Z}_p}(C)^{-1} = \mathbf{1}_{\mathbf{Z}_p}$. Then modulo \mathbf{Z}_p^\times the composite $\mathbf{1}_{\mathbf{Q}_p} \xrightarrow{\text{acyc}} \text{Det}_{\mathbf{Z}_p}(C)_{\mathbf{Q}_p} \xrightarrow{\sim} \mathbf{1}_{\mathbf{Q}_p}$ corresponds to the cardinality $|A|^{-1} \in \mathbf{Q}_p^\times$.

REFERENCES

- [BB08] Denis Benois and Laurent Berger, *Théorie d'Iwasawa des représentations cristallines. II*, Comment. Math. Helv. **83** (2008), no. 3, 603–677. MR 2410782 [1](#)
- [BC08] Laurent Berger and Pierre Colmez, *Familles de représentations de de Rham et monodromie p -adique*, Astérisque (2008), no. 319, 303–337, Représentations p -adiques de groupes p -adiques. I. Représentations galoisiennes et (ϕ, Γ) -modules. [2](#)
- [Bel15] Rebecca Bellovin, *p -adic Hodge theory in rigid-analytic families*, Algebra & Number Theory **9** (2015), 371–433. [2](#), [6](#), [11](#), [17](#), [31](#), [39](#)
- [Ber02] Laurent Berger, *Représentations p -adiques et équations différentielles*, Invent. Math. **148** (2002), 219–284. [4](#), [17](#)
- [Ber04] ———, *Limites de représentations cristallines*, Compos. Math. **140** (2004), no. 6, 1473–1498. MR 2098398 [1](#), [3](#), [7](#), [10](#), [11](#), [15](#), [17](#), [18](#), [19](#), [20](#), [31](#)
- [BGR84] S. Bosch, U. Güntzer, and R. Remmert, *Non-Archimedean analysis*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 261, Springer-Verlag, Berlin, 1984, A systematic approach to rigid analytic geometry. MR 746961 [27](#)
- [CHT08] Laurent Clozel, Michael Harris, and Richard Taylor, *Automorphy for some l -adic lifts of automorphic mod l Galois representations*, Publ. Math. Inst. Hautes Études Sci. (2008), no. 108, 1–181, With Appendix A, summarizing unpublished work of Russ Mann, and Appendix B by Marie-France Vignéras. [37](#)
- [Col08] Pierre Colmez, *Espaces vectoriels de dimension finie et représentations de de Rham*, Astérisque (2008), no. 319, 117–186, Représentations p -adiques de groupes p -adiques. I. Représentations galoisiennes et (ϕ, Γ) -modules. MR 2493217 [4](#)
- [Dee01] Jonathan Dee, *Φ - Γ modules for families of Galois representations*, J. Algebra **235** (2001), no. 2, 636–664. MR 1805474 [2](#), [7](#)
- [Del87] Pierre Deligne, *Le déterminant de la cohomologie*, Current trends in arithmetical algebraic geometry (Arcata, Calif., 1985), Contemp. Math., vol. 67, Amer. Math. Soc., Providence, RI, 1987, pp. 93–177. [41](#)
- [dJ95] A.J. de Jong, *Crystalline dieudonné theory via formal and rigid geometry*, Inst. Hautes Études Sci. Publ. Math. (1995), no. 82, 5–96 (1996). MR 1383213 (97f:14047) [19](#), [30](#), [39](#)
- [FK06] Takako Fukaya and Kazuya Kato, *A formulation of conjectures on p -adic zeta functions in noncommutative Iwasawa theory*, Proceedings of the St. Petersburg Mathematical Society. Vol. XII (Providence, RI), Amer. Math. Soc. Transl. Ser. 2, vol. 219, Amer. Math. Soc., 2006, pp. 1–85. [1](#), [2](#), [22](#), [34](#), [35](#), [36](#), [41](#), [42](#), [43](#)
- [Fon90] Jean-Marc Fontaine, *Représentations p -adiques des corps locaux. I*, The Grothendieck Festschrift, Vol. II, Progr. Math., vol. 87, Birkhäuser Boston, Boston, MA, 1990, pp. 249–309. MR 1106901 (92i:11125) [5](#)
- [Her98] Laurent Herr, *Sur la cohomologie galoisienne des corps p -adiques*, Bull. Soc. Math. France **126** (1998), no. 4, 563–600. MR 1693457 [21](#)
- [Kie67] Reinhardt Kiehl, *Theorem A und Theorem B in der nichtarchimedischen Funktionentheorie*, Invent. Math. **2** (1967), 256–273. MR 0210949 (35 #1834) [38](#)
- [Kis08] Mark Kisin, *Potentially semi-stable deformation rings*, J. Amer. Math. Soc. **21** (2008), no. 2, 513–546. MR 2373358 [2](#), [8](#), [11](#)
- [Kis09] ———, *Moduli of finite flat group schemes, and modularity*, Ann. of Math. (2) **170** (2009), no. 3, 1085–1180. MR 2600871 [1](#), [8](#)
- [KL10] Kiran Kedlaya and Ruochuan Liu, *On families of ϕ, Γ -modules*, Algebra Number Theory **4** (2010), no. 7, 943–967. MR 2776879 [2](#), [38](#)
- [KM76] F. F. Knudsen and D. Mumford, *The projectivity of the moduli space of stable curves. I. Preliminaries on “det” and “Div”*, Math. Scand. **39** (1976), no. 1, 19–55. [42](#)
- [LVZ15] David Loeffler, Otmar Venjakob, and Sarah Livia Zerbes, *Local epsilon isomorphisms*, Kyoto J. Math. **55** (2015), no. 1, 63–127. MR 3323528 [1](#), [2](#), [3](#), [27](#), [28](#), [29](#), [30](#), [31](#), [32](#), [35](#)
- [LZ14] David Loeffler and Sarah Livia Zerbes, *Iwasawa theory and p -adic L -functions over \mathbf{Z}_p^2 -extensions*, Int. J. Number Theory **10** (2014), no. 8, 2045–2095. MR 3273476 [3](#), [23](#), [24](#)
- [Mat89] Hideyuki Matsumura, *Commutative ring theory*, second ed., Cambridge Studies in Advanced Mathematics, vol. 8, Cambridge University Press, Cambridge, 1989, Translated from the Japanese by M. Reid. MR 1011461 (90i:13001) [5](#), [14](#), [16](#)
- [Nak13] K. Nakamura, *A generalization of Kato’s local epsilon-conjecture for (φ, Γ) -modules over the Robba ring*, ArXiv e-prints (2013). [1](#), [2](#), [38](#)
- [Nak14] K. Nakamura, *Iwasawa theory of de Rham (φ, Γ) -modules over the Robba ring*, J. Inst. Math. Jussieu **13** (2014), no. 1, 65–118. [1](#)

- [Nek06] Jan Nekovář, *Selmer complexes*, Astérisque (2006), no. 310, viii+559. MR 2333680 [26](#), [34](#)
- [SRV13] Peter Schneider, Sujatha Ramdorai, and Otmar Venjakob, *Discussions among Otmar, Peter, and Sujatha on the paper “wach modules and iwasawa theory for modular forms” by lei/loeffler/zerbes (february 2013, ubc)*, 2013. [23](#)
- [ST03] Peter Schneider and Jeremy Teitelbaum, *Algebras of p -adic distributions and admissible representations*, Invent. Math. **153** (2003), no. 1, 145–196. MR 1990669 (2004g:22015) [38](#)
- [SU14] C. Skinner and E. Urban, *The Iwasawa main conjectures for GL_2* , Invent. Math. **195** (2014), no. 1, 1–277. [3](#)
- [Ven07] Otmar Venjakob, *From the Birch and Swinnerton-Dyer conjecture to non-commutative Iwasawa theory via the equivariant Tamagawa number conjecture—a survey*, *L-functions and Galois representations*, London Math. Soc. Lecture Note Ser., vol. 320, Cambridge Univ. Press, Cambridge, 2007, pp. 333–380. [2](#)
- [Ven13] ———, *On Kato’s local ϵ -isomorphism conjecture for rank-one Iwasawa modules*, Algebra Number Theory **7** (2013), no. 10, 2369–2416. MR 3194646 [1](#), [2](#), [29](#)
- [Wan15] Carl Wang Erickson, *Algebraic families of Galois representations and potentially semi-stable pseudodeformation rings*, preprint, 2015. [3](#)
- [Wei13] C. A. Weibel, *The K-book*, Graduate Studies in Mathematics, vol. 145, American Mathematical Society, Providence, RI, 2013, An introduction to algebraic K -theory. [42](#)

(Bellovin) 180 QUEEN’S GATE, IMPERIAL COLLEGE LONDON, SW7 2AZ

E-mail address: r.bellovin@imperial.ac.uk

URL: <http://wwwf.imperial.ac.uk/~rbellovi/>

(Venjakob) MATHEMATISCHES INSTITUT, UNIVERSITÄT HEIDELBERG, IM NEUENHEIMER FELD 288, 69120 HEIDELBERG, GERMANY

E-mail address: otmar@mathi.uni-heidelberg.de

URL: <http://www.mathi.uni-heidelberg.de/~otmar/>